


# COLORECTAL CANCER IDAHO 2018

Colorectal cancer is **preventable** through screening and early intervention.

**Idaho Goal:** 80% screening rate by 2020

**Screening 2016**



Adults that are current on screening

<b>Idaho</b>	<b>U.S.</b>
<b>62%</b>	<b>68%</b>

Public Health Districts 3, 5 and 6 have a lower screening rate than the state overall.

Idaho ranks

**44th**

in the nation for **screening**  
(Includes DC & Puerto Rico)

**646**

**new cases** of colorectal cancer in 2016


**256**

**deaths** from colorectal cancer in 2017

**56%**

of Idahoans will **survive** at least 5 years after diagnosis

Colorectal cancer screening increased from 2014 to 2016



## Screening Recommendation

Begin screening at age 50 and continue until age 75.

### Most Common Screening Options

- Colonoscopy: at least every 10 years
- Stool-based tests (gFOBT and FIT): annually

Patients at high risk for colorectal cancer should be referred directly for a colonoscopy

An abnormal stool-based test should always be followed-up with a colonoscopy

### Priority Populations

- Greatest impact on screening rates can be made by focusing on:
  - Ages 50-64
  - Adults with health insurance
  - Hispanic and American Indian/Alaska Native populations

For more information and sources visit: [www.cancer.dhw.idaho.gov](http://www.cancer.dhw.idaho.gov) or [www.ccaidaho.org](http://www.ccaidaho.org)

