

References

Breast, Cervical and Colorectal

1. Idaho Behavioral Risk Factor Surveillance System (BRFSS), 2014, and CDC BRFSS public use dataset, 2014. Boise, ID: Idaho Department of Health and Welfare, Division of Public Health, Bureau of Vital Records and Health Statistics; June 2016.
2. Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence - SEER 18 Regs Research Data + Hurricane Katrina Impacted Louisiana Cases, Nov 2015 Sub (2000-2013) <Katrina/Rita Population Adjustment> - Linked To County Attributes - Total U.S., 1969-2014 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2016, based on the November 2015 submission.
3. Johnson, C.J., Carson, S.L., *Cancer in Idaho, 2013*. Boise, ID: Cancer Data Registry of Idaho; December 2015.
4. Johnson, C.J., *Unpublished*, Cancer Data Registry of Idaho; April 2016.
5. Comprehensive Cancer Alliance of Idaho, *Idaho Comprehensive Cancer Strategic Plan 20016-2020*. Boise, ID: Idaho Department of Health and Welfare, Bureau of Community and Environmental Health; July 2016.

Notes

- Incidence and mortality rates are per 100,000 and age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) standard.
- U.S. incidence rates are White only because data are more comparable to the population demographics of Idaho than the U.S. All Races data.
- Screening rates for breast, cervical and colorectal cancers are measured through the Behavioral Risk Factor Surveillance System (BRFSS). Each specific measure is defined below.
- Idaho goals are set by the Comprehensive Cancer Alliance of Idaho (CCAI).
- Survival estimates are based on crude survival and include both cancer as well as other competing causes of death.

Breast Cancer

- BRFSS breast cancer screening measure is defined as percent of females aged 50-74 who had a mammogram in the past two years.
- BRFSS data are age-adjusted to 2000 US Standard Population age groups, 50-64 and 65-74.

Idaho Cancer Fact Sheets

- Screening recommendations vary. The U.S. Preventive Services Task Force recommendation summary and supplemental information can be found here:
<http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/breast-cancer-screening1?ds=1&s=breast%20cancer>
- Late stage incidence is included because the CCAI goal is specific to that measure.

Cervical Cancer

- Idaho ranks 51st because all states and the District of Columbia are included.
- Public Health District data are not available for mortality due to a small population size.
- BRFSS cervical cancer screening measure is defined as percent of females aged 21-65 who had a pap test in the past three years.
- BRFSS data are age-adjusted to 2000 US Standard Population age groups 21-34, 35-44, and 45-65.
- The U.S. Preventive Services Task Force screening recommendation summary and supplemental information can be found here:
<http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/cervical-cancer-screening>

Colorectal Cancer

- BRFSS colorectal cancer screening measure is defined as percentage of adults aged 50-75 who received a blood stool test in the past year, sigmoidoscopy in the past five years and blood stool test in the past three years, or a colonoscopy in the past 10 years.
- BRFSS data are age-adjusted to 2000 US Standard Population age groups, 50-64 and 65-75.
- Screening recommendation and screening options by the U.S. Preventive Services Task Force were updated in June 2016. Recommendation summary and supplemental information can be found here:
<http://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/colorectal-cancer-screening2>