


# IDAHO Colorectal CANCER 2016

Colorectal cancer is **preventable** through screening.

**Idaho Goal:** 80% screening rate by 2020

**Screening 2014**



Adults that are up to date on screening.

<b>Idaho</b>	<b>U.S.</b>
<b>61%</b>	<b>64%</b>

Public Health Districts 1, 3, 5 and 6 have lower screening rates than the state overall

Idaho ranks **44th** in the nation for screening

**604** new cases of colorectal cancer in 2013

**232** deaths from colorectal cancer in 2014

**55%** of Idahoans will survive at least 5 years after diagnosis

## Screening Recommendation


Begin screening at age 50 and continue until age 75.

### Screening Options

- **Stool-based tests**  
*gFOBt, FIT, and FIT-DNA*
- **Direct visualization tests**  
*Flexible sigmoidoscopy, alone or combined with FIT; colonoscopy; and CT colonography*
- **Serology tests**  
*SEPT9 DNA test*


### Take Action

- Know and use screening guidelines
- Provide patients with screening options
- Implement provider and patient reminders for timely screening
- Identify system improvements to streamline screening processes



### Incidence 2013

Public Health Districts 1 and 3 have higher incidence rates than the state and national overall rates.



### Mortality 2014

Public Health Districts 1, 2, 3, 5 and 6 have death rates from colorectal cancer that are higher than the state overall rate.

For more information and sources visit: [www.cancer.dhw.idaho.gov](http://www.cancer.dhw.idaho.gov) or [www.ccaidaho.org](http://www.ccaidaho.org)

