


Sexually Transmitted Diseases Reported Cases Among Residents of Idaho January - March 2013

| District | HIV Infection¶ | Chlamydia | Gonorrhea | Early Syphilis* | Congenital Syphilis | Late or Unknown Duration Syphilis** | TOTAL |
|----------|-------------------|-----------|-----------|--------------------|------------------------|--|-------|
| 1 | 0 | 192 | 1 | 0 | 0 | 1 | 194 |
| 2 | 0 | 75 | 3 | 0 | 0 | 0 | 78 |
| 3 | 2 | 308 | 2 | 3 | 1 | 2 | 318 |
| 4 | 3 | 417 | 10 | 1 | 0 | 3 | 434 |
| 5 | 0 | 125 | 1 | 0 | 0 | 0 | 126 |
| 6 | 0 | 81 | 0 | 1 | 0 | 1 | 83 |
| 7 | 0 | 60 | 1 | 0 | 0 | 0 | 61 |
| Total | 5 | 1,258 | 18 | 5 | 1 | 7 | 1,294 |

Sexually Transmitted Diseases Reported Cases Among Residents of Idaho January – March 2012

| District | HIV Infection¶ | Chlamydia | Gonorrhea | Early Syphilis* | Congenital Syphilis | Late or Unknown Duration Syphilis** | TOTAL |
|----------|-------------------|-----------|-----------|--------------------|------------------------|--|-------|
| 1 | 0 | 169 | 3 | 0 | 0 | 0 | 172 |
| 2 | 0 | 95 | 7 | 0 | 0 | 0 | 102 |
| 3 | 1 | 192 | 4 | 0 | 0 | 0 | 197 |
| 4 | 2 | 425 | 11 | 10 | 0 | 0 | 448 |
| 5 | 1 | 111 | 0 | 0 | 0 | 1 | 113 |
| 6 | 0 | 99 | 5 | 0 | 0 | 0 | 104 |
| 7 | 0 | 74 | 0 | 0 | 0 | 0 | 74 |
| Total | 4 | 1,165 | 30 | 10 | 0 | 1 | 1,210 |



| Idaho District Health Departments | | | | | | |
|-----------------------------------|---------------|----------|--|--|--|--|
| District 1 | Coeur d'Alene | 415-5100 | | | | |
| District 2 | Lewiston | 799-3100 | | | | |
| District 3 | Caldwell | 455-5300 | | | | |
| District 4 | Boise | 375-5211 | | | | |
| District 5 | Twin Falls | 734-5900 | | | | |
| District 6 | Pocatello | 233-9080 | | | | |
| District 7 | Idaho Falls | 522-0310 | | | | |

| Reported Idaho HIV and AIDS Cases Presumed Living* by Diagnostic Status as of March 31, 2013 | | | | | | | | | |
|--|----------------|-----------------------------------|--------------------------------|---------------|------------------------------------|---------------------------------|--|--|--|
| District | HIV Cases** | HIV Cases Reported Deceased | Presumed Living with HIV | AIDS Cases | AIDS Cases Reported Deceased | Presumed Living with AIDS | Reported HIV and AIDS Cases Presumed Living*** | | |
| 1 | 78 | 11 | 67 | 164 | 76 | 88 | 155 | | |
| 2 | 55 | 6 | 49 | 72 | 24 | 48 | 97 | | |
| 3 | 102 | 11 | 91 | 150 | 52 | 98 | 189 | | |
| 4 | 372 | 37 | 335 | 497 | 217 | 280 | 615 | | |
| 5 | 65 | 9 | 56 | 109 | 44 | 65 | 121 | | |
| 6 | 84 | 10 | 74 | 119 | 52 | 67 | 141 | | |
| 7 | 58 | 4 | 54 | 96 | 41 | 55 | 109 | | |
| TOTAL | 814 | 88 | 726 | 1,207 | 506 | 701 | 1,427 | | |

All reported HIV and AIDS cases in Idaho, regardless of residence at first diagnosis or death.

^{*}Reported cases of HIV not reported with AIDS EVER REPORTED in Idaho and have not been reported as deceased. Previous prevalence tables only included cases first reported in Idaho. Reported AIDS cases presumed living - Reported cases of AIDS EVER REPORTED in Idaho and have not been reported as deceased. Previous prevalence tables only included cases first reported in Idaho.

^{**}Number of reported individuals with HIV (not yet AIDS) EVER REPORTED in Idaho, excluding those who were concurrently diagnosed with AIDS or have progressed to AIDS

^{***}Reported HIV/AIDS cases presumed living - Total number of HIV and AIDS cases EVER REPORTED in Idaho that have not been reported as deceased.

| HIV and AIDS (HIV Stage 3) reports among residents of Idaho at diag | | | nosis, by case s | | status through N | | March 31, 2013 COMBINED | |
|---|----------------------------------|-----------|------------------|------|------------------|-------|----------------------------|--|
| Sex | | | V | All | | OOME | INCO | |
| Male | | 425 | 83% | 635 | 85% | 1,060 | 84% | |
| Female | | 88 | 17% | 115 | 15% | 203 | 16% | |
| Total | 513 | 100% | 750 | 100% | 1,263 | 100% | | |
| | ılt & Adolescent | | | | | 1,200 | 100,0 | |
| Adults/Adoles | scents (13 or above) | 510 | 99% | 741 | 99% | 1,251 | 99% | |
| Pediatric (<13 | , | 3 | 1% | 5 | 1% | 8 | 1% | |
| Total | | 513 | 100% | 746 | 100% | 1,259 | 100% | |
| Race/Ethnicit | ty | | | | | | | |
| Hispanic - All | Races | 60 | 12% | 83 | 11% | 143 | 11% | |
| Not Hispanic | American Indian/Alaska Native | 11 | 2% | 11 | 1% | 22 | 2% | |
| | Asian | 3 | 1% | 1 | 0% | 4 | 0% | |
| | Black | 21 | 4% | 23 | 3% | 44 | 3% | |
| | Native Hawaiian/Pacific Islander | 0 | 0% | 0 | 0% | 0 | 0% | |
| | White | 404 | 79% | 654 | 84% | 1,058 | 82% | |
| | Legacy Asian/Pacific Islander | 0 | 0% | 2 | 0% | 2 | 0% | |
| | Multi-race | 4 | 1% | 6 | 1% | 10 | 0% | |
| | Unknown | 10 | 2% | 3 | 0% | 13 | 1% | |
| Total | | 513 | 100% | 783 | 100% | 1,296 | 100% | |
| Exposure Categories - Adults/Adolescents | | | | | | · | | |
| Men who have sex with men (MSM) | | 248 | 49% | 382 | 51% | 630 | 50% | |
| Injection drug | use | 52 | 10% | 82 | 11% | 134 | 11% | |
| MSM and inje | MSM and inject drugs | | 11% | 65 | 9% | 122 | 9% | |
| Hemophilia/C | oagulation disorder | 1 | 0% | 18 | 2% | 19 | 2% | |
| High-risk hete | erosexual contact | 58 | 11% | 95 | 13% | 153 | 12% | |
| Presumed he | terosexual contact | 11 | 2% | 14 | 2% | 25 | 2% | |
| Receipt of blo | ood, components, or tissue | 2 | 0% | 11 | 1% | 13 | 1% | |
| Other/risk not | reported or identified | 81 | 16% | 78 | 10% | 159 | 13% | |
| Total | | 510 | 100% | 745 | 100% | 1,255 | 101% | |
| | tegories - Pediatric | | | | | | | |
| • | oagulation disorder | 0 | 0% | 0 | 0% | 0 | 0% | |
| | t risk for HIV infection | 2 | 67% | 2 | 40% | 4 | 50% | |
| | ood, components, or tissue | 0 | 0% | 2 | 40% | 2 | 25% | |
| | reported or identified | 1 | 33% | 1 | 20% | 2 | 25% | |
| Total | | 3 | 100% | 5 | 100% | 8 | 100% | |
| Cases by Age Group | | 3 | 4.04 | | 00/ | | 001 | |
| Under 5 | | | 1% | 2 | 0% | 5 | 0% | |
| 5-12 | | | 0% | 1 | 0% | 1 | 0% | |
| 13-19 | | | 3% | 7 | 1% | 20 | 2% | |
| 20-29 | | | 41% | 113 | 15% | 324 | 26% | |
| 30-39 40-49 | | 156 | 30% | 306 | 41% | 462 | 37% | |
| | | 99 | 19% | 216 | 29% | 315 | 25% | |
| Over 49 | | | 6% | 105 | 14% | 136 | 11% | |
| Unknown Total | | | 100% | 750 | 100% | 1,263 | 0% 100% | |
| Rate (Cumulative) per 100,000 Residents | | 513 32 | 100% | | 100% '.8 | 1,203 | 100% | |

This table excludes cases of HIV or AIDS (HIV Stage 3) first diagnosed in other states. Data are provisional. Percentages may not add up to 100% due to rounding. Reports of HIV positive and AIDS cases should be made to the Bureau of Communicable Disease Prevention and Control as required by Idaho Code 39-602. Data collection and publication is made possible through a federal grant from the Centers for Disease Control and Prevention, United States – Department of Health and Human Services. Idaho statistics are posted on the Internet four times per year. To be placed on an e-mail alert group to signal you when a new issue is posted, please e-mail Allen Haumann at haumanna@dhw.idaho.gov.