Selective Screening Criteria

Rev 1/2012

Chlamydia (CT)

All Males

Females < 25

Mucopurulence

Friability

Sex with 2 or more partners (60 days)

Ectopy with inflammation

Gonorrhea (GC)

Sex worker or partner of sex worker

Sex worker

Sex worker or partner of MSM

Cervicitis, urethritis, or PID

PID Exposed to GC
Pregnant IUD insert planned

Exposed to CT History of repeated GC infections IUD insert planned Symptomatic partner (60 days)

Positive CT last 12 months Client with HIV

Symptomatic partner (60 days) Re-screen (3-4 months after +) Re-screen (3-4 months after +) Meets any criteria for CT

Meets any criteria for GC

Herpes

Undiagnosed, painless, ulcerative oral/genital lesion

Hepatitis B (HBV)

2 or more sex partners (60 days)

Pregnant

Commercial sex worker or partner of sex worker

Sexually active MSM or partner of MSM

Client with HIV

Client Injection drug user (IDU) past or present

Sexual contact with IDU

Exposed to HBV

Household contact of person with HBV

HIV Counseling, Testing, and Referral (CTR)

MSM (Men who have sex with men)

MSM / IDU

Heterosexual IDU

Sex partner at risk (Partner IDU, MSM, or HIV+)

Child of HIV+ woman

STD or active TB diagnosis

Sex for drugs / money

Pregnant

Hepatitis C (HCV)

Exposed to HCV IDU Past or Present

Sexual Contact of IDU

Syphilis

Sex worker or partner of sex worker

Sex for money/drugs

Sexually active MSM or partner of MSM

F/U post treatment

Pregnant

Symptomatic partner (60 days)

Exposed to syphilis
Person with HIV or GC

Undiagnose painless ulcerative genital -

or oral lesions

Unexplained adult rash

Unexplained lymphadenopathy

Suspicious condylomata lata