



IDAHO DEPARTMENT OF HEALTH & WELFARE
DIVISION OF PUBLIC HEALTH

Influenza Surveillance 2015-2016 Season Update


Week ending: **11/14/2015** CDC MMWR week: **45**

Synopsis

- Only **0.60%** of visits to ILINet-participating healthcare providers in Idaho were for influenza-like illness (ILI) this week. This frequency of visits for ILI is considered low.
- **2** specimens were tested by the Idaho Bureau of Laboratories (IBL) this week for influenza; **0** were positive.
- **0** influenza-associated deaths were reported this week.
- Several Idaho public health districts have reported receiving anecdotal reports of positive rapid influenza test results from local healthcare providers during this week.
- State activity code: **Sporadic**

Outpatient Surveillance Data

Data on outpatient visits to health care providers for ILI are collected through the Centers for Disease Control and Prevention's (CDC) U.S. Outpatient ILI Surveillance Network (ILINet). ILI is defined as fever (temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat without a KNOWN cause other than influenza.



During MMWR week **45** the percentage of visits for ILI in Idaho remains very low, at **0.60%**. This frequency is slightly higher than the % ILI visits recorded during week **44** (**0.43%**).

A small number of healthcare sites from every state, including Idaho, provide weekly situational awareness on the geographic distribution and current level of ILI activity in their area. Data collected include the total number of patients seen for any reason, and the number of those patients specifically seen for ILI. Information is reported by age group (0–4 years, 5–24 years, 25–49 years, 50–64 years, and ≥ 65 years) to understand the population most affected. The percentage of weekly visits to Idaho ILINet providers for ILI during the 2015–2016 influenza season (all ages) are compared with weekly visits during the 2012–2013, 2013–2014, and 2014–2015 influenza seasons in Figure 1.

Visit <http://www.cdc.gov/flu/weekly/overview.htm> to learn more about ILINet

Laboratory Data

- During MMWR week 45 (From 11/08/2015 to 11/14/2015), IBL reported results for 2 sample; 0 samples were positive for influenza
- For the 2015-2016 influenza season (week 40 to date), IBL reported results on 7 samples tested for influenza; 1 sample was positive for influenza A(H3).

Mortality Data

Data regarding influenza-associated deaths are reviewed weekly during the influenza season. The number and age groups affected provide information on populations most severely affected.

0 influenza-associated deaths were reported during MMWR week 45; 0 influenza-related deaths have been reported so far for the 2015-2016 season (see Table 1.)

Table 1. Total influenza-related deaths occurring in Idaho during the 2015-2016 influenza season, by age group, and public health district of residence (as of week ending 10/10/2015)

Residence	Total	Age		
		<18 yrs	18-49 yrs	50+ yrs
Panhandle Health District (PHD-1)	-	-	-	-
North Central Health District (PHD-2)	-	-	-	-
Southwest District Health (PHD-3)	-	-	-	-
Central District Health Department (PHD-4)	-	-	-	-
South Central Public Health (PHD-5)	-	-	-	-
Southeastern Idaho Public Health (PHD-6)	-	-	-	-
Eastern Idaho Public Health District (PHD-7)	-	-	-	-
Non-resident	-	-	-	-
Total	0	-	-	-

Source: Bureau of Vital Records and Health Statistics, Idaho Department of Health and Welfare
 Link to Idaho Public Health Districts: <http://www.healthandwelfare.idaho.gov/?TabId=97>

Only deaths occurring in Idaho are reported in Table 1; data include deaths in Idaho residents and non-residents, by public health district of residence. Deaths in Idaho

residents occurring while out-of-state are not shown here. Data shown in Table 1 may differ from statistics based on Idaho resident records, data based on underlying cause of death only, and data based on calendar year. Deaths are considered influenza-related based on ICD coding for Influenza in Part I or Part II on the death certificate. Influenza may have been the underlying cause of death or contributed to death.

Table 2 lists the number of Idaho deaths recorded during recent past influenza seasons; with an average of 22 deaths recorded annually over the previous six years.

Influenza season	Influenza-associated deaths
2014-2015	32
2013-2014	19
2012-2013	35
2011-2012	5
2010-2011	21
2009-2010	22

State Activity Code

For week 45: [Sporadic](#)

State health departments report estimated levels of geographic spread of influenza weekly to CDC, based on surveillance findings.

- **No Activity:** No lab-confirmed cases, no reported increase in ILI activity.
- **Sporadic:** Small numbers of lab-confirmed cases or a single laboratory-confirmed outbreak. No reported increase in ILI activity.
- **Local:** Outbreaks or increases in ILI activity, lab-confirmed influenza in a single region* of the state.
- **Regional:** Outbreaks of influenza or increases in ILI activity and recent lab-confirmed influenza in at least two but less than half the regions of the state.
- **Widespread:** Outbreaks or increases in ILI activity and recent lab-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

*In Idaho regions are defined by public health district boundaries.

This page, generated by the Idaho Department of Health and Welfare, Division of Public Health, Bureau of Communicable Disease Prevention, is updated regularly during the traditional influenza season. Additional postings occur during an early, late, or prolonged season.