

National Immunization Survey—Child Data for United States and Idaho, 2006 through 2013

	4:3:1:3:3:1				4:3:1:3:3:1:4				4:3:1-:3:1:4				4:3:1:3 <sup>†</sup> :3:1:4			
	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI
2006, n=222	76.9	1.0	68.2	6.9												
2007, n=183	77.4	1.1	65.7	7.2	66.5	1.3	52.9	7.6								
2008, n=282	76.1	1.1	60.4	6.8	68.4	1.2	54.2	6.7	70.6	1.2	60.4	6.6				
2009, n=207	69.9	1.2	51.7	7.7	63.6	1.3	48.0	7.7	70.5	1.2	<b>70.5</b>	7.0	44.3	1.4	33.6	7.2
2010, n=271	74.9	1.1	61.2	6.7	70.2	1.3	57.4	6.8	72.7	1.2	66.3	6.4	56.6	1.3	42.6	6.8
2011, n=317	77.0	1.1	<b>68.8</b>	7.6	73.3	1.2	66.9	7.7	73.6	1.2	<b>66.9</b>	7.7	68.5	1.3	57.1	8.4
2012, n=289	71.9	1.4	64.5	8.1									68.4	1.4	63.0	8.2
2013, n=233	77.7	1.4	76.9	6.4					72.7	1.5	76.2	6.4	70.4	1.5	70.2	6.9

	4+ DTaP/DTP/DT				3+ Polio vaccine				1+ MMR			
	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI
2006, n=222	85.2	0.9	82.4	5.8	92.8	0.6	92.2	3.9	92.3	0.6	88.0	4.8
2007, n=183	84.5	0.9	77.2	6.3	92.6	0.7	87.6	4.9	92.3	0.7	86.1	5.2
2008, n=282	84.6	1.0	77.6	5.9	93.6	0.6	91.8	3.7	92.1	0.7	86.1	5.4
2009, n=207	83.9	1.0	84.0	5.5	92.8	0.7	94.7	3.5	90.0	0.8	88.1	4.8
2010, n=271	84.4	1.0	77.9	5.7	93.3	0.7	89.2	4.7	91.5	0.7	87.2	5.0
2011, n=317	84.6	1.0	79.0	6.6	93.9	0.6	94.5	3.1	91.6	0.8	89.5	4.8
2012, n=289	82.5	1.2	76.6	6.7	92.8	0.7	92.6	3.9	90.8	0.8	93.3	3.6
2013, n=233	83.1	1.3	84.2	5.3	92.7	1.0	92.6	3.7	91.9	0.9	91.1	4.3

	HIB-FS				3+ Hib vaccine				HIB-PS			
	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI
2006, n=222					93.4	0.6	91.3	4.2				
2007, n=183					92.6	0.7	86.3	5.1				
2008, n=282					90.9	0.7	<b>77.6</b>	3.4				
2009, n=207					86.6	1.0	<b>64.1</b>	7.5	92.1	0.8	90.9	4.5
2010, n=271	66.8	1.3	53.3	6.9	90.4	0.9	<b>77.7</b>	6.2	92.2	0.8	<b>82.3</b>	5.9
2011, n=317	80.4	1.1	75.0	7.3	94.0	0.6	<b>94.7</b>	3.1	94.2	0.6	<b>94.9</b>	3.1
2012, n=289	80.9	1.2	77.3	7.6					93.3	0.7	92.7	3.9
2013, n=233	82.0	1.3	80.0	5.9	92.8	0.9	91.2	4.3	93.7	0.9	94.7	3.1

	1+ VAR				4+ PCV				2+ Hep A vaccine			
	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI
2006, n=222	89.2	0.7	77.4	6.1	68.4	1.1	72.1	6.5				
2007, n=183	90.0	0.7	75.5	6.4	75.3	1.2	66.6	7.2				
2008, n=282	90.7	0.7	80.7	5.8	80.1	1.1	74.8	6.3	40.4	1.2	38.6	6.5
2009, n=207	89.6	0.8	84.1	5.7	80.4	1.1	82.5	6.0	46.6	1.4	37.3	7.5
2010, n=271	90.4	0.8	83.7	5.1	83.3	1.0	79.7	5.5	49.7	1.4	44.2	6.8
2011, n=317	90.8	0.7	86.0	5.6	84.4	1.0	84.1	5.7	52.2	1.4	45.2	8.6
2012, n=289	90.2	0.1	90.3	4.3	81.9	1.1	83.9	4.8	53.0	1.1	52.8	8.6
2013, n=233	91.2	0.9	87.9	4.8	82.0	1.3	87.3	4.6	54.7	1.6	60.7	7.3


	Rotavirus				Hep B vaccine birth dose				3+ Hep B vaccine			
	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI	U.S.	± CI	Idaho	± CI
2006, n=222									93.3	0.6	92.9	3.5
2007, n=183									92.7	0.7	88.8	4.7
2008, n=282					55.3	1.3	64.0	6.6	93.5	0.7	93.3	3.4
2009, n=207	43.9	1.4	30.3	7.2	60.8	1.3	70.3	7.1	92.4	0.7	94.3	3.6
2010, n=271	59.2	1.4	<b>50.4</b>	6.9	64.1	1.3	65.2	6.4	91.8	0.7	<b>86.8</b>	4.9
2011, n=317	67.3	1.3	<b>62.0</b>	8.2	68.6	1.3	70.2	7.5	91.1	0.7	86.3	5.4
2012, n=289	68.6	1.4	68.2	7.2	71.6	1.4	70.1	7.8	89.0	0.9	87.7	5.0
2013, n=233	72.6	1.5	74.6	6.4	74.2	1.4	72.7	6.5	90.8	1.0	90.7	4.3

**Glossary of Abbreviations and Terms**

- 4:3:1:3:3:1 The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MMR vaccinations, 3 or more Hib vaccinations of any type, 3 or more hepatitis B vaccinations, and 1 or more varicella vaccinations given at age 12 months or older
- 4:3:1:3<sup>†</sup>:3:1:3 The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MMR vaccinations, 2 or 3 Hib vaccinations depending on brand received, 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 4 or more pneumococcal vaccinations
- 4:3:1:3:3:1:4 The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MMR vaccinations, 3 or more Hib vaccinations of any type, 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 4 or more pneumococcal vaccinations
- 4:3:1-:3:1:4 The series of 4 or more DTaP vaccinations, 3 or more polio vaccinations, 1 or more MMR vaccinations, 3 or more hepatitis B vaccinations, 1 or more varicella vaccinations given at age 12 months or older, and 4 or more pneumococcal vaccinations
- DTaP Diphtheria and tetanus toxoids and acellular pertussis vaccine
- DTP Diphtheria and tetanus toxoids and pertussis vaccine
- DT Diphtheria and tetanus toxoids vaccine
- FS Full series
- Hep A Hepatitis A
- Hep B Hepatitis B
- Hib *Haemophilus influenzae* type b
- MMR Measles, mumps, and rubella vaccine
- NIS National Immunization Survey
- PCV Pneumococcal conjugate vaccine
- PS Primary series
- VAR Varicella vaccine

<sup>†</sup> Full series *Haemophilus influenzae* type b (Hib) vaccine: ≥3 or ≥4 doses, depending on brand received  
 Idaho data points in **BOLD** indicate a statistically significant change compared to the previous year, p=0.05  
 Blue cells indicate NIS did not collect data during that year  
 CI = Confidence Interval

## 4:3:1:3:3:1\* Vaccination Coverage, 2006–2013†



\* Refers to 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids (DTP/DT/DTaP), 3 or more doses of polio virus vaccine, 1 or more doses a measles-mumps-rubella (MMR) vaccine, 3 or more doses of *Haemophilus influenzae* type b (Hib) vaccine of any type, 3 or more doses of hepatitis B vaccine, and 1 or more doses of varicella vaccine


† From the National Immunization Survey (NIS), children aged 19 through 35 months

‡ Refers to a statistically significant change compared to the previous year,  $p=0.05$

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## 4:3:1:3:3:1:4\* Vaccination Coverage, 2007–2011<sup>†§</sup>




\* Refers to 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids (DTP/DT/DTaP), 3 or more doses of polio virus vaccine, 1 or more doses of measles-mumps-rubella vaccine (MMR), 3 or more doses of *Haemophilus influenzae* type b (Hib) vaccine of any type, 3 or more doses of hepatitis B vaccine, 1 or more doses of varicella vaccine, and 4 or more doses of pneumococcal conjugate vaccine (PCV)  
<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months

<sup>§</sup> This series was sampled during 2007 through 2011

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## 4:3:1:-:3:1:4\* Vaccination Coverage, 2008–2013<sup>†§</sup>



\* Refers to 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids (DTP/DT/DTaP), 3 or more doses of polio virus vaccine, 1 or more doses of measles-mumps-rubella vaccine (MMR), *Haemophilus influenzae* type b (Hib) vaccine is excluded, 3 or more doses of hepatitis B vaccine, 1 or more doses of varicella vaccine, and 4 or more doses of pneumococcal conjugate vaccine (PCV)

<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months


<sup>§</sup> This series was sampled during 2008 through 2011 to exclude the impact of the national Hib vaccine shortage and not sampled during 2012

¥ Refers to a statistically significant change compared to the previous year, p=0.05

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## 4:3:1:3<sup>‡</sup>3:1:4\* Vaccination Coverage, 2009–2013<sup>†§</sup>



\*Refers to 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids (DTP/DT/DTaP), 3 or more doses of polio virus vaccine, 1 or more doses of a measles-mumps-rubella vaccine (MMR), plus full series of *Haemophilus influenzae* type b (Hib), 3 or more doses of hepatitis B vaccine, 1 or more doses of varicella vaccine, and 4 or more doses of pneumococcal conjugate vaccine (PCV)

<sup>‡</sup> Full series *Haemophilus influenzae* type b (Hib) vaccine:  $\geq 3$  or  $\geq 4$  doses, depending on brand received


<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months

<sup>§</sup> This series was sampled beginning in 2012, however in 2013 data was published for 2009–2011 rates

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

# 4+ DTaP\* Vaccination Coverage, 2006–2013†




\* Refers to 4 or more doses of diphtheria and tetanus toxoids and pertussis vaccine, or diphtheria and tetanus toxoids (DTP/DT/DTaP)

† From the National Immunization Survey (NIS), children aged 19 through 35 months

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

### 3+ Polio\* Vaccination Coverage, 2006–2013†




\* Refers to 3 or more doses of any polio virus vaccine

† From the National Immunization Survey (NIS), children aged 19 through 35 months

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

# 1+ MMR\* Vaccination Coverage, 2006–2013†




\* Refers to 1 or more doses of a measles, mumps, and rubella vaccine (MMR)

† From the National Immunization Survey (NIS), children aged 19 through 35 months

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## Hib-FS\* Vaccination Coverage, 2010–2013<sup>†§</sup>



\* Refers to full series *Haemophilus influenzae* type b (Hib) vaccine:  $\geq 3$  or  $\geq 4$  doses, depending on brand received


<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months

<sup>§</sup> This series was sampled beginning in 2010

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

### 3+ Hib\* Vaccination Coverage, 2006–2013<sup>†§</sup>



\* Refers to 3 or more doses of *Haemophilus influenzae* type b (Hib) vaccine

<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months


<sup>§</sup> This series was sampled during 2006 through 2013 except 2012

<sup>¥</sup> Refers to a statistically significant change compared to the previous year, p=0.05

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## Hib\* -PS Vaccination Coverage, 2009–2013<sup>†§</sup>



\* Refers to primary series *Haemophilus influenzae* type b (Hib) vaccine:  $\geq 2$  or  $\geq 3$  doses, depending on brand received

<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months


<sup>§</sup> This series was sampled beginning in 2009

¥ Refers to a statistically significant change compared to the previous year,  $p=0.05$

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## 1+ VAR\* Vaccination Coverage, 2006–2013†




\* Refers to 1 or more doses of varicella vaccine at or after child's first birthday, unadjusted for history of varicella illness

† From the National Immunization Survey (NIS), children aged 19 through 35 months

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

### 4+ PCV\* Vaccination Coverage, 2006–2013†




\* Refers to 4 or more doses of pneumococcal conjugate vaccine (PCV)

† From the National Immunization Survey (NIS), children aged 19 through 35 months

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## 2+ Hep A\* Vaccination Coverage, 2008–2013<sup>†§</sup>



\* Refers to 2 or more doses of hepatitis A vaccine


<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months

<sup>§</sup> This series was sampled beginning in 2008

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## Rotavirus\* Vaccination Coverage, 2009–2013<sup>†§</sup>



\* Refers to  $\geq 2$  or  $\geq 3$  doses of rotavirus vaccine depending on brand received

<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months


<sup>§</sup> This series was sampled beginning in 2009

<sup>¥</sup> Refers to a statistically significant change compared to the previous year,  $p=0.05$

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

## Hep B Birth Dose\* Vaccination Coverage, 2008–2013<sup>†§</sup>



\* Refers to 1 or more doses of hepatitis B vaccine administered between birth and age 3 days


<sup>†</sup> From the National Immunization Survey (NIS), children aged 19 through 35 months

<sup>§</sup> This series was sampled beginning in 2008

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval

### 3+ Hep B\* Vaccination Coverage, 2006–2013†



\* Refers to 3 or more doses of hepatitis B vaccine

† From the National Immunization Survey (NIS), children aged 19 through 35 months

‡ Refers to a statistically significant change compared to the previous year,  $p=0.05$

n = number of children sampled for each survey year in Idaho

Vertical bars above and below each point estimate indicate the 95% confidence interval