

# Smoking During Pregnancy

Idaho PRATS, 2010-2015\*



## Quick Facts


14.2 percent of moms smoked in the 3 months prior to pregnancy.

86.5 percent of moms who smoked in the 3 months prior to pregnancy either quit or decreased smoking levels during pregnancy.

6.3 percent of moms reported smoking in the third trimester of pregnancy.

During the aggregate period from 2010-2015, 14.2 percent of mothers smoked in the three months prior to pregnancy. The majority (58.3 percent) of those mothers who smoked in the three months prior to pregnancy quit smoking during pregnancy, and over a quarter (28.2 percent) decreased their smoking level during pregnancy.


Pregnancy smoking levels among Idaho moms who smoked prior to pregnancy, 2010-2015



Between 2010 and 2015, 6.3 percent of Idaho mothers smoked in the third trimester of pregnancy. A higher proportion of Idaho mothers who smoked in the third trimester:

- Utilized Medicaid during pregnancy
- Had annual incomes of less than \$25,000
- Participated in WIC
- Did not have a high school diploma
- Experienced an unintended pregnancy
- Were aged 25 and older
- Were non-Hispanic.

Demographic characteristics of Idaho moms who smoked in the third trimester, 2010-2015




\*Except 2012. No data were collected in 2012.

# Smoking During Pregnancy

## Idaho PRATS, 2010-2015\*

Public Health District 2 had the highest concentration of moms who smoked in the third trimester from 2010-2015 with 9.4 percent, and District 7 had the lowest concentration with 3.6 percent.



For more information please contact:  
Health Statistics Section  
Bureau of Vital Records and  
Health Statistics  
Idaho Department of Health  
and Welfare  
450 W. State Street 1st Floor  
P.O. Box 83720  
Boise, ID 83720-0036  
(208) 332-7366  
[www.prats.dhw.idaho.gov](http://www.prats.dhw.idaho.gov)

### What is PRATS?

The Pregnancy Risk Assessment Tracking System (PRATS) is an annual survey of new mothers in Idaho. It is a population-based tracking system that identifies and examines maternal experiences and health practices associated with pregnancy which may affect pregnancy outcomes and infant health. PRATS was first conducted in Idaho in 1999, and has been conducted annually since 2001\*. PRATS is modeled after the Centers for Disease Control and Prevention (CDC) Pregnancy Risk Assessment Monitoring System (PRAMS).

\*Except 2012

This project was funded by grants through the State Systems Development Initiative (SSDI) and the Title V MCH Service Block Grant Program (Maternal and Child Health Bureau, Health Resources and Services Administration, U.S. Department of Health and Human Services). This and other health statistics publications can be found online at [www.healthstatistics.dhw.idaho.gov](http://www.healthstatistics.dhw.idaho.gov).



Costs associated with this publication are available from the Idaho Department of Health and Welfare, HW-1206.

•January 2018•



IDAHO DEPARTMENT OF HEALTH & WELFARE  
DIVISION OF PUBLIC HEALTH