



## Hemophilia Agents Therapeutic Class Review (TCR)

---

September 19, 2015

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Magellan Rx Management.

All requests for permission should be mailed to:

Magellan Rx Management  
Attention: Legal Department  
6950 Columbia Gateway Drive  
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to [PSTCReDitor@magellanhealth.com](mailto:PSTCReDitor@magellanhealth.com).

---

September 2015

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.  
© 2013-2015 Magellan Rx Management. All Rights Reserved.

**MagellanRx**  
MANAGEMENT<sup>SM</sup>

## FDA-APPROVED INDICATIONS

### Factor VIII Products

Drug	Manufacturer	Indications
antihemophilic factor VIII - recombinant (Advate®) <sup>1</sup>	Baxter Healthcare	Control and prevention of bleeding episodes in adults and children with hemophilia A Perioperative management in adults and children with hemophilia A Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children with hemophilia A
antihemophilic factor VIII – plasma derived (Alphanate®) <sup>2</sup>	Grifols Biologics	Control and prevention of bleeding episodes in hemophilia A Surgical and/or invasive procedures in adult and pediatric patients with von Willebrand’s disease (vWD) in whom desmopressin is ineffective or contraindicated; it is not indicated for patients with severe vWD (type 3) undergoing major surgery
antihemophilic factor VIII – recombinant, Fc fusion protein (Eloctate®) <sup>3</sup>	Biogen Idec	Indicated for prevention and control of bleeding episodes in adults and children with hemophilia A (classical hemophilia) Perioperative management in adults and children with hemophilia A Routine prophylaxis therapy to prevent or reduce the frequency of bleeding episodes in adults and children with hemophilia A Not indicated for vWD
antihemophilic factor VIII – recombinant (Helixate FS®) <sup>4</sup>	Bayer Healthcare (distributed by CSL Behring)	Control and prevention of bleeding episodes in adults and children with hemophilia A Perioperative management in adults and children with hemophilia A Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in children with hemophilia A and no preexisting joint damage Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults with hemophilia A
antihemophilic factor VIII – plasma derived (Hemofil M®) <sup>5</sup>	Baxter Healthcare	Control and prevention of hemorrhagic episodes in (classical hemophilia) hemophilia A Not indicated for vWD
antihemophilic factor VIII – plasma derived (Humate-P®) <sup>6</sup>	CSL Behring	Treatment and prevention of bleeding in adults with hemophilia A (classical hemophilia) Treatment of spontaneous and trauma-induced bleeding episodes, and prevention of excessive bleeding during and after surgery for adults and pediatric patients with vWD; this applies to patients with severe and mild to moderate vWD where the use of desmopressin is known or suspected to be inadequate Not indicated for the prophylaxis of spontaneous bleeding episodes in vWD
antihemophilic factor VIII – plasma derived (Koate DVI®) <sup>7</sup>	Grifols Therapeutics (distributed by Kendrion Biopharma)	Treatment of hemophilia A (classical hemophilia) in which there is a demonstrated deficiency of activity of the plasma clotting Factor VIII to control or prevent bleeding episodes, or in order to perform emergency and elective surgery on individuals with hemophilia Not approved for use in vWD

**Factor VIII Products (continued)**

Drug	Manufacturer	Indications
antihemophilic factor VIII – recombinant (Kogenate FS®) <sup>8,9</sup>	Bayer Healthcare	Control and prevention of bleeding episodes in adults and children with hemophilia A Perioperative management in adults and children with hemophilia A Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in children with hemophilia A and reduce the risk of joint damage in children showing no preexisting joint damage Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults with hemophilia A <b>Not indicated for vWD</b>
antihemophilic factor VIII – plasma derived (Monoclate-P®) <sup>10</sup>	CSL Behring	Treatment of (classical hemophilia) hemophilia A <b>Not indicated or effective for vWD</b>
antihemophilic factor VIII – recombinant (Novoeight®) <sup>11</sup>	Novo Nordisk	Control and prevention of bleeding episodes in adults and children with hemophilia A Perioperative management in adults and children with hemophilia A Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children with hemophilia A <b>Not indicated for vWD</b>
antihemophilic factor VIII – recombinant, porcine sequence (Obizur®) <sup>12</sup>	Baxter Healthcare	Treatment of bleeding episodes in adults with acquired hemophilia A <b>Not indicated for the treatment of congenital hemophilia A or vWD</b>
antihemophilic factor VIII – recombinant (Recombinate®) <sup>13</sup>	Baxter Healthcare	Control and prevention of hemorrhagic episodes in (classical hemophilia) hemophilia A Perioperative management in patients with hemophilia A Not indicated for vWD
antihemophilic factor VIII – plasma derived (Wilate®) <sup>14</sup>	Octapharma USA	<b>For on-demand treatment and control of bleeding episodes children and adults with vWD</b> Perioperative management of bleeding in children and adults with vWD Not indicated for treatment of hemophilia A
antihemophilic factor VIII – recombinant (Xyntha®) <sup>15,16</sup>	Wyeth Pharmaceuticals	Control and prevention of bleeding episodes in <b>adults and children</b> with hemophilia A Perioperative management in <b>adults and children</b> with hemophilia A

## Factor IX Products

Drug	Manufacturer	Indications
coagulation factor IX – plasma derived (AlphaNine SD®) <sup>17</sup>	Grifols Biologics	Indicated for the prevention and control of bleeding in patients with Factor IX deficiency due to hemophilia B AlphaNine SD contains low, non-therapeutic levels of Factors II, VII, and X, and is not indicated for the treatment of Factor II, VII, or X deficiencies Not indicated for the treatment of hemophilia A patients with inhibitors to Factor VIII
coagulation factor IX – recombinant, FC fusion protein (Alprolix®) <sup>18</sup>	Biogen Idec	Indicated for prevention and control of bleeding episodes in adults and children with Factor IX deficiency, hemophilia B Perioperative management in adults and children with hemophilia B Routine prophylaxis therapy to prevent or reduce the frequency of bleeding episodes in adults and children with Factor IX deficiency, hemophilia B Not indicated for induction of immune tolerance therapy in patients with hemophilia B.
coagulation factor IX – plasma derived (Bebulin®/Bebulin VH®*) <sup>19</sup>	Baxter Healthcare	Indicated for the prevention and control of hemorrhagic episodes in hemophilia B patients; it is not indicated for use in the treatment of Factor VII deficiency; no clinical studies have been conducted to show benefit from this product for treating deficiencies other than Factor IX deficiency
coagulation factor IX – recombinant (BeneFIX®) <sup>20</sup>	Wyeth Pharmaceuticals	Indicated for the prevention and control of bleeding episodes in adults and pediatric patients with hemophilia B Indicated for perioperative management in adults and pediatric patients with hemophilia B Not indicated for the treatment of other factor deficiencies (e.g., Factors II, VII, VIII, and X), or the treatment of hemophilia A patients with inhibitors to Factor VIII
coagulation factor IX – recombinant (Ixinity®) <sup>21</sup>	Emergent BioSolutions	Indicated for the prevention and control of bleeding episodes in adults and pediatric patients 12 years of age or older with hemophilia B Indicated for perioperative management in adults and pediatric patients with hemophilia B Not indicated for the induction of immune tolerance therapy in patients with hemophilia B
coagulation factor IX – plasma derived (Mononine®) <sup>22</sup>	CSL Behring	Prevention and control of bleeding in Factor IX deficiency, also known as hemophilia B Not indicated in the treatment or prophylaxis of hemophilia A in patients with inhibitors to Factor VIII Mononine contains non-detectable levels of Factors II, VII, and X and is, therefore, not indicated for replacement therapy of these clotting factors
coagulation factor IX – plasma derived (Profilnine SD®) <sup>23</sup>	Grifols Biologics	Indicated for the prevention and control of bleeding in patients with Factor IX deficiency due to hemophilia B Not indicated for use in the treatment of Factor VII deficiency

### Factor IX Products (continued)

Drug	Manufacturer	Indications
coagulation factor IX – recombinant (Rixubis®) <sup>24</sup>	Baxter Healthcare	Indicated for prevention and control of bleeding episodes in adults and children with Factor IX deficiency, hemophilia B Perioperative management in adults and children with hemophilia B Routine prophylaxis therapy to prevent or reduce the frequency of bleeding episodes in adults and children with Factor IX deficiency, hemophilia B Not indicated for induction of immune tolerance therapy in patients with hemophilia B.

\*Bebulin VH is in the process of being phased out in favor of Bebulin.

### Factor VII Products\*\*

Drug	Manufacturer	Indications
anti-inhibitor coagulation complex containing factors II, IX, X and primarily activated factor VII – plasma derived (Feiba NF®) <sup>25</sup>	Baxter Healthcare	Indicated for the control and prevention of bleeding episodes in hemophilia A and hemophilia B patients with inhibitors Perioperative management in hemophilia A and hemophilia B patients with inhibitors Routine prophylaxis to prevent or reduce the frequency of bleeding episodes in hemophilia A and hemophilia B patients with inhibitors
coagulation factor VIIa - recombinant (Novo Seven RT®) <sup>26</sup>	Novo Nordisk	Treatment of bleeding episodes in hemophilia A or B with inhibitors and in acquired hemophilia Prevention of bleeding in surgical interventions or invasive procedures in hemophilia A or B with inhibitors and in acquired hemophilia Treatment of bleeding episodes in congenital Factor VII (FVII) Deficiency Prevention of bleeding in surgical interventions or invasive procedures in congenital FVII deficiency Treatment of Glanzmann's Thrombasthenia with refractoriness to platelet transfusions with or without antibodies

\*\*MasAc recommendations refer to these as factor VII products.

## OVERVIEW<sup>27,28,29</sup>

Hemophilia is a rare, inherited bleeding disorder where the blood does not clot properly due to an absence of one of the coagulation factors present in normal blood. Hemophilia is identified as an X-linked congenital bleeding disorder that has an estimated frequency of one in 10,000 births. The World Foundation of Hemophilia estimates the global prevalence of hemophilia at around 400,000 persons. It is estimated there are approximately 17,000 to 20,000 persons in the United States are afflicted with hemophilia.

Patients born with hemophilia may have varying degrees of coagulation or clotting factor deficiencies which serve to characterize the severity of their disease. There are two main types of hemophilia, type A and type B. Patients with type A hemophilia exhibit low or missing levels of clotting factor VIII (8), while those with type B have low or missing levels of clotting factor IX (9). Hemophilia A is also known as factor VIII deficiency, classical hemophilia, or standard hemophilia. Hemophilia B is also known as factor IX deficiency or Christmas disease. Hemophilia A is far more common than hemophilia B with

hemophilia A presenting in 80 to 85% of all patients. Hemophilia also typically affects males on the maternal side due to X-linked inheritance; however, females may also rarely be affected but are more commonly carriers of the disease. Up to 30% of newly diagnosed cases occur with no prior family history and are attributed to spontaneous mutations in either the F8 or F9 gene. While a diagnosis of hemophilia may be made from specific tests from an umbilical cord blood sample if there is a family history or suspicion of hemophilia, the disease is more commonly diagnosed when a recurrence of excessive bleeding problems is noted. The extent of bleeding episodes depends on the severity of hemophilia and these may be instances of external or internal bleeding. Persons who have mild hemophilia may not normally exhibit problems with clotting until after occurrences such as dental procedures, surgery, or an accident. More commonly, signs of hemophilia leading to a positive diagnosis include external bleeds such as: mouth bleeds following a cut, losing a tooth, or teething; heavy bleeding from minor cuts; bleeding from cuts that resume after short stoppages; and bleeding following circumcision. Internal signs of excessive bleeding that may indicate hemophilia include: easy and large areas of bruising, particularly in the large muscles of the body; spontaneous bleeding into joints (knees, elbows, and ankles), muscles, and soft tissues; joints that suddenly swell, are hot to the touch, and painful; blood in the stool or urine; long-lasting painful headaches; neck stiffness; double vision; repeated vomiting; and convulsions or seizures. Bleeding manifestations can lead to substantial morbidity, as well as mortality, if not properly treated.

Hemophilia, regardless of type (hemophilia A or B), is classified as mild, moderate, or severe depending on the intrinsic amount of clotting factor, either factor VII or factor IX, in the patient's blood. The following table details the severity, general clotting factor level, and bleeding episode characteristics for hemophilia. Bleeding into joints or hemarthrosis is more common than bleeds into muscles, the central nervous system, or other major bleeds and comprise between 70% to 80% of bleeds. Bleeding is more common in hinged joints such as ankles, elbows, and knees and less frequent on multi-axial joints such as hips, shoulders, and wrists. Frequent and prolonged bleeds in the hinged joints may cause these to become less functional over time.

Severity	Level of Clotting Factor	Bleeding Characteristics
Mild hemophilia	5% to less than 40% of normal (5-40 IU/dL)	Patient may never exhibit bleeds; spontaneous bleeds are rare; rarely has joint problems; may have severe bleeding with surgery, injections, or major trauma
Moderate hemophilia	1% to 5% of normal (1-5 IU/dL)	Patient may have rare or occasional spontaneous bleeds; may have prolonged bleeds with minor trauma or surgery; may experience joint problems
Severe hemophilia	Less than 1% of normal (less than 1 IU/dL)	Patient can experience spontaneous bleeds without injury; joint and muscle bleeds are common; may experience bleeds 1 to 2 times a week

In July 2012, the World Federation of Hemophilia published its comprehensive treatment guidelines for the disease with certain documents within the guidelines updated in May and June of 2014. The guidelines list the general principles of care for treatment of bleeding episodes. These include the prevention and treatment of bleeds with the specific factor concentrate for that patient's clotting deficiency; treatment of active bleeds as soon as possible and preferably within a two hour window of onset; help in teaching patients to recognize a bleeding aura which is often experienced prior to outward evidence of a bleed; the use of adjunctive therapies to help control bleeds, including compression and cold therapy; ensuring that patients seek experienced medical care, including a pediatric or adult hematologist and continuing care through a recognized hemophilia treatment center; use of patient training and home therapy to treat non-life-threatening bleeding episodes; comprehensive care plans that encourage and promote regular exercise to improve overall fitness while avoiding activities likely to cause trauma; maintaining good oral health and regular appointments to monitor health status.

The recommended treatment of bleeding episodes is dependent on several factors, including the patient's severity level, the location, and type of the injury or trauma, as well as the patient's overall status. Providing immediate treatment reduces the risk of lasting damage, the need for additional medication, and the reduction of pain, as well as additional treatments. It is important to note that a person with a bleeding disorder will not bleed faster than anyone else; however, the bleeding, if untreated, will last longer.

Von Willebrand's disease (vWD) is related to hemophilia A and is a group of inherited bleeding disorders related to defects of von Willebrand's Factor, which is needed to achieve hemostasis. Von Willebrand's factor includes Factor VIII. Unlike hemophilia A, vWD occurs equally in males and females. The disease leads to bleeding from impaired platelet adhesion and aggregation, which may be accompanied by reduced levels of Factor VIII. Mucous membrane and skin bleeding symptoms, as well as bleeding with surgical or other hemostatic challenges, may occur in these patients. The prevalence of the disease is estimated to affect between one in 100 to 10,000 individuals, but is noted to be higher in patients with type O blood. Pregnancy can increase vWF levels and make diagnosis difficult. There are three major subtypes of vWD identified. Type 1 is a partial quantitative deficiency of vWF deficiency and accounts for 75% of all patients. Type 2 is a more pronounced qualitative deficiency and comprises almost all the remaining 25% of patients. Type 3 is characterized as a complete vWF deficiency and occurs very rarely. Type 2 disease is further divided into four variants named 2A, 2B, 2M, 2N on the basis of identified phenotypes. For type 3 vWD patients, their inherent Factor VIII levels are typically very low. Acquired von Willebrand's syndrome may also occur, but is highly rare, occurring in fewer than one in 100,000 adults. Three main approaches for the treatment of vWD are used and include increasing plasma concentrations of vWF through stimulation with desmopressin (DDAVP); replacing vWF by using human plasma-derived viral inactivated concentrates; and promoting hemostasis by utilizing hemostatic agents with mechanisms other than increasing vWF. Regular prophylaxis for von Willebrand's patients is seldom required.

## PHARMACOLOGY<sup>30,31,32,33,34,35,36,37,38,39,40,41,42</sup>

Factor VIII and IX, along with Factor VII-containing products are intravenously administered and designed to supplement endogenous coagulation factors in patients with hemophilia A and B.

Products are differentiated based upon factor type, whether they are derived from pooled human plasma in the manufacturing process, as well as their level of purity or purification process.

Among hemophilia A or Factor VIII products, those produced by cloning of the Factor VIII gene, or the recombinant process, include Advate, Eloctate, **Novoeight, Obizur**, Helixate FS, Kogenate FS, Recombinate, and Xyntha. Kogenate FS and Helixate FS are both manufactured by Bayer Corporation who markets the recombinant Factor VIII product as Kogenate FS while CSL Behring distributes the product as Helixate FS. **Since Obizur** and Recombinate have been found to contain animal and/or human plasma-derived proteins in the cell culture medium, as well as in the final product formulation, they are considered first-generation products. Helixate FS and Kogenate FS, second-generation recombinant products, are considered an advance over first-generation products as they contain animal or human plasma in the medium but not in the final formulation vial. Advate, Eloctate, **Novoeight**, and Xyntha are considered third-generation Factor VIII products due to the fact that they do not contain any animal or human plasma-derived proteins in either the culture medium or final formulation. Xyntha is a B-domain deleted product that has the B-domain deleted from the Factor VIII gene prior to its insertion into Chinese hamster ovary cells. There is no known functional impact or role for the B-domain on factor VIII activity and its removal simply serves to increase the manufacturing output without affecting the resultant product's biological activity.

**In 2014, Obizur was introduced as the only Factor VIII product indicated for the treatment of acquired hemophilia A. In acquired hemophilia A, patients develop autoantibodies to their normal Factor VIII genes neutralizing circulating human Factor VIII thereby creating a functional factor deficiency. Obizur is a recombinant analogue of porcine Factor VIII. Once activated in the body, the porcine Factor VIII demonstrates comparable activity to normal endogenous human Factor VIII.**

Pooled human plasma-derived Factor VIII products for the treatment of hemophilia A include Alphanate, Hemofil M, Humate-P, Koate-DVI, Monoclate-P, and Wilate. With the exception of Wilate, the products are indicated for use in hemophilia A. Wilate is only indicated for patients with vWD and, although Wilate is not indicated in hemophilia A, it does contain Factor VIII. The ratio of vW factor to Factor VIII for Wilate is 1:1. Alphanate and Humate-P also carry an indication for treatment of vWD patients and their ratios of vW factor to factor VIII are 1.3:1 and 1.8-2.4:1, respectively. While both products may be used to treat vWD patients, they are not identical and should not be regarded as interchangeable.

When considering Factor IX products for the treatment of hemophilia B, Alprolix, BeneFIX, **Ixinity**, and Rixubis are products made by recombinant technology; all other products are derived from human plasma. **Additionally, the manufacturing process for these four products does not use any human or animal proteins making the risk of human blood-borne viral contamination extremely low.** AlphaNine SD and Mononine are human plasma-derived products that are considered to be of high purity in comparison to other plasma-derived Factor IX agents, employing added and improved methods of screening and viral depletion. While these additional methods serve to reduce the risk, there does remain the slight possibility of viral transmission. Transmission of human parvovirus B19 and hepatitis A have occurred; however, the risk has been reduced with the use of additional viral attenuation

methods such as ultrafiltration. Bebulin and Profilnine SD are also human plasma-derived products considered to be of lower purity. One of these product variations, Bebulin VH, is being phased-out in favor of Bebulin, which employs nanofiltration in addition to vapor heat for the purification process. These last two products are also referred to as Prothrombin Complex Concentrates (PCCs) as they contain detectable levels of other clotting factors including Factors II, VII, IX, and X. Bebulin and Profilnine SD can, in some instances, be used to treat patients with deficiencies of Factors II and X but they vary significantly in the amounts of factors that they contain. Further complicating the additional use is the fact that there commonly may be a marked differences in factor content between the different lots of the same agent produced by the same manufacturer.

In mid-2014, two new FC fusion protein factor products were introduced to the market; Eloctate (a recombinant Factor VIII product) and Alprolix (a recombinant Factor IX product). The theory behind the FC protein is to prolong the half-life of the coagulation factor in clinical use, thereby potentially increasing the interval of administration in prophylactic dosing regimens. The fusion protein does not increase the potency of the factor in correcting bleeds; it is only intended to attempt to prolong the half-life. While clinical trials did indicate the half-life may be increased in adults, the pharmacologic variability in pediatric hemophilia patients made any prolongation less certain. Manufacturer recommendations deferred to individual patient response and physician evaluation (see *Dosages* section).

The National Hemophilia Foundation Medical and Safety Advisory Committee (MASAC) published a revised document with recommendations regarding products used for the treatment of hemophilia and other bleeding disorders in June 2015. The recommendations state the risk of human viral contamination associated with recombinant Factor VIII products is reduced in comparison with the risk with plasma-derived Factor VIII products. There have been no reported instances of seroconversion of human immunodeficiency virus (HIV), hepatitis B virus (HBV), or hepatitis C virus (HCV) with recombinant derived products. Recombinant Factor VIII products are the recommended treatment of choice by MASAC for patients with hemophilia A. MASAC also recommended that manufacturers should work toward the removal of human albumin, as well as human and/or bovine proteins, from their manufacturing processes for the recombinant Factor VIII products.

The MASAC recommendations also address the use of plasma-derived Factor VIII products. The MASAC recommendations stated improved purification and viral-depleting processes along with increased donor screening processes have greatly reduced the risk of transmission from plasma-derived factor products of HIV, HBV, and HCV. There have not been any reported instances of seroconversion of HIV, HBV, or HCV as a result of the use of any plasma-derived Factor VIII products currently available in the United States. This includes products that undergo the following purification/viral inactivation measures: heating in aqueous solution (pasteurization), solvent-detergent treatment, and/or immunoaffinity chromatography purification. Each of these methods, whether employed alone or in conjunction with another process, appears to have greatly reduced the risk of viral transmission compared to older methods. Despite the improvements and lack of any documented transmission, MASAC maintains there remains a possibility of HIV-1, HIV-2, HBV, or HCV virus transmission with the use of currently marketed, viral-inactivated, plasma-derived products. The MASAC guidelines also caution that other non-lipid enveloped viruses may also be transmitted by plasma-derived Factor VIII products. MASAC acknowledges that additional measures have been taken, such as viral filtration, to further reduce any potential transmission risks.

Similar to the recommendations for Factor VIII products, MASAC stated that recombinant Factor IX products should also be considered the treatment of choice for hemophilia B patients due to the greatly reduced pathogen transmission risk compared with the plasma-derived Factor IX concentrates and may be considered the safest of the Factor IX agents. The recombinant products include Alprolix, BeneFIX, and Rixubis.

The factor product table located in the *Dosages* section of this review contains information about purification methods utilized by the various factor products, both recombinant and plasma-derived.

## PHARMACOKINETICS

The pharmacokinetics of various factor products dosed on an episodic or prophylactic basis is generally stable; however, this can vary depending on individual patient characteristics and factors, such as the presence of inhibitors.

## CONTRAINDICATIONS/WARNINGS

Contraindications for factor products are similar. All the factor agents are contraindicated in patients who have known anaphylactic or severe hypersensitivity reactions to the components of each product or in patients who are known to have a normal coagulation mechanism. Preliminary indications of allergic reactions, which may escalate to anaphylaxis, can include angioedema, chest tightness, hypotension, rash, nausea, vomiting, paresthesia, restlessness, wheezing, and dyspnea. Should any of these symptoms occur, use of the factor product should be discontinued and appropriate treatment administered.

Factor VIII products are contraindicated in patients who do not have hemophilia A or von Willebrand's disease.

Factor IX products are contraindicated in patients who do not have hemophilia B.

Feiba is contraindicated for the treatment of bleeding episodes resulting from coagulation factor deficiencies in the absence of inhibitors to coagulation Factor VIII or coagulation Factor IX. It is also contraindicated in patients with significant signs of disseminated intravascular coagulation (DIC) or in patients with acute thrombosis or embolism (including myocardial infarction).

Similarly, the warnings for all factor products are nearly equivalent. Initial treatment with all bleeding factor products should be performed under the supervision of a qualified healthcare professional experienced in the treatment of bleeding disorders. Recommended monitoring of plasma factor activity levels following administration of factor products is by one-stage clotting assay or a chromogenic substrate assay to confirm that adequate levels have been achieved and maintained, when clinically indicated.

The emergence of nephrotic syndrome has been reported following immune tolerance induction with Factor IX products in hemophilia B patients with Factor IX inhibitors and a history of allergic reactions to Factor IX.

In July 2014, NovoSeven added a boxed warning for thrombosis. Serious arterial and venous thrombotic events have been reported following the administration of NovoSeven RT. The risk appears greater in patients presenting with disseminated intravascular coagulation (DIC), advanced atherosclerotic disease, a crush injury, septicemia, or concomitant treatment with aPCCs/PCCs (activated or non-activated prothrombin complex concentrates) and uncontrolled post-partum hemorrhage due to circulating tissue factor (TF) or predisposing coagulopathy. The potential benefit of use in these patients should be strongly considered in light of the increased risk. Patients should be carefully monitored for any signs or symptoms indicating activation of the coagulation system or development of thrombosis.

## DRUG INTERACTIONS

There are no known or reported drug interactions with Factor VIII or Factor IX products. In many cases, drug interaction studies were not performed for coagulation factor products.

## ADVERSE EFFECTS<sup>43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67</sup>

All factor products, Factor VIII, Factor IX, and Factor VII-containing products, derived from pooled human plasma carry a risk of transmitting infectious diseases. These risks have been attenuated by various purification methods employed in their manufacture. In general, for all these products, the benefits of these products outweigh the risks.

The plasma-derived Factor IX products may present a greater risk of disseminated intravascular coagulation (DIC) or thrombosis when administered at frequent or prolonged intervals. This may be more critical for the Prothrombin Complex Concentrates (PCCs), Bebulin and Profilnine SD, that contain significant amounts of activated Factors VII, X, and prothrombin. While the risk may be increased with plasma-derived products, the recombinant Factor IX products Alprolix, BeneFIX, **Ixinity**, and Rixubis have also been associated with the development of thromboembolic complications.

Severe allergic reactions may occur in up to 50% of hemophilia B patients who have inhibitors. These reactions may include anaphylaxis to Factor IX administration; it is possible that reactions of this type could represent the first signs and symptoms of inhibitor development. It is recommended that all newly diagnosed hemophilia B patients, and, in particular, patients who have a family history of factor inhibitors, as well as those patients with genetic defects predisposed to inhibitor development, should receive their initial doses and factor administration treatments in a clinic, hemophilia treatment center, or hospital setting with the resources to immediately treat severe allergic reactions. These precautions should persist throughout the first 10 to 20 infusion treatments with Factor IX concentrates. It is possible that reactions can be delayed or occur later; however, in these cases, they may be less severe.

The most significant adverse event that may occur for hemophilia A or B patients is the development of inhibitors. For hemophiliacs, inhibitors are IgG antibodies that neutralize clotting factors administered to control bleeding episodes. With the development of products with a greater level of purification, the development of inhibitors to Factor VIII or Factor IX coagulation factors is recognized as the most severe treatment-related complication in hemophilia.

In situations where a patient who has previously responded positively to factor infusion therapy fails to respond clinically to clotting factors, the presence of inhibitors should be suspected and appropriate testing performed. In situations where inhibitors are present, the expected response, clinical effect, and half-life of the transfused clotting factor are severely compromised. There are no documented cases where inhibitor development was attributed to a change in the specific factor product used.

Inhibitors are more frequently encountered in persons with severe hemophilia in comparison to patients diagnosed those with moderate or mild hemophilia regardless of phenotype. The overall lifetime risk for the development of inhibitors in patients identified with severe hemophilia A is in the range between 20% and 30%. The risk is significantly lower for patients diagnosed with mild to moderate hemophilia falling to between 5% to 10% over a lifetime. Finally, the prevalence of inhibitor development is much less frequent in hemophilia B than hemophilia A, occurring in less than 5% of affected individuals. Regardless of hemophilia phenotype, inhibitors render the treatment of bleeding episodes with replacement factor concentrates difficult. Patients receiving clotting factor therapy should be screened for the potential of inhibitor development.

In severe hemophilia A, the median age of inhibitor development in patients with severe hemophilia A is three years or less as seen in most developed countries. For those patients with moderate to mild hemophilia A, the median age of inhibitor development is closer to 30 years of age. Development in mild to moderate patients is often seen in following intensive Factor VIII exposure as a result of surgery.

In patients with severe hemophilia, the presence of inhibitors does not change the site, frequency, or severity of bleeds. Conversely, in patients with moderate or mild hemophilia, the development of inhibitors may serve to neutralize endogenously synthesized Factor VIII, effectively converting the patient's phenotype to severe hemophilia. Bleeding manifestations in moderate to mild hemophilia that are complicated by the development of an inhibitor are frequently reminiscent of those seen in patients with acquired hemophilia A occurring as a result of the presence of auto-antibodies to a patient's own Factor VIII. In such cases, there is a greater predominance of mucocutaneous, urogenital, and gastrointestinal bleeding sites. As a result, the risk of severe complications or even death from bleeding may be greater in these patients.

The presence of an inhibitor and the level of inhibitor involvement (quantification of the titer), should be performed in a qualified laboratory, preferably using the Nijmegen-modified Bethesda assay method. When the patients are children, screening for inhibitors should be performed once every five exposure days up to 20 factor exposure days, then every 10 factor exposure days between 21 and 50 exposure days. Following the initial phase of testing, evaluations should be performed at least twice a year until 150 exposure days have occurred. When dealing with adults who have experienced more than 150 factor exposure days, a review should be performed approximately every 6 to 12 months. If there is an instance of failure to adequately respond to normal factor concentrate replacement dose in a patient who previously demonstrated a positive clinical response, this is an indication to assess for the emergence of an inhibitor.

Inhibitor measurement should also be done for all patients who have been intensively treated with factor therapy for more than five days duration, within four weeks of their last infusion. Inhibitors should be assessed prior to surgery or if recovery assays are not as expected. Again, in any case where the clinical response to treatment of bleeding is sub-optimal in the post-operative period, an assessment should be performed.

There are two primary types of inhibitors identified based upon their measurement in Bethesda units (BU). Low responding inhibitors are defined as those presenting an inhibitor level that is persistently < 5 BU/mL, while high responding inhibitors are defined as those having a Bethesda level  $\geq$  5 BU/mL.

Although high responding inhibitors tend to be persistent, if the patient does not undergo factor infusion therapy for a long period, the inhibitor titer levels may drop or even become undetectable. Unfortunately, this effect is only temporary and there will be a recurrence of an anamnestic response along with a rise in the inhibitor titer level generally in three to five days following exposure to specific factor products. In contrast, certain low titer inhibitors may be transient and disappear commonly within six months of their initial detection and documentation. In these cases, this occurs despite any recent antigenic challenge with factor concentrates. In certain instances, very low titer inhibitors may not be detected by the Bethesda inhibitor assay, but are identified by a poor clinical response, sub-optimal recovery, and/or a shortened half-life for the agent utilized following clotting factor infusions.

## SPECIAL POPULATIONS

### Pediatrics

Please refer to the Food and Drug Administration (FDA)-approved indications table.

### Pregnancy

The Factor VIII products Advate, Alphanate, Eloctate, Helixate FS, Hemofil M, Humate-P, Kogenate FS, Monoclate-P, **Novoeight**, **Obizur**, Recombinate, Wilate, and Xyntha all carry a pregnancy category designation C. Similarly, Factor IX products AlphaNine SD, Alprolix, Bebulin/Bebulin VH, BeneFIX, **Ixinity**, Mononine, Profilnine SD, and Rixubis also carry a pregnancy category C. Feiba NF/Feiba VH and NovoSeven RT are also pregnancy category C.

### Geriatrics

Clinical studies of factor products did not include subjects 65 years of age or older, or did not have sufficient patients to determine differences in response or efficacy. Dosing for elderly patients should be individualized.

## DOSAGES<sup>68,69,70,71,72,73,74,75,76,77,78</sup>

There are two primary methodologies employed in factor replacement therapy. Commonly, mild to moderate hemophilia A and B patients may be treated with episodic or 'on demand' treatment of bleeds. Severe hemophilia patients, particularly younger hemophilia patients, are recommended to begin prophylactic factor replacement therapy where individualized dosages are given on a scheduled basis.

General dosing information for hemophilia A is commonly included in the respective product labels. These recommendations are usually provided as a range or percentage based on Factor VIII or Factor IX activity in the blood. The percentages or range can vary for each individual due to a variety of reasons including hemorrhage types, different surgical procedures, the presence or absence of inhibitors, and acute treatment versus prophylactic therapy. To assure optimal treatment outcomes, it is recommended that factor activity levels be monitored during replacement therapy. Ultimately, dosing decisions should be under the direction of the physician treating the condition.

In 2007, the National Hemophilia Foundation Medical and Scientific Advisory Council (MASAC) made recommendations regarding prophylaxis dosing (regular administration of clotting factor concentrate to prevent bleeding). Because of the benefits demonstrated by prophylactic therapy, MASAC recommends that prophylaxis be considered optimal therapy for individuals with severe hemophilia A or B. Prophylaxis therapy should be initiated early, with the goal of keeping trough values for Factor VIII or Factor IX levels above 1% between doses, although benefit is still seen when trough levels fall below the target goal. This can usually be accomplished by following one of two commonly accepted prophylaxis protocols. The first is the Malmö protocol where 25-40 IU/kg per dose are administered three times per week for patients with hemophilia A or twice a week for patients with hemophilia B. The second is the Utrecht protocol where 15-30 IU/kg per dose are given three times per week for patients with hemophilia A or twice a week for patients with hemophilia B. Some variations of these dosing schedules call for every other day administration. Protocols should be individualized to the extent possible taking into account age, bleeding phenotype, activity, cost, and factor availability. Individuals should have regular follow-up visits for evaluation of joint status. Where applicable, prophylaxis dosing recommendations from manufacturers for specific factor products is summarized in the table below.

There are not specific guidelines regarding when to stop prophylaxis or at what age prophylaxis benefit decreases, although some data suggest that some young adults do well after prophylaxis has ended. Recombinant factors products are the most appropriate choice for prophylaxis because of markedly reduced risk of blood-borne infection; however, as noted earlier, there are data that suggest that plasma-derived Factor VIII products may have a marginally-reduced risk although the conclusion is not definitive. Potential reasons to discontinue prophylaxis include development of an inhibitor or patient preference (with physician agreement).

## MANUFACTURER RECOMMENDED PROPHYLAXIS DOSING

### Factor VIII Products

Drug	Manufacturer Recommended Starting Doses for Prophylaxis
antihemophilic factor VIII - recombinant (Advate®) <sup>79</sup>	For children and adults – 20-40 IU/kg every other day (3 to 4 times a week) or every third day to maintain factor VIII trough levels greater than or equal to 1% Adjust dosing based on patient’s clinical response
antihemophilic factor VIII – plasma derived (Alphanate®) <sup>80</sup>	Dose (units) = body weight (kg) x desired FVIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL). The dosing frequency should be determined by the type of bleeding episode and the recommendation of the physician.
antihemophilic factor VIII – recombinant, Fc fusion protein (Eloctate) <sup>81</sup>	For children and adults – 50 IU/kg every 4 days Adjust dosing based upon patient’s clinical response Dosing range based upon response may be adjusted within 25-65 IU/kg every 3 to 5 days Doses of up to 80 IU/kg with greater frequency may be required for children less than 6 years of age
antihemophilic factor VIII – recombinant (Helixate FS®) <sup>82</sup>	For children 15 years and younger – 25 IU/kg every other day For adults 15 years and older - 25 IU/kg 3 times a week Adjust dosing based on patient’s clinical response
antihemophilic factor VIII – plasma derived (Hemofil M®) <sup>83</sup>	N/A
antihemophilic factor VIII – plasma derived (Humate-P®) <sup>84</sup>	N/A
antihemophilic factor VIII – plasma derived (Koate DVI®) <sup>85</sup>	N/A
antihemophilic factor VIII – plasma derived (Monoclate-P®) <sup>86</sup>	N/A
antihemophilic factor VIII – recombinant (Novoeight®) <sup>87</sup>	For children less than 12 years and younger: 20-50 IU/kg every other day or 20-60 IU/kg 3 times a week For adults 12 years and older: 20-40 IU/kg every other day or 20-50 IU/kg 3 times a week Adjust dosing based on patient’s clinical response
antihemophilic factor VIII – recombinant, porcine sequence (Obizur®) <sup>88</sup>	N/A
antihemophilic factor VIII – recombinant (Recombinate®) <sup>89</sup>	N/A
antihemophilic factor VIII – plasma derived (Wilate®) <sup>90</sup>	N/A
antihemophilic factor VIII – recombinant (Xyntha®) <sup>91,92</sup>	N/A

## Factor IX Products

Drug	Manufacturer Recommended Starting Doses for Prophylaxis
coagulation factor IX – plasma derived (AlphaNine SD®) <sup>93</sup>	N/A
coagulation factor IX – Recombinant, FC fusion protein (Alprolix) <sup>94</sup>	For children and adults – 50 IU/kg once a week or 100 IU/kg once every 10 days Adjust dosing based upon patient’s clinical response
coagulation factor IX – plasma derived (Bebulin®/Bebulin VH®*) <sup>95</sup>	N/A
coagulation factor IX – recombinant (BeneFIX®) <sup>96</sup>	N/A
coagulation factor IX – recombinant (Ixinity®) <sup>97</sup>	N/A
coagulation factor IX – plasma derived (Mononine®) <sup>98</sup>	N/A
coagulation factor IX – plasma derived (Profilnine SD®) <sup>99</sup>	N/A
coagulation factor IX – Recombinant (Rixubis) <sup>100</sup>	For patients less than 12 years of age previously treated with Factor IX – 60-80 IU/kg twice a week For patients 12 years and older previously treated with Factor IX – 40-60 IU/kg twice a week Adjust dosing based upon patient’s clinical response

\*Bebulin VH is in the process of being phased out in favor of Bebulin.

## Factor VII Products\*\*

Drug	Manufacturer Recommended Starting Doses for Prophylaxis
anti-inhibitor coagulation complex containing factors II, IX, X and primarily activated factor VII – plasma derived (Feiba NF) <sup>101</sup>	N/A Adjust dosing based on patient’s clinical response
coagulation factor VIIa - recombinant (Novo Seven RT) <sup>102</sup>	N/A Adjust dosing based on patient’s clinical response

\*\*MASAC recommendations refer to these as Factor VII products.

Episodic dosing is also often based on physician preference taking into account the patient's severity, location, and type of trauma. The goal of episodic treatment is to raise the factor level in the blood from 40% to 100% depending on the location and level of injury. These calculations include the patient's current weight, the factor level to be reached, and a constant depending on whether the patient had hemophilia A or B. As previously stated, final dosing decisions should be under the direction of the physician treating the condition and take into account patient and product specifics.

For hemophilia A patients a commonly accepted calculation is: patient weight pounds divided by 4.4 multiplied by the factor level or correction factor desired. The result is the number of Factor VIII units required:

$$\text{Weight in pounds} \div 4.4 \times \text{factor level desired} = \text{Factor VIII units required.}$$

For hemophilia B patients, a similar commonly accepted calculation is: patient weight pounds divided by 2.2 multiplied by the factor level or correction factor desired. The result is the number of Factor IX units required:

$$\text{Weight in pounds} \div 2.2 \times \text{factor level desired} = \text{Factor IX units required.}$$

In general, every attempt should be made to match the vial strength to the final correction required. It is permissible for hemophilia A patients to receive a bit more factor than the maximum calculated dose; however, hemophilia B patients should be cautioned not to exceed the maximum units required.<sup>103</sup> Depending on severity, if bleeding persists beyond 24 hours, the physician should be consulted and additional doses may be required.

One of the most complex and problematic aspects of treating hemophilia patients is the development of inhibitors, IgG antibodies, that neutralize exogenous clotting factors. If a patient does not respond clinically to the usual dose of factor which has previously been responsive, the development of an inhibitor must be suspected as these may be considered the most severe treatment-related complication in hemophilia. In these cases, care must be coordinated with the expertise of a hematologist who specializes in bleeding disorders as well as a hemophilia treatment center.

There are three primary methodologies for the treatment of persons with inhibitors including: High-Dose Clotting Factor Concentrates; Bypassing Agents; and Immune Tolerance Induction Therapy. Bypassing agents are special blood products used to treat bleeding in people with high titer inhibitors. They are called bypassing agents since, instead of replacing the missing factor, they go around (or bypass) the factors that are blocked by the inhibitor to help the body form a normal clot. People taking bypassing agents should be monitored closely to make sure the blood is not clotting too much or clotting in the wrong place in the body; and Immune Tolerance Induction (ITI) Therapy: The goal of ITI therapy is to reduce/eliminate the inhibitor reaction from the blood and to teach the body to accept clotting factor concentrate treatments. With ITI therapy, people receive large amounts of clotting factor concentrates every day for many weeks or months.

ITI therapy requires specialized medical expertise, is costly, and may take a long time to work. In many cases, ITI gets rid of the inhibitor. However, patients may need to continue taking frequent, large amounts of factor concentrates for many years to keep the inhibitor from coming back.

The choice of a specific product in treating persons with inhibitors involves consideration of the titer of the inhibitor, clinical response to product, site, and nature of bleed. Inhibitor activity is measured in Bethesda units (BU). Patients who have a confirmed low-responding inhibitor, (<5 BU/mL), may be treated with factor replacement at a much higher dose than normal. This increased dosing serves, if possible, to neutralize the inhibitor with excess factor activity and stop and active bleeding. Also known as High-Dose Clotting Factor Concentrate, it is important to test the blood and measure the factor level after this new treatment schedule is established to see if the inhibitor is gone.

Patients with a documented high responding inhibitor ( $\geq 5$ BU/mL), but who have low titers, may be treated in a similar manner in emergency situations until an anamnestic response occurs. This usually occurs in three to five days, with the goal of eliminating additional treatment with concentrates that only contain the missing factor.

Patients who have a Bethesda assay inhibitor level  $\geq 5$  BU are unlikely to show an effective clinical response to factor replacement designed to overwhelm the inhibitor without continuous infusion therapy at ultra high doses. In these instances, the use of alternative bypassing agents, such as Factor VII-containing products (Feiba NF, Novo Seven RT) or, in some cases, prothrombin complex concentrates (PCC), including the activated forms (aPCC), may prove more successful. It has been shown that efficacy of two doses of Factor VII and one dose of aPCC for management of joint bleeding are essentially equivalent. As with all factor therapy, however, some patients may respond better to one agent than the other, underscoring the importance of individualized therapy.

One additional therapy available for patients with severe hemophilia A is the potential eradication of inhibitors via Immune Tolerance Induction (ITI) therapy. One primary issue is that, prior to the initiation of ITI therapy, high-responding patients diagnosed with high-responding inhibitors should avoid the use of any Factor VIII products so inhibitor titers may fall in the hope of avoiding a persistent anamnestic rise. To date, no optimal regimen for ITI, either through product selection or dose requirements, has been defined or achieved a consensus. An international trial comparing the dosage of 50 IU/kg three times a week to 200 IU/kg daily was recently stopped due to safety concerns although interpretation of the data collected is an ongoing process.

Overall, the response to ITI may be less favorable in patients with moderate/mild hemophilia and current experience with ITI for hemophilia B inhibitor patients is limited. The principles for treatment of hemophilia B inhibitor patients is similar, but rate of success has proven is much lower. There is a possibility of the development of nephrotic syndrome by hemophilia B patients who have a documented history of severe allergic reactions to Factor IX and undertake ITI therapy. The resultant nephrotic syndrome is not always reversible upon cessation of the ITI therapy. Alternative treatment schedules, which include the addition of immunosuppressive therapies, are reported to have been successful.

While there has been interest in the use of immunosuppressive therapies for hemophilia patients with inhibitors, their role is not yet defined. At present, there is no consensus whether they have a place in the management of these patients.

## Factor VIII Products

Drug	Purification	Storage	Availability
antihemophilic factor VIII - recombinant (Advate®) <sup>104</sup>	Immunoaffinity chromatography (IC) and solvent/detergent (SD)	Up to 6 months but the temperature should not to exceed 30°C (86°F); the powder form should be refrigerated	Single-dose vials of 250 IU, 500 IU, 1,000 IU, 1,500 IU, 2,000 IU, 3,000 IU, and 4,000 IU
antihemophilic factor VIII – plasma derived (Alphanate®) <sup>105</sup>	Affinity chromatography (AC), solvent/detergent (SD), and dry heat 72 hours at 80°C	Up to the expiration date as long as storage temperature does not exceed 25 °C (77 °F) - do not freeze	Single-dose vials in strengths of 250 IU, 500 IU, 1,000 IU, 1,500 IU and 2,000 IU
antihemophilic factor VIII – recombinant, Fc fusion protein (Eloctate) <sup>106</sup>	Solvent/detergent (SD) and nanofiltration	Store between 2°C to 8°C – do not freeze. May store at room temperature not to exceed 30°C (86°F) for up to 6 months. Following storage at room temperature, do not refrigerate again.	Single-use vials of 250 IU, 500 IU, 750 IU, 1,000 IU, 1,500 IU, 2,000 IU, and 3,000 IU with diluent
antihemophilic factor VIII – recombinant (Helixate FS®) <sup>107</sup>	Immunoaffinity chromatography (IC) and solvent/detergent (SD)	Up to 12 months but the temperature should not exceed 25°C (77°F)	Single-use glass vials in strengths of 250 IU, 500 IU, 1,000 IU, 2,000 IU, and 3,000 IU
antihemophilic factor VIII – plasma derived (Hemofil M®) <sup>108</sup>	Immunoaffinity chromatography (IC) and solvent/detergent (SD)	Based upon the bottle expiration date but the temperature should not exceed 30°C (86°F)	Single-dose bottles in strengths of 250 IU, 500 IU, 1,000 IU, and 1,700 IU
antihemophilic factor VIII – plasma derived (Humate-P®) <sup>109</sup>	Pasteurization 10 hours at 60°C	Up to 24 months or the expiration date on the bottle but the temperature should not exceed 25°C (77°F)	Single-dose vials in strengths of 250, 500 and 1,000 IU of Factor VIII/vial and 600, 1,200 and 2,400 IU vWF:RCo/vial
antihemophilic factor VIII – plasma derived (Koate DVI®) <sup>110</sup>	Solvent/detergent (SD) and dry heat 72 hours at 80°C	Up to 6 months but the temperature should not exceed 30°C (86°F)	Single-dose bottles in strengths of 250 IU, 500 IU, and 1,000 IU
antihemophilic factor VIII – recombinant (Kogenate FS®) <sup>111,112</sup>	Immunoaffinity chromatography (IC) and solvent/detergent (SD)	At +2°C to +8°C (36°F to 46°F) for up to 30 months from manufacture date. Within the 30-month period, may be stored for a period of up to 12 months at up to +25°C or 77°F	Single-use glass vials in strengths of 250 IU, 500 IU, 1,000 IU, 2,000 IU, and 3,000 IU
antihemophilic factor VIII – plasma derived (Monoclate-P®) <sup>113</sup>	Immunoaffinity chromatography (IC) and pasteurization 10 hours at 60°C	At +2°C to +8°C (36°F to 46°F) up to expiration date Up to 6 months but the temperature should not exceed 25°C (77°F)	Single-dose vials in strengths of 250 IU, 500 IU, 1,000 IU, and 1,500 IU

**Factor VIII Products (continued)**

<b>Drug</b>	<b>Purification</b>	<b>Storage</b>	<b>Availability</b>
antihemophilic factor VIII – recombinant (Novoeight®) <sup>114</sup>	Immunoaffinity chromatography (AC), solvent/detergent (SD), anion-exchange chromatography, gel filtration and nanofiltration	At +2°C to +8°C (36°F to 46°F) for up to 30 months from manufacture date. Within the 30 month period, may be stored for a period of up to 12 months at up to +30°C or 86°F. Following storage at room temperature, do not refrigerate again. Do not use after the end of the 12-month period at room temperature storage, or after the expiration date stated on the vial, whichever occurs earlier.	Single-use glass vials in strengths of 250 IU, 500 IU, 1,000 IU, 1,500 IU, 2,000 IU, and 3,000 IU
antihemophilic factor VIII – recombinant, porcine sequence (Obizur®) <sup>115</sup>	Chromatography, solvent/detergent (SD) and nanofiltration	At +2°C to +8°C (36°F to 46°F) up to expiration date	Single-dose vials in strengths of 500 U
antihemophilic factor VIII – recombinant (Recombinate®) <sup>116</sup>	Immunoaffinity chromatography (IC)	Based upon the bottle expiration date but the temperature should not exceed 30°C (86°F)	Single-dose vials in the following strengths: 220 to 400 IU; 401 to 800 IU; 801 to 1,240 IU; 1,241 to 1,800 IU; and 1,801 IU to 2,400 IU
antihemophilic factor VIII – plasma derived (Wilate®) <sup>117</sup>	Solvent/detergent (SD) and dry heat 2 hours at 100°C	Up to 36 months at +2°C to +8°C (36°F to 46°F) in the original containers to protect from light from the date of manufacture  May be stored for a period of up to 6 months at room temperature. Once stored at room temperature, do not return to the refrigerator.	Single-dose vial in strengths of 500 IU VWF:RCo and 500 IU Factor VIII activities and 1,000 IU VWF:RCo and 1,000 IU Factor VIII activities
antihemophilic factor VIII – recombinant (Xyntha®) <sup>118,119</sup>	Solvent/detergent (SD) and nanofiltration	Up to 3 months but the temperature should not exceed 25°C (77°F)	Single-use vials and prefilled dual-chamber syringes (Solofuse) in strengths of 250 IU, 500 IU, 1,000 IU, and 2,000 IU. Solofuse is available in 3,000 IU

## Factor IX Products

Drug	Purification	Storage	Availability
coagulation factor IX – plasma derived (AlphaNine SD®) <sup>120</sup>	Dual affinity chromatography (AC), solvent/detergent (SD), and nanofiltration	Store between 2°C to 8°C. May store at room temperature not to exceed 30°C (86°F) for up to 1 month	Single-dose vials of 500 IU, 1,000 IU, and 1,500 IU with diluent
coagulation factor IX – recombinant, FC fusion protein (Alprolix) <sup>121</sup>	Column chromatography and nanofiltration	Store between 2°C to 8°C – do not freeze. May store at room temperature not to exceed 30°C (86°F) for up to 6 months. Following storage at room temperature, do not refrigerate again.	Single-use vials of 500 IU, 1,000 IU, 2,000 IU, and 3,000 IU with diluent
coagulation factor IX – plasma derived (Bebulin®/Bebulin VH®*) <sup>122</sup>	Nanofiltration (Bebulin) and vapor heat (Bebulin VH)	Store at a temperature between 36°F to 46°F	Single-dose vials contain 200 to 1,200 IU and diluent
coagulation factor IX – recombinant (BeneFIX®) <sup>123</sup>	Affinity chromatography (AC) and viral filtration	Store between 2°C to 8°C. May store at room temperature not to exceed 30°C (86°F) for up to 6 months.	Single-use vials contain 250 IU, 500 IU, 1,000 IU, 2,000 IU, and 3,000 IU with diluent
coagulation factor IX – recombinant (Ixinity®) <sup>124</sup>	Chromatography, solvent/detergent (SD) and nanofiltration	Store between 2°C to 25°C (36°F to 77°F) Do not freeze	Single-use vials contain 500 IU, 1,000 IU and 1,500 IU with diluent Multi-vial (2) Kit with 1,000 IU and 1,500 IU vials
coagulation factor IX – plasma derived (Mononine®) <sup>125</sup>	Immunoaffinity chromatography (IC), sodium thiocyanate and ultrafiltration	Store between 2°C to 8°C. May store at room temperature not to exceed 25°C (77°F) for up to 1 month.	Single-dose vials contain approximately 500 IU and 1,000 IU with diluent
coagulation factor IX – plasma derived (Profilnine®) <sup>126</sup>	Solvent/detergent (SD) and nanofiltration	Store between 2°C to 8°C. May store at room temperature not to exceed 30°C (86°F) for up to 3 months.	Single-dose vials of 500 IU, 1,000 IU, and 1,500 IU
coagulation factor IX – recombinant (Rixubis) <sup>127</sup>	Affinity chromatography (AC), solvent/detergent (SD), and nanofiltration	Store between 2°C to 8°C for up to 24 months – do not freeze. May store at room temperature not to exceed 30°C (86°F) for up to 12 months within the 24-month time frame. Following storage at room temperature, do not refrigerate again.	Single-use vials contain 250 IU, 500 IU, 1,000 IU, 2,000 IU, and 3,000 IU with diluent

## Factor VII Products

Drug	Purification	Storage	Availability
anti-inhibitor coagulation complex containing factors II, IX, X and primarily activated factor VII – plasma derived (Feiba NF) <sup>128</sup>	Nanofiltration and vapor heat	Store at room temperature not to exceed 25°C (77°F) up to the label expiration; do not freeze	Single-dose vials of 500 U, 1,000 U, and 2,500 U with diluent
coagulation factor VIIa – recombinant (Novo Seven RT) <sup>129</sup>	Immunoaffinity chromatography (IC)	Store between 2°C to 25°C (36°F to 77°F) up to the label expiration date; do not freeze	Single-dose vials of 1 mg, 2 mg, 5 mg and 8 mg with diluent

MASAC recommendations refer to these as Factor VII products

## CLINICAL TRIALS

### Search Strategies

Studies were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the FDA-approved use of all drugs in this class. Studies included for analysis in the review were published in English, performed with human participants, and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question, and include follow-up (endpoint assessment) of at least 80% of participants entering the investigation. Despite some inherent bias found in all studies including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship and/or funding must be considered, the studies in this review have also been evaluated for validity and importance. Many of the trials with agents in this class were performed in an open-label manner; introduction of bias must be considered when evaluating study findings.

The method of administration and associated monitoring makes it difficult to perform properly blinded studies with these drugs.

For the majority of Factor VIII, Factor IX, and Factor VII products, no direct comparisons to other coagulation products have been conducted; therefore, no conclusions regarding the comparative safety or efficacy can be made.

## SUMMARY

There are little direct comparative data for both the Factor VIII or the Factor IX products. In general, for Factor VIII, there are no apparent differences in efficacy among the various products for controlling bleeding episodes with all products demonstrating a success rate of 90% when dosed according to the patient's requirements. Factor products, with the exception of Obizur indicated only for the extremely rare acquired hemophilia A, may be considered interchangeable in efficacy, although other issues are also important when considering product selection. These may include purification processes and the perception of purity, as well as potential inhibitor development. Additional considerations include, but are not limited to, patient and/or physician preference, vial size for dosing convenience, cost, and product availability. The impact of the new FC protein products designed to increase factor half-life has yet to be defined as patient pharmacologic variability and potential product cost may prevent widespread utilization.

## REFERENCES

- 1 Advate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2014.
- 2 Alphanate [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; March 2015.
- 3 Elocatate [package insert]. Cambridge, MA: Biogen Idec, Inc.; June 2014
- 4 Helixate FS [package insert]. Kankakee, IL: CSL Behring, LLC; May 2014.
- 5 Hemofil M [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2012.
- 6 Humate-P [package insert]. Kankakee, IL: CSL Behring, LLC; June 2014.
- 7 Koate-DVI [package insert]. Fort Lee, NJ: Kendrion Biopharma, Inc.; August 2012.
- 8 Kogenate FS [package insert]. Whippany, NJ: Bayer Healthcare LLC; August 2015.
- 9 Kogenate FS with Bio-Set [package insert]. Whippany, NJ: Bayer Healthcare LLC; August 2015.
- 10 Monoclate-P [package insert]. Kankakee, IL: CSL Behring, LLC; February 2014.
- 11 Novoeight [package insert]. Plainsboro, NJ: Novo Nordisk, Inc.; March 2015
- 12 Obizur [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; October 2014
- 13 Recombinate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; December 2010.
- 14 Wilate [package insert]. Hoboken, NJ: Octapharma USA; August 2015.
- 15 Xyntha [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 16 Xyntha Solofuse [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 17 AlphaNine SD [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; January 2013.
- 18 Alprolix [package insert]. Cambridge, MA: Biogen Idec, Inc.; March 2014.
- 19 Bebulin [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; July 2012.
- 20 BeneFIX [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; August 2015.
- 21 Ixinity [package insert]. Baltimore, MD: Emergent BioSolutions, Inc.; April 2015
- 22 Mononine [package insert]. Kankakee, IL: CSL Behring, LLC; February 2013.
- 23 Profilnine [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; May 2014.
- 24 Rixubis [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; September 2014
- 25 Feiba NF [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; November 2013.
- 26 NovoSeven RT [package insert]. Plainsboro NJ: Novo Nordisk, Inc.; June 2015.
- 27 NHLBI Working Group: Hemophilia and Thrombosis Disorders Executive Summary. Available at: <http://www.nhlbi.nih.gov/meetings/workshops/hrht.htm>. Accessed September 19, 2015.
- 28 WFH Guidelines for the Management of Hemophilia. Available at: <http://www.wfh.org/en/resources/wfh-treatment-guidelines>. Accessed September 19, 2015.
- 29 The Hemophilia, von Willebrand Disease and Platelet Disorders Handbook. Available at: <http://www.hog.org/handbook>. Accessed September 19, 2015.
- 30 Advate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2014.
- 31 Helixate FS [package insert]. Kankakee, IL: CSL Behring, LLC; May 2014.
- 32 Kogenate FS [package insert]. Whippany, NJ: Bayer Healthcare LLC; August 2015.
- 33 Recombinate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; December 2010.
- 34 Xyntha [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 35 MASAC (Medical and Scientific Advisory Council) Recommendations concerning products licensed for the treatment of hemophilia and other bleeding disorders (Revised May 2014). MASAC Document #235. Available at <https://www.hemophilia.org/Researchers-Healthcare-Providers/Medical-and-Scientific-Advisory-Council-MASAC/MASAC-Recommendations/MASAC-Recommendations-Concerning-Products-Licensed-for-the-Treatment-of-Hemophilia-and-Other-Bleeding-Disorders>. Accessed September 19, 2015.
- 36 MASAC (Medical and Scientific Advisory Council) Recommendations concerning products licensed for the treatment of hemophilia and other bleeding disorders (Revised May 2014). MASAC Document #235. Available at <https://www.hemophilia.org/Researchers-Healthcare-Providers/Medical-and-Scientific-Advisory-Council-MASAC/MASAC-Recommendations/MASAC-Recommendations-Concerning-Products-Licensed-for-the-Treatment-of-Hemophilia-and-Other-Bleeding-Disorders>. Accessed September 19, 2015.

- 37 Elocate [package insert]. Cambridge, MA: Biogen Idec, Inc.; June 2014.
- 38 Alprolix [package insert]. Cambridge, MA: Biogen Idec, Inc.; March 2014.
- 39 Rixubis [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; September 2014
- 40 MASAC (Medical and Scientific Advisory Council) Recommendation regarding the use of recombinant clotting factor products with respect to pathogen transmission (Revised May 2014). MASAC Document #226. Available at: <https://www.hemophilia.org/Researchers-Healthcare-Providers/Medical-and-Scientific-Advisory-Council-MASAC/MASAC-Recommendations/MASAC-Recommendation-Regarding-the-Use-of-Recombinant-Clotting-Factor-Products-with-Respect-to-Pathogen-Transmission>. Accessed September 19, 2015.
- 41 Ixinity [package insert]. Baltimore, MD: Emergent BioSolutions, Inc.; April 2015
- 42 Novoeight [package insert]. Plainsboro, NJ: Novo Nordisk, Inc.; March 2015
- 43 AlphaNine SD [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; January 2013.
- 44 Bebulin [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; July 2012.
- 45 BeneFIX [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; August 2015.
- 46 Mononine [package insert]. Kankakee, IL: CSL Behring, LLC; February 2013.
- 47 Profilnine [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; May 2014.
- 48 Chitlur M, Warrior I, Rajpurkar M, Lusher JM. Inhibitors in factor IX deficiency a report of the ISTH-SSC international FIX inhibitor registry (1997-2006). *Haemophilia* 2009;15(5):1027-31.
- 49 Recht M, Pollmann H, Tagliaferri A, et al. A retrospective study to describe the incidence of moderate to severe allergic reactions to factor IX in subjects with haemophilia B. *Haemophilia* 2011;17(3):494-9.
- 50 Astermark J, Altisent C, Batorova A, et al; European Haemophilia Therapy Standardisation Board. Non-genetic risk factors and the development of inhibitors in haemophilia: a comprehensive review and consensus report. *Haemophilia* 2010;16(5):747-66.
- 51 Wight J, Paisley S. The epidemiology of inhibitors in haemophilia A: a systematic review. *Haemophilia* 2003;9(4):418-35.
- 52 Eckhardt CL, Menke LA, Van Ommen CH, et al. Intensive peri-operative use of factor VIII and the Arg593->Cys mutation are risk factors for inhibitor development in mild/moderate hemophilia A. *J Thromb Haemost* 2009;7:930-37.
- 53 Kempton CL, Soucie JM, et al. In non-severe hemophilia A the risk of inhibitor after intensive factor treatment is greater in older patients: a case-control study. *JTH* 2010 Oct;8(10):2224-31.
- 54 Hay CR. Factor VIII inhibitors in mild and moderate-severity haemophilia A. *Haemophilia* 1998;4(4):558-63.
- 55 Bolton-Maggs PH, Pasi KJ. Haemophilias A and B. *Lancet* 2003 May 24;361(9371):1801-9.
- 56 Berntorp E, Collins P, D'Oiron R, et al. Identifying non-responsive bleeding episodes in patients with haemophilia and inhibitors: a consensus definition. *Haemophilia* 2011;17(1):e202-10.
- 57 Hay CR, Brown S, Collins PW, Keeling DM, Liesner R. The diagnosis and management of factor VIII and IX inhibitors: a guideline from the United Kingdom Haemophilia Centre Doctors Organisation. *Br J Haematol* 2006;133:591-605.
- 58 McMillan CW, Shapiro SS, Whitehurst D, et al. The natural history of factor VIII:C inhibitors in patients with hemophilia A: a national cooperative study. II. Observations on the initial development of factor VIII:C inhibitors. *Blood* 1988;71(2):344-8.
- 59 Sharathkumar A, Lillicrap D, Blanchette VS, et al. Intensive exposure to factor VIII is a risk factor for inhibitor development in mild hemophilia A. *J Thromb Haemost* 2003;1(6):1228-36.
- 60 Teitel JM, Carcao M, Lillicrap D, et al. Orthopaedic surgery in haemophilia patients with inhibitors: a practical guide to haemostatic, surgical and rehabilitative care. *Haemophilia* 2009; 15(1):227-39.
- 61 Meijer P, Verbruggen B. The between-laboratory variation of factor VIII inhibitor testing: the experience of the external quality assessment program of the ECAT foundation. *Semin Thromb Hemost* 2009;35(8):786-93.
- 62 Verbruggen B, van Heerde WL, Laros-van Gorkom BA. Improvements in factor VIII inhibitor detection: from Bethesda to Nijmegen. *Semin Thromb Hemost* 2009;35:752-9.
- 63 de Moerloose P, Fischer K, Lambert T, Windyga J, Batorova A, Lavigne-Lissalde G, Rocino A, Astermark J, Hermans C. Recommendations for assessment, monitoring and follow-up of patients with haemophilia. *Haemophilia* 2012 May; 18(3): 319-25.
- 64 Rixubis [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; September 2014
- 65 Alprolix [package insert]. Cambridge, MA: Biogen Idec, Inc.; March 2014.
- 66 Elocate [package insert]. Cambridge, MA: Biogen Idec, Inc.; June 2014.
- 67 Ixinity [package insert]. Baltimore, MD: Emergent BioSolutions, Inc.; April 2015
- 68 WFH Guidelines for the Management of Hemophilia. Available at: <http://www.wfh.org/en/resources/wfh-treatment-guidelines>. Accessed September 19, 2015.
- 69 MASAC recommendation concerning prophylaxis (regular administration of clotting factor concentrate to prevent bleeding). MASAC document #179. Available at: <https://www.hemophilia.org/Researchers-Healthcare-Providers/Medical-and-Scientific-Advisory-Council-MASAC/MASAC-Recommendations/MASAC-Recommendation-Concerning-Prophylaxis>. Accessed September 19, 2015.
- 70 The Hemophilia, von Willebrand Disease & Platelet Disorders Handbook. Available at: <http://www.hog.org/handbook>. Accessed September 19, 2015.
- 71 Fischer K, Van Der Bom JG, Prejs R, et al. Discontinuation of prophylactic therapy in severe haemophilia: incidence and effects on outcome. *Haemophilia* 2001;7(6):544-50.
- 72 Hay CR, Brown S, Collins PW, Keeling DM, Liesner R. The diagnosis and management of factor VIII and IX inhibitors: a guideline from the United Kingdom Haemophilia Centre Doctors Organisation. *Br J Haematol* 2006;133:591-605.
- 73 McMillan CW, Shapiro SS, Whitehurst D, et al. The natural history of factor VIII:C inhibitors in patients with hemophilia A: a national cooperative study. II. Observations on the initial development of factor VIII:C inhibitors. *Blood* 1988;71(2):344-8.
- 74 Teitel JM, Berntorp E, Collins P, et al. A systematic approach to controlling problem bleeds in patients with severe congenital haemophilia A and high-titre inhibitors. *Haemophilia* 2007;13: 256-63.
- 75 Astermark J, Donfield SM, DiMichele DM, et al. A randomized comparison of bypassing agents in hemophilia complicated by an inhibitor: the FEIBA Novoseven Comparative (FENOC) Study. *Blood* 2007;109(2):546-51.
- 76 Berntorp E, Shapiro A, Astermark J, et al. Inhibitor treatment in haemophilias A and B: summary statement for the 2006 international consensus conference. *Haemophilia* 2006;12(Suppl 6):1-7.
- 77 Hay CR, Dimichele DM. The principal results of the International Immune Tolerance Study: a randomized dose comparison. *Blood* 2012;119:1335-1344.

- 
- 78 Beutel K, Hauch H, Rischewski J, Kordes U, Schneppenheim J, Schneppenheim R. ITI with high-dose FIX and combined immunosuppressive therapy in a patient with severe haemophilia B and inhibitor. *Hamostaseologie* 2009 May;29(2):155-7.
- 79 Advate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2014.
- 80 Alphanate [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; March 2015.
- 81 Eloctate [package insert]. Cambridge, MA: Biogen Idec, Inc.; June 2014
- 82 Helixate FS [package insert]. Kankakee, IL: CSL Behring, LLC; May 2014.
- 83 Hemofil M [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2012.
- 84 Humate-P [package insert]. Kankakee, IL: CSL Behring, LLC; June 2014.
- 85 Koate-DVI [package insert]. Fort Lee, NJ: Kendrion Biopharma, Inc.; August 2012.
- 86 Monoclate-P [package insert]. Kankakee, IL: CSL Behring, LLC; February 2014.
- 87 Novoeight [package insert]. Plainsboro, NJ: Novo Nordisk, Inc.; March 2015
- 88 Obizur [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; October 2014
- 89 Recombinate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; December 2010.
- 90 Wilate [package insert]. Hoboken, NJ: Octapharma USA; August 2015.
- 91 Xyntha [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 92 Xyntha Solofuse [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 93 AlphaNine SD [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; January 2013.
- 94 Alprolix [package insert]. Cambridge, MA: Biogen Idec, Inc.; March 2014.
- 95 Bebulin [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; July 2012.
- 96 BeneFIX [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; August 2015.
- 97 Ixinity [package insert]. Baltimore, MD: Emergent BioSolutions, Inc.; April 2015
- 98 Mononine [package insert]. Kankakee, IL: CSL Behring, LLC; February 2013.
- 99 Profilnine [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; May 2014
- 100 Rixubis [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; September 2014
- 101 Feiba NF [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; November 2013.
- 102 NovoSeven RT [package insert]. Plainsboro NJ: Novo Nordisk, Inc.; June 2015.
- 103 Calculating the Dose The Hemophilia, von Willebrand Disease & Platelet Disorders Handbook. Available at: <http://www.hog.org/handbook/article/3/29/calculating-the-dose>. Accessed September 19, 2015.
- 104 Advate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2014.
- 105 Alphanate [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; March 2015.
- 106 Eloctate [package insert]. Cambridge, MA: Biogen Idec, Inc.; June 2014
- 107 Helixate FS [package insert]. Kankakee, IL: CSL Behring, LLC; May 2014.
- 108 Hemofil M [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; April 2012.
- 109 Humate-P [package insert]. Kankakee, IL: CSL Behring, LLC; June 2014.
- 110 Koate-DVI [package insert]. Fort Lee, NJ: Kendrion Biopharma, Inc.; August 2012.
- 111 Kogenate FS [package insert]. Whippany, NJ: Bayer Healthcare LLC; August 2015.
- 112 Kogenate FS with Bio-Set [package insert]. Whippany, NJ: Bayer Healthcare LLC; August 2015.
- 113 Monoclate-P [package insert]. Kankakee, IL: CSL Behring, LLC; February 2014.
- 114 Novoeight [package insert]. Plainsboro, NJ: Novo Nordisk, Inc.; March 2015
- 115 Obizur [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; October 2014
- 116 Recombinate [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; December 2010.
- 117 Wilate [package insert]. Hoboken, NJ: Octapharma USA; August 2015.
- 118 Xyntha [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 119 Xyntha Solofuse [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; October 2014.
- 120 AlphaNine SD [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; January 2013.
- 121 Alprolix [package insert]. Cambridge, MA: Biogen Idec, Inc.; March 2014.
- 122 Bebulin [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; July 2012.
- 123 BeneFIX [package insert]. Philadelphia, PA: Wyeth Pharmaceuticals, Inc; August 2015.
- 124 Ixinity [package insert]. Baltimore, MD: Emergent BioSolutions, Inc.; April 2015
- 125 Mononine [package insert]. Kankakee, IL: CSL Behring, LLC; February 2013.
- 126 Profilnine [package insert]. Los Angeles, CA: Grifols Biologics, Inc.; May 2014.
- 127 Rixubis [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; September 2014
- 128 Feiba NF [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; November 2013.
- 129 NovoSeven RT [package insert]. Plainsboro NJ: Novo Nordisk, Inc.; June 2015.