



Sedative / Hypnotic Therapeutic Class Review (TCR)

December 14, 2018

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Magellan Rx Management.

All requests for permission should be mailed to:

Magellan Rx Management
Attention: Legal Department
6950 Columbia Gateway Drive
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to PSTCREditor@magellanhealth.com.

December 2018

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
© 2004-2018 Magellan Rx Management. All Rights Reserved.

MagellanRx
MANAGEMENTSM

FDA-APPROVED INDICATIONS

Drug	Manufacturer	Short-term Treatment of Insomnia	Treatment of Insomnia	Treatment of Non-24-hour Sleep-wake Disorder
doxepin (Silenor®) ¹	Pernix		X	
estazolam ²	generic	X		
eszopiclone (Lunesta®) ³	generic, Sunovion		X	
flurazepam ⁴	generic	X		
quazepam (Doral®) ⁵	generic, Galt	X		
ramelteon (Rozerem®) ⁶	Takeda		X	
suvorexant (Belsomra®) ⁷	Merck		X	
tasimelteon (Hetlioz®) ⁸	Vanda			X
temazepam (Restoril®) ⁹	generic, Mallinckrodt	X		
triazolam (Halcion®) ¹⁰	generic, Pfizer	X		
zaleplon (Sonata®) ¹¹	generic, Pfizer	X		
zolpidem (Ambien®) ¹²	generic, Sanofi Aventis	X		
zolpidem sublingual (Edluar®) ¹³	Meda/Mylan	X		
zolpidem sublingual (Intermezzo®) ¹⁴	generic, Purdue		X Middle of the night awakening	
zolpidem (Zolpimist®) ¹⁵	Magna	X		
zolpidem ER (Ambien® CR) ¹⁶	generic, Sanofi Aventis		X	

ER = extended-release

- Eszopiclone (Lunesta), suvorexant (Belsomra), zaleplon (Sonata), zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) and the benzodiazepines (estazolam, flurazepam, quazepam [Doral], temazepam [Restoril], and triazolam [Halcion]) are Scheduled IV controlled substances.
- Doxepin (Silenor), ramelteon (Rozerem), tasimelteon (Hetlioz) are not controlled substances.

OVERVIEW

Insomnia is a symptom complex that comprises difficulties falling asleep, staying asleep, or non-refreshing sleep in combination with daytime dysfunction or distress. The symptom complex can be an independent disorder (primary insomnia) or the result of another condition (secondary insomnia).¹⁷ Insomnia is commonly divided into 3 types based on duration. Transient insomnia lasts up to 1 week and is often referred to as adjustment sleep disorder because it is caused most often by an acute situational stress, such as a test or deadline. It is often recurrent with the same or similar stresses. The second type, short-term insomnia, by definition lasts 1 to 6 months and is usually associated with more persistent stressful situational (death or illness) or environmental (noise) factors. Finally, chronic insomnia is insomnia lasting more than 6 months.

Treatment for insomnia should first consist of identification and treatment and/or control of secondary sources. Whenever possible, non-pharmacological measures should be used to treat insomnia. When such measures fail to address the condition, use of pharmacologic hypnotics may be necessary.¹⁸ The 2017 American Academy of Sleep Medicine (AASM) guidelines recommend psychological and behavioral strategies, which are effective in both primary and secondary insomnia, as are pharmacological interventions.¹⁹ The guidelines recommend that initial behavioral interventions should include stimulus control therapy or relaxation therapy, or a combination of therapies referred to as cognitive behavioral therapy for insomnia (CBT-I). Cognitive behavioral therapy for insomnia includes traditional cognitive behavioral therapy (CBT), stimulus control, and sleep restriction therapy (with or without relaxation therapy). Paradoxical intention and biofeedback therapy may also be effective. Behavioral therapy should always include good sleep hygiene in combination with other therapies. Additionally, the AASM guideline recommends that pharmacotherapy should be used to treat patients who failed to respond to CBT (Grade: weak recommendation, low-quality evidence). AASM recommends zaleplon, triazolam, and ramelteon versus no treatment for sleep onset insomnia (weak recommendations), suvorexant and doxepin over no treatment for sleep maintenance insomnia (weak recommendations), and eszopiclone, zolpidem, and temazepam for both sleep onset and sleep maintenance insomnia. The AASM guidelines recommend against the use of trazodone or tiagabine for sleep onset or sleep maintenance insomnia in adults (Grade: weak recommendation, low-quality evidence). AASM recommends against the use of over-the-counter (OTC) medications or supplements (e.g., diphenhydramine, tryptophan, melatonin) or herbal products (e.g., valerian) as a treatment for sleep onset and sleep maintenance for chronic insomnia (Grade: weak recommendation, low-quality evidence). Choice of agent should be based on symptom pattern, treatment goals, past treatment response, patient preference, cost, availability of other treatments options, comorbid conditions, contraindications, potential interaction with concurrent medication, and side effects.

The American College of Physicians (ACP) released 2016 clinical practice guidelines on the management of chronic insomnia disorder in adults.²⁰ Recommendations include CBT-I as the initial treatment for chronic insomnia disorder (Grade: strong recommendation, moderate-quality evidence). Clinicians should use a shared decision-making approach with the patient with a discussion of benefits, harms, costs of short-term use of medications when deciding whether to add pharmacological therapy in adults with chronic insomnia in whom CBT-I alone was not successful (Grade: weak recommendation, low-quality evidence). The ACP guidelines state that the FDA has approved medications for short-term use (4 to 5 weeks), and patients should not continue taking them for

extended periods of time. Patients should be further evaluated if insomnia does not remit within 7 to 10 days of treatment.

The incidence of insomnia in children ranges from 1% to 6%; in children with neurodevelopmental or psychiatric comorbidities, the incidence is as high as 50% to 75%.^{21,22,23} Insomnia in children may result in irritability, restlessness, lack of concentration, suicide risk, and poor memory.²⁴ The American Academy of Sleep Medicine's task force on Pharmacotherapy in Pediatric Sleep Medicine published guidelines in 2005 that do not recommend any one hypnotic over another for use in children.²⁵ Rather, the consensus statement urges caution when using any of these drugs for the pediatric patient and calls for additional research to be completed.

Guidelines from the 2006 National Sleep Foundation state that there is a need for pharmacologic management of pediatric insomnia.²⁶ Acknowledging that there is an absence of pharmaceuticals indicated for hypnotic use in the pediatric population, this organization stated that there is a need for trials to confirm the safety and efficacy of such agents in these patients.

Non-24-hour sleep-wake disorder (N24SWD or non-24) is a chronic circadian rhythm disorder that causes problems with the timing of sleep and sleep patterns.²⁷ It occurs in approximately 55% to 70% of people who are completely blind, but can also be experienced in sighted people; prevalence among people with sight is unknown. The National Organization for Rare Disorders (NORD) states that the condition is characterized by the failure of a person's biological clock to synchronize to a 24-hour day light-dark cycle. In people who are completely blind (e.g., have no perception of light), this is due to their eyes inability to register light signals. In sighted people N24SWD may be due to a number of factors, such as altered sensitivity of light on circadian rhythm; self-selected changes in light exposure late in the day; and hormonal factors. Those with the disorder may have difficulty falling or staying asleep, and may wake up feeling as if they need more rest. People with N24SWD may find their sleep patterns reversed (e.g., needing to sleep during the day and to be awake at night). N24SWD onset most often occurs in late teen or early twenties but can occur at any age and appears to be a life-long effect.

In 2015, the AASM updated their guidelines for the treatment of intrinsic circadian rhythm sleep-wake disorders, which includes N24SWD.²⁸ They endorse strategically-timed melatonin or light therapy for select patients with Circadian Rhythm Sleep-Wake Disorders (CRSWD), including N24SWD. The AASM Task Force also strongly recommends avoiding the use of sleep-promoting medications to treat elderly patients with dementia and Irregular Sleep-Wake Rhythm Disorder (ISWRD). Tasimelteon (Hetlioz), FDA approved in October 2014 to treat N24SWD, was not addressed in the guidelines.

This review will focus on those agents indicated for sleep disorders and contemporary treatment in adults.

PHARMACOLOGY

Benzodiazepines are believed to potentiate gamma aminobutyric acid (GABA) neuronal inhibition. The sedative and anticonvulsant actions of these drugs involve GABA receptors located in the limbic, neocortical, and mesencephalic reticular systems. At least 2 benzodiazepine receptor subtypes have been identified in the brain, BZ-1 and BZ-2. BZ-1 is thought to be associated with sleep mechanisms while BZ-2 is thought to be associated with memory, motor, sensory, and cognitive functions. Benzodiazepines generally decrease the time to onset of persistent sleep (sleep onset latency, SOL)

and reduce the number of awakenings.²⁹ Benzodiazepines in this class review include estazolam, flurazepam, quazepam, temazepam, and triazolam.

Although structurally different from the benzodiazepines and from one another, the cyclopyrrolone hypnotics (Z-drugs), eszopiclone (Lunesta), zaleplon (Sonata), and zolpidem (Ambien, Ambien CR, Edluar, Zolpimist, Intermezzo), are all active at the GABA-BZ receptor complex.³⁰ Unlike the benzodiazepines, these agents bind selectively to the BZ-1 receptor.

Ramelteon (Rozerem) is a highly selective and potent agonist of the MT1 and MT2 melatonin receptors, which are believed to be involved in the regulation of the circadian rhythm.³¹ Tasimelteon (Hetlioz) is also an agonist at the MT1 and MT2 receptors, with greater affinity for the MT2 receptor.³² The MT1 receptor is believed to regulate sleepiness, whereas the MT2 receptor is thought to help the body shift between day and night. Ramelteon has been reported to have greater affinity, selectivity, and potency than melatonin for the MT1 receptor, resulting in a better ability to induce sleep onset. Ramelteon has shown no affinity for the GABA-receptor complex, which is the primary target area for most of the other agents in this class.³³

Doxepin (Silenor) is a new formulation of the sedative tricyclic antidepressant that is approved for treatment of insomnia, in particular in patients with sleep maintenance problems. Doxepin binds with a high affinity to H₁ histamine receptors where it functions as an antagonist and may result in its sedative effect.³⁴

Suvorexant (Belsomra) is an orexin receptor antagonist. The orexin neuropeptide signaling system is a central promoter of wakefulness. Blocking the orexin receptor suppresses wakefulness.³⁵

PHARMACOKINETICS^{36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51}

Drug	Onset of Action (minutes)	Duration of Action (hours)	Half-Life of Parent Compound (hours)	Active Metabolite(s) (Half-Life)	Metabolism
doxepin (Silenor)	n/a	3 (delayed in presence of high fat meal)	15.3	N-desmethyldoxepin	oxidation and demethylation
estazolam ⁵²	15-120	6-8	14.4-15	None	CYP 3A4
eszopiclone (Lunesta) ⁵³	≤ 30	6-8	6	(S)-zopiclone-N-oxide (S)-N-desmethylzopiclone	CYP 3A4 CYP 2E1
flurazepam	30-60	7-10	2.3	N ₁ -desalkylflurazepam (47-100 hrs; up to 160 hrs in elderly) N ₁ -hydroxyethylflurazepam (2-4 hrs)	oxidation
quazepam (Doral)	20-60	7-10	39	2-oxoquazepam (39 hrs) N-desalkyl-2-oxoquazepam (73 hrs)	hepatic
ramelteon (Rozerem)	≤ 30	nr	1-2.6	MII (2-5 hrs)	oxidation CYP 1A2

Pharmacokinetics (continued)

Drug	Onset of Action (minutes)	Duration of Action (hours)	Half-Life of Parent Compound (hours)	Active Metabolite(s) (Half-Life)	Metabolism
suvorexant (Belsomra)	30 mins to 6 hrs (delayed by 1.5 hrs in presence of food)	nr	12	None	CYP3A (major) CYP2C19 (minor)
tasimelteon (Hetlioz)	n/a	nr	1.3±0.4	None	oxidation and oxidative dealkylation
temazepam (Restoril)	15-120	6-8	3.5-18.4	None	conjugation
triazolam (Halcion)	15-30	1.7-3	1.5-5.5	None	oxidation CYP 3A4
zaleplon (Sonata) ⁵⁴	10-30	4	1	None	aldehyde oxidase CYP 3A4
zolpidem (Ambien) zolpidem ER (Ambien CR)	<30-96 (delayed in presence of food)	8	2.5-2.6 2.8	None	CYP 3A4
zolpidem sublingual (Edluar)	delayed in presence of food	8	2.65-2.85	None	CYP 3A4
zolpidem sublingual (Intermezzo) ⁵⁵	35 (delayed in presence of food)	4	2.5	None	CYP 3A4

hrs = hours

Tasimelteon (Hetlioz) effects may not occur for weeks or months due to of individual differences in circadian rhythms.

Zolpidem ER (Ambien CR) is a coated 2-layer tablet with 1 layer that releases the drug content immediately and another layer that slowly releases additional drug beyond 3 hours after administration. Compared to the immediate-release formulation (Ambien), the peak concentration of zolpidem ER is reached at a later time (2.4 versus 2 hours; $p < 0.004$) and is approximately 13% lower.⁵⁶

Zolpidem (Zolpimist) is bioequivalent to zolpidem (Ambien) tablets.

The difference in zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, and Zolpimist) dosing between men and women is due to the lower rate of clearance by women compared to men.

CONTRAINDICATIONS/WARNINGS^{57,58,59,60,61,62,63,64,65,66,67,68,69,70,71}

All agents in this class, except doxepin (Silenor), suvorexant (Belsomra), and tasimelteon (Hetlioz), have a warning regarding their potential for anaphylaxis and angioedema that can occur as early as the first dose.

These drugs, with the exception of zolpidem SL (Intermezzo), should be administered immediately before going to bed or once the patient experiences difficulty falling asleep. Zolpidem SL (Intermezzo) should be utilized for middle of the night awakenings when the patient has more than 4 hours before planned waking time.

All of these agents may result in short-term memory impairment, hallucinations, impaired coordination, dizziness, and light-headedness. The FDA-approved hypnotics, including non-benzodiazepine sedative hypnotics, hypnotics, and doxepin, carry a warning regarding complex sleep-related behaviors such as sleep-driving, making phone calls, sexual activity, and preparing and eating food while asleep; often patients have no memory of these events. These behaviors are more likely to occur when the sedative-hypnotic is taken concurrently with alcohol or other central nervous system (CNS) depressants.

All drugs in this class should be used at the lowest effective dose. In January 2013, the FDA announced lower dosing recommendations for zolpidem due to new data showing that blood levels in some patients may be high enough the morning after use to impair activities that require alertness, including driving.⁷² Women appear to be more susceptible to this risk as they eliminate zolpidem more slowly compared to men. The FDA lowered the recommended dose of zolpidem for women from 10 mg to 5 mg for immediate-release products (Ambien, Edluar, and Zolpimist) and from 12.5 mg to 6.25 mg for extended-release products (Ambien CR). Prescribers should consider lower zolpidem doses for men, but it is not required. At the time of Intermezzo's approval, the label already recommended a lower dosage for women than for men. For zolpidem and other insomnia drugs, the lowest dose should be prescribed that treats the patients' symptoms.

All sedative/hypnotics should be administered with caution to patients exhibiting signs and symptoms of depression. Suicidal tendencies may be present in such patients, and protective measures may be required. Intentional overdose is more common in this group of patients; therefore, the smallest amount of drug that is feasible should be prescribed for the patient at any one time. Special attention should be given to doxepin (Silenor), because the active ingredient is a tricyclic antidepressant. Administration of antidepressants in children, adolescents, and young adults with major depressive disorder (MDD) and other psychiatric disorders could also increase suicidal thoughts and actions

All drugs in this class should be used only after careful assessment of the cause of sleep, behavioral changes, amnesia, withdrawal symptoms, and emergence or worsening of psychiatric or physical disorders. Patients whose insomnia fails to remit after 7 to 10 days of treatment with a sedative-hypnotic may have a primary psychiatric or medical illness that should be evaluated.

In 2016, the FDA informed healthcare professionals that concurrent use of opioids and benzodiazepines or other CNS depressants has resulted in serious adverse reactions such as profound sedation, respiratory depression, coma, and death.⁷³ Providers should limit prescribing of opioids with benzodiazepines (e.g., estazolam, flurazepam, quazepam, temazepam, triazolam) to patients without alternative treatment options. If used together, dosages and duration of therapy should be minimized and patients should be monitored for signs and symptoms of respiratory depression and sedation. Boxed warnings were added to the labeling of all prescription opioids, including those indicated for pain and for cough, and benzodiazepines. In 2017, the safety warning was updated to urge prescribers to consider the risks and benefits for the use of opioid addiction medications such as buprenorphine and methadone in combination with benzodiazepines. Careful medication management is recommended over withholding opioid addiction medications for patients using benzodiazepines.⁷⁴

All sedative hypnotics can cause drowsiness and a decreased level of consciousness, which may lead to falls and, consequently, to severe injuries such as hip fractures and intracranial hemorrhage. This risk is especially concerning in the elderly. In addition, a meta-analysis of sleep medications in older adults reviewed the risk for hip fracture during short-term and long-term use of benzodiazepines and non-benzodiazepines.⁷⁵ The short-term use (≤ 14 days) was associated with the highest risk (RR, 2.4; [95% confidence interval, 1.88 to 3.05]; and RR, 2.39 [95% confidence interval, 1.74 to 3.29]). In the absence of direct clinical trials comparing agents in this class and the increased risk of falls and hip fractures, benzodiazepines and non-benzodiazepines should be prescribed with equal caution for short-term use in older adults.

Suvorexant is contraindicated in patients with narcolepsy.

Benzodiazepines, including estazolam, flurazepam, quazepam, temazepam, and triazolam, are contraindicated in patients with suspected or established sleep apnea. Doxepin, suvorexant, or tasimelteon have not been studied in subjects with severe sleep apnea or severe chronic obstructive pulmonary disease (COPD) and they are not recommended for use in these populations. Single doses of ramelteon did not result in respiratory depression in patients with mild to severe COPD or in patients with mild to moderate sleep apnea; there is no data available on the effect of multiple doses in patients with either condition. Ramelteon has not been studied in patients with severe obstructive sleep apnea; therefore, it is not recommended in this patient population.

Patients with compromised respiratory function should be cautioned prior to the use of zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) since respiratory depression can occur.

Eszopiclone (Lunesta), suvorexant (Belsomra), zaleplon (Sonata), zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) and benzodiazepines (estazolam, flurazepam, quazepam [Doral], temazepam [Restoril], and triazolam [Halcion]) are Scheduled IV controlled substances. Patients with a history of addiction to, or abuse of, drugs or alcohol are at increased risk for misuse, abuse, and addiction with these agents; patients should be monitored carefully when taking these agents. Doxepin (Silenor), ramelteon (Rozerem), and tasimelteon (Hetlioz) are not controlled substances and are not associated with abuse or physical dependence.

Patients with untreated narrow angle glaucoma or severe urinary retention should not use doxepin (Silenor).

Sleep paralysis, an inability to move or speak for up to several minutes during sleep-wake transitions, hypnagogic/hypnopompic hallucinations, and cataplexy-like symptoms can occur with suvorexant.

DRUG INTERACTIONS^{76,77,78,79,80,81,82,83,84,85,86,87,88,89,90}

Drugs in this class should be used with caution in patients receiving other CNS depressants as the effects may be additive, resulting in decreased alertness and impaired psychomotor performance.

Increased CNS depressant effects of the benzodiazepines that are metabolized by oxidation have been reported when co-administered with isoniazid, oral contraceptives, cimetidine, and disulfiram.

Estazolam, eszopiclone (Lunesta), ramelteon (Rozerem) and its active MII metabolite, tasimelteon (Hetlioz), triazolam, zaleplon (Sonata), and zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) are substrates for the CYP450 3A4 enzyme. As such, inducers of CYP450 3A4 (e.g., rifampin) increase the clearance and reduce the bioavailability of these agents by approximately 80%. Inhibitors

of CYP450 3A4 (e.g., cimetidine, clarithromycin, ketoconazole) increase the bioavailability of these drugs by up to 84%.

Doxepin (Silenor) is primarily metabolized by CYP2C19 and CYP2D6 hepatic cytochrome P450 isozymes. Inhibitors of these isozymes may increase the exposure of doxepin in patients. Concomitant administration of doxepin with CNS depressants, including alcohol, and sedative antihistamines has shown increased sedative effects. Severe hypoglycemia has been reported with the simultaneous use of tolazamide. Using doxepin and cimetidine together has caused an increased exposure to doxepin. Patients should not use monoamine oxidase inhibitor (MAOI) medications within 14 days of doxepin.

Ramelteon (Rozerem) is contraindicated for concomitant use with fluvoxamine, a strong CYP1A2 inhibitor. It should be used with caution in patients taking less strong CYP1A2 inhibitors. Administration of ramelteon with fluconazole increases the bioavailability of ramelteon and the MII metabolite by approximately 150%.

Suvorexant (Belsomra) is not recommended with strong inhibitors of CYP450 3A (e.g., ketoconazole, itraconazole, posaconazole, clarithromycin, nefazodone, ritonavir, saquinavir, nelfinavir, indinavir, boceprevir, telaprevir, telithromycin, and conivaptan). The recommended dose in patients receiving moderate CYP450 3A inhibitors (amprenavir, aprepitant, atazanavir, ciprofloxacin, diltiazem, erythromycin, fluconazole, fosamprenavir, grapefruit juice, imatinib, verapamil) is 5 mg; the dosage may be increased to 10 mg if needed for efficacy. CYP3A inducers (e.g., rifampin, carbamazepine, and phenytoin) can substantially decrease suvorexant exposure thereby minimizing the efficacy.

Concurrent administration of triazolam with efavirenz, delavirdine, azole antifungals, nefazodone, protease inhibitors, and any drug that significantly impairs the CYP3A mediated oxidative metabolism are contraindicated. Caution is recommended when administering triazolam with grapefruit juice, fluvoxamine, diltiazem, verapamil, amiodarone, nifedipine, nifedipine, cimetidine, and ranitidine because these agents can increase plasma concentration of triazolam.

The efficacy of tasimelteon (Hetlioz) may be reduced in patients on concomitant beta adrenergic receptor antagonists.

The concurrent use of a CYP3A4 inducer (e.g., St. John's wort, rifampin) with zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) is not recommended as it may reduce the blood levels of zolpidem.

ADVERSE EFFECTS^{91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106}

Drug	Headache	Myalgia	Amnesia	Dizziness	Daytime Drowsiness/ Somnolence
doxepin (Silenor)	Nr	<2	nr	reported	6-9
estazolam	16 (27)	≤ 1	≤ 1	7 (3)	3 (2)
eszopiclone (Lunesta)	16-20 (13)	nr	nr	5-7 (4)	8-10 (3)
flurazepam	reported	nr	reported	reported	reported
quazepam (Doral)	4.5 (2.2)	nr	reported	1.5 (< 1)	12 (3.3)

Adverse Effects (continued)

Drug	Headache	Myalgia	Amnesia	Dizziness	Daytime Drowsiness/ Somnolence
ramelteon (Rozerem)	< 1	nr	nr	4 (3)	3 (2)
suvorexant (Belsomra)	7 (6)	nr	reported	3 (2)	7 (3)
tasimelteon (Hetlioz)	17 (7)	nr	nr	nr	nr
temazepam (Restoril)	8.5 (9.1)	nr	< 0.5	4.5 (3.3)	2.5 (1.1)
triazolam (Halcion)	9.7 (8.4)	nr	reported	7.8 (3.1)	nr
zaleplon (Sonata)	30-42 (35)	7 (4)	2-4 (1)	7-9 (7)	nr
zolpidem (Ambien, Edluar, Zolpimist)	7 (6)	> 1	1 (0)	5 (1)	8 (6)
zolpidem (Intermezzo)	3 (1)	nr	nr	nr	nr
zolpidem ER (Ambien CR)	14-19 (11-16)	< 1-4 (0)	1-3 (0)	8-12 (3-5)	6-15 (2-5)

Adverse effects are reported as a percentage. Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. Incidences for the placebo group are indicated in parentheses. nr = not reported.

For eszopiclone (Lunesta), 16% to 34% of patients reported a dose-related unpleasant taste as compared to 3% of placebo patients. Dose-related respiratory infection has been reported in 5% to 10% of patients taking eszopiclone compared to 3% of patients taking placebo. Anxiety has been reported in 1% to 3.7% of patients receiving eszopiclone compared to 0 patients taking placebo in one 6-week placebo controlled trial.

Nightmares have been reported with the use of tasimelteon (Hetlioz) and zaleplon (Sonata) and abnormal dreams with suvorexant (Belsomra). Additionally, postmarketing reports for suvorexant (Belsomra) have identified several possible adverse reactions, including palpitations, tachycardia, psychomotor hyperactivity, and anxiety.

SPECIAL POPULATIONS^{107,108,109,110,111,112,113,114,115,116,117,118,119}

Pediatrics

Safety and effectiveness of agents in this class review have not been established in pediatric patients.

In an 8-week study, in patients aged 6 to 17 years with insomnia associated with attention-deficit/hyperactivity disorder (ADHD) zolpidem oral solution at a dose of 0.25 mg/kg at bedtime did not decrease sleep latency compared to placebo. The most common treatment-related adverse effects reported were (> 5%) with zolpidem versus placebo and included dizziness (23.5% versus 1.5%), headache (12.5% versus 9.2%), and hallucinations (7% versus 0%).

Eszopiclone (Lunesta) failed to demonstrate efficacy in controlled clinical studies of pediatric patients with Attention Deficit/Hyperactivity (ADHD)-associated insomnia.

A survey of 671 primary care pediatricians found that more than 75% had prescribed nonprescription medications, and more than half had prescribed prescription medications for pediatric insomnia.¹²⁰ Most commonly, these agents were prescribed for acute pain and travel, followed by children with special needs. Antihistamines were the most common nonprescription medications for sleep, followed by melatonin and herbal remedies. Alpha-agonists were the most frequently prescribed prescription sleep medication.

Pregnancy

Eszopiclone (Lunesta), ramelteon (Rozerem), and zolpidem (Ambien, Ambien CR, Edluar) were previously Pregnancy Category C and have now been updated to contain descriptive text in compliance with the Pregnancy and Lactation Labeling Rule (PLLR). Available data for these agents are insufficient to identify a risk of major birth defects or miscarriage due to exposure to these drugs. There have been reports of respiratory depression and sedation in neonates born to mothers using zolpidem during the third trimester of pregnancy.

Doxepin (Silenor), eszopiclone (Lunesta), suvorexant (Belsomra), tasimelteon (Hetlioz), zaleplon (Sonata), and zolpidem SL (Intermezzo, Zolpimist) are Pregnancy Category C. While the other benzodiazepine hypnotics are Pregnancy Category X, flurazepam has not been assigned a pregnancy category but is contraindicated during pregnancy.

Hepatic Impairment

Patients with hepatic impairment may display higher doxepin (Silenor) concentrations than healthy patients. Patients with hepatic impairment should initiate treatment at the lowest recommended daily dosage and monitor for adverse daytime effects.

In patients with severe hepatic impairment, the bioavailability of eszopiclone is increased 2-fold compared with healthy volunteers; time-to-peak and peak concentrations remain unchanged.

Exposure to ramelteon was increased nearly 4 times normal amounts when given to patients with mild hepatic impairment and 10 times normal amounts in those with moderate impairment. Ramelteon should be used with caution in patients with moderate hepatic impairment and should not be used in those with severe impairment.

Dose adjustment of suvorexant (Belsomra) is not required in patients with mild or moderate hepatic impairment. Suvorexant is not recommended in patients with severe hepatic impairment.

Dose adjustment of tasimelteon (Hetlioz) is not necessary in patients with mild or moderate hepatic impairment. Tasimelteon has not been studied in patients with severe hepatic impairment (Child-Pugh Class C). Therefore, tasimelteon is not recommended for use in patients with severe hepatic impairment.

The bioavailability of zaleplon is increased up to 400% in patients with compensated cirrhosis and 700% with decompensated cirrhosis. Zaleplon is not recommended for use in patients with severe hepatic impairment; dose reductions should be employed with mild to moderate impairment.

In patients receiving zolpidem (Ambien, Edluar, Zolpimist) who have hepatic insufficiency, the daily dose should not exceed 5 mg and, for the controlled-release tablet (Ambien CR), the maximum dose should not exceed 6.25 mg. Ambien CR should be avoided in patients with severe hepatic impairment

as it may precipitate hepatic encephalopathy. For zolpidem sublingual (Intermezzo) in those with hepatic insufficiency, the maximum once daily dose should not exceed 1.75 mg.

Ethnicity

In Japanese adults, the maximum plasma concentration of zaleplon is increased by 37%. The effect of other ethnic groups has not been widely studied.¹²¹

Geriatrics^{122,123,124}

Patients over the age of 65 years may demonstrate an increase in total exposure to sedative/hypnotic agents. Dosing for the benzodiazepines should commonly begin at the lowest effective dose for these patients. Ramelteon and doxepin may be exceptions as they did not show any overall differences in safety and efficacy between elderly and younger adult patients.

Smokers

The clearance of benzodiazepines is accelerated in smokers compared to nonsmokers.

Smoking causes induction of CYP1A2 levels. The exposure of tasimelteon (Hetlioz) in smokers was lower than in non-smokers and, therefore, the efficacy of tasimelteon may be reduced in smokers.

DOSAGES^{125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141}

Drug	Bedtime Dose	Dose Adjustment	Availability
doxepin (Silenor)	6 mg Take 30 minutes before bedtime and not within 3 hours of a meal	Elderly: begin with 3 mg and increase to 6 mg, if required 3 mg if clinically indicated for individual patients	Tablet: 3 mg, 6 mg
estazolam	1 mg to 2 mg	Elderly, underweight, or debilitated: 0.5 mg to 1 mg	Tablet: 1 mg, 2 mg
eszopiclone* (Lunesta)	1 mg	Dosing can be raised to 2 mg or 3 mg if clinically indicated Elderly: total dose should not exceed 2 mg Severe hepatic impairment: should not exceed 2 mg Concurrent use with strong CYP 3A4 inhibitor: should not exceed 2 mg	Tablet: 1 mg, 2 mg, 3 mg
flurazepam	Women [†] : 15 mg Men: either 15 mg or 30 mg	Women: May increase to 30 mg Elderly and debilitated patients: 15 mg	Capsule: 15 mg, 30 mg
quazepam (Doral)	Initial dose: 7.5 mg	May be increased to 15 mg if necessary for efficacy	Tablet: 15 mg
ramelteon (Rozerem) [‡]	8 mg 30 minutes prior to going to bed	--	Tablet: 8 mg
suvorexant (Belsomra)	10 mg within 30 minutes of going to bed, with at least 7 hours remaining before the planned time of awakening	Dose can be increased to 20 mg if the 10 mg dose is well-tolerated but not effective	Tablet: 5 mg, 10 mg, 15 mg, 20 mg
tasimelteon (Hetlioz)	20 mg	--	Capsule: 20 mg
temazepam (Restoril)	7.5 mg to 30 mg	Elderly and debilitated: start at 7.5 mg initially until individual responses are determined	Capsule: 7.5 mg, 15 mg, 22.5 mg, 30 mg
triazolam (Halcion)	0.125 mg to 0.5 mg	Elderly or debilitated: 0.125 mg to 0.25 mg	Tablet: 0.125 mg (generic only), 0.25 mg
zaleplon (Sonata) [§]	10 mg to 20 mg May also use 5 mg for low weight individuals May be taken immediately before bedtime or after the patient has gone to bed and has experienced difficulty falling asleep	Elderly, low-weight, or hepatic impairment: 5 mg to 10 mg Concurrent use of cimetidine: Initial dose: 5 mg Mild to moderate hepatic impairment: 5 mg	Capsule: 5 mg, 10 mg

Dosages (continued)

Drug	Bedtime Dose	Dose Adjustment	Availability
zolpidem* (Ambien, Edluar, Zolpimist)	Men: 5 mg or 10 mg Women: 5 mg Use the lowest dose effective for patient. Should be taken immediately before bedtime with at least 7–8 hours remaining before the planned time of awakening The 5 mg dose can be increased to 10 mg, if needed, but the higher dose is more likely to impair next morning driving and other activities that require full alertness. The total dose should not exceed 10 mg once daily.	Elderly, debilitated, or hepatic insufficiency: 5 mg once daily in both men and women immediately before bedtime Concurrent CNS depressants: dosage adjustment may be necessary	Tablet: 5 mg, 10 mg Sublingual Tablet: 5 mg, 10 mg Oral Spray: 5 mg (30 and 60 metered-doses)
zolpidem ER (Ambien CR)*	Men: 6.25 mg or 12.5 mg Women: 6.25 mg Use the lowest dose effective for patient. Should be taken immediately before bedtime with at least 7 to 8 hours remaining before the planned time of awakening The 6.25 mg dose can be increased to 12.5 mg, if needed, but the higher dose is more likely to impair next morning driving and other activities that require full alertness. The total dose should not exceed 12.5 mg daily.	Elderly, debilitated, or hepatic insufficiency: 6.25 mg once daily in both men and women immediately before bedtime Concurrent CNS depressants: dosage adjustment may be necessary	Tablet: 6.25 mg, 12.5 mg
zolpidem sublingual (Intermezzo)	Men: 3.5 mg Women: 1.75 mg Take if there is more than 4 hours remaining before planned time of waking	Elderly, hepatic insufficiency, concurrent CNS depressants: 1.75 mg	Sublingual Tablets: 1.75 mg, 3.5 mg

* Should be taken on empty stomach to avoid delayed onset of action.

† The recommended initial doses for women and men are different because flurazepam clearance is lower in women.

‡ Due to a 31% increase in bioavailability when given with a high-fat meal, ramelteon (Rozerem) should not be taken with or immediately after such a meal.

§ Taking zaleplon (Sonata) or eszopiclone (Lunesta) with or immediately after a heavy, high-fat meal results in slower absorption and would be expected to reduce its effect on sleep latency.

Since many of the adverse effects to the sedative/hypnotics appear to be dose-related, therapy should usually be initiated with a low dose and then maintained at the lowest effective dose, especially in the elderly.

Continuous use should be avoided; patients should be encouraged to use these medications only when necessary. Use for more than 3 weeks should be avoided and monitored if a longer duration is necessary. These drugs should never be combined with alcohol consumption.

CLINICAL TRIALS

Search Strategies

Articles were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the use of all drugs in this class. Randomized, controlled comparative trials for FDA-approved indications are considered the most relevant in this category. Studies included for analysis in the review were published in English, performed with human participants, and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question, and include follow-up (endpoint assessment) of $\geq 80\%$ of participants entering the investigation. Despite some inherent bias found in all studies including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship and/or funding must be considered, the studies in this review have also been evaluated for validity and importance. Many of the trials with agents in this class were performed in an open-label manner; introduction of bias must be considered when evaluating study findings.

Comparisons of Benzodiazepines

estazolam versus flurazepam

The hypnotic efficacy of estazolam 1 mg and 2 mg was compared to flurazepam 30 mg and placebo in a randomized, double-blind, 7-night study that involved 223 patients with insomnia.¹⁴² On subjective assessments of the patients, no differences were noted between estazolam 2 mg and flurazepam 30 mg on any of 6 sleep parameters. Patients who received estazolam 1 mg rated their sleep significantly better than did patients who were receiving placebo on all parameters except sleep onset latency (SOL). Global evaluation of the physicians indicated significant improvement in quality of sleep, sleep duration, and nocturnal awakenings in all 3 active treatment groups; estazolam 2 mg and flurazepam 30 mg decreased SOL significantly. Adverse events were reported by 54% of patients receiving estazolam 1 mg, 58% of those receiving estazolam 2 mg, and 68% of those receiving flurazepam 30 mg. The incidence of adverse events in the placebo group was 43%.

In a double-blind trial, 229 patients with insomnia were randomized to receive estazolam 2 mg, flurazepam 30 mg, or placebo for 7 consecutive nights.¹⁴³ The analysis of efficacy was based on patients' daily assessments of sleep and investigators' global evaluations. The patient subjective questionnaire indicated that estazolam and flurazepam significantly improved all parameters ($p < 0.05$) as compared to placebo. A marked or moderate improvement in sleep was reported by 81%, 78%, and 36% estazolam, flurazepam, and placebo recipients, respectively. There were no significant differences in hypnotic effect between estazolam and flurazepam. All efficacy parameters of the investigators' global evaluation, except quality of sleep, improved significantly more ($p < 0.05$) for patients receiving estazolam or flurazepam than for those receiving placebo. The percentage of patients reporting any adverse experience was 59% for estazolam, 72% for flurazepam, and 43% for placebo. Somnolence and hypokinesia were the most commonly reported adverse events. An analysis of the global evaluation of side effects showed that flurazepam had a significantly worse side effect profile than estazolam ($p < 0.05$) or placebo ($p = 0.001$).

flurazepam versus quazepam (Doral)

Daytime residual drowsiness and psychomotor performance were assessed for quazepam and flurazepam in 2 randomized, parallel, double-blind studies in insomniacs.¹⁴⁴ In the first study, 17 middle-aged patients took quazepam 15 mg or 30 mg or flurazepam 30 mg nightly for 4 weeks. Subjects were given placebo for 4 nights before and 15 nights after active treatment. In the second study, 48 geriatric patients took quazepam 15 mg, flurazepam 15 mg, or placebo nightly for 1 week. Subjects were given placebo for 1 night before and 7 nights after active treatment. In the first study, flurazepam patients were significantly ($p < 0.05$) sleepier the day after the 7th and 14th treatment nights when compared to baseline, whereas quazepam patients were not. In the second study, flurazepam patients were sleepier in the late afternoon ($p < 0.05$) after the 7th treatment night than were quazepam and placebo patients. There were no significant differences among the groups in the performance test results.

quazepam versus triazolam (Halcion)

In a double-blind study, 45 patients were randomized to receive either quazepam 15 mg to 30 mg (median 15 mg) or triazolam 0.25 mg to 0.5 mg (median 0.25 mg) for 4 weeks.¹⁴⁵ The subjects, who had insomnia based on a mild to moderate generalized anxiety disorder, received placebo for 1 week before and 2 weeks after treatment with active drug. Anxiety improved significantly with both drugs and remained improved throughout the 2-week post-drug placebo phase; quazepam was slightly superior to triazolam. Polysomnography, an objective measure of SOL, demonstrated a shortened sleep onset only after quazepam. Sleep efficiency improved after acute administration of both drugs, but improvement was maintained only by quazepam as tolerance developed to triazolam. Rebound insomnia was observed only in the first post-triazolam placebo night. Subjective sleep quality behaved very similarly to objective sleep efficiency. Awakening quality improved after acute therapy with both drugs. Somatic complaints were reported only with quazepam.

A randomized, double-blind, 3-compartment, parallel-group study comparing quazepam 15 mg, triazolam 0.5 mg, and placebo was conducted in 65 insomniac subjects over 5 weeks.¹⁴⁶ Using sleep questionnaires for evaluation, no differences were noted between quazepam and triazolam on treatment nights. Evidence of carryover effectiveness with quazepam and rebound effects with triazolam was noted on off-treatment nights.

Comparisons of Benzodiazepines and Non-benzodiazepines

temazepam versus zolpidem (Ambien)

A randomized, double-blind trial compared zolpidem 10 mg to temazepam 20 mg with respect to subjective rebound insomnia after cessation of 4 weeks of treatment of 163 patients with chronic insomnia.¹⁴⁷ Both agents improved total sleep time (TST), as well as SOL significantly during the 4 treatment weeks. Prevalence rates for rebound insomnia, defined as a worsening of TST or SOL of more than 40% compared to baseline, were 27% for TST and 53% for SOL in the zolpidem group and 26% and 58%, respectively, in the temazepam group. No significant differences were found between the agents for rebound insomnia, nor with respect to their efficacy or safety.

triazolam versus zaleplon (Sonata)

Zaleplon and triazolam were compared in a double-blind, placebo-controlled trial enrolling 132 patients with primary insomnia.¹⁴⁸ Patients received zaleplon 5 mg or 10 mg, triazolam 0.25 mg, or

placebo for 14 nights. Median SOL was shorter in both zaleplon groups and triazolam group compared to placebo during the first week of therapy, but not during the second week due to a significant placebo effect. The effects of zaleplon on SOL were similar in the first and second weeks. Total sleep time did not differ between zaleplon and placebo groups. Total sleep time was increased during the first week of triazolam treatment, but not the second. On subjective assessment of SOL, zaleplon 10 mg and triazolam were more effective than placebo during the first week. Only zaleplon 10 mg produced lower subjective SOL during the second week. On the last night of assessment, none of the active treatments were judged more effective than placebo. Negative residual morning psychomotor or memory effects were not observed in any treatment group.

triazolam versus zolpidem (Ambien)

In a parallel-group, double-blind, placebo-controlled, polysomnographic study, the possible occurrence of rebound insomnia was evaluated in 24 patients suffering from moderate to severe chronic insomnia. Patients were randomized to either triazolam 0.5 mg, zolpidem 10 mg, or placebo.¹⁴⁹ Treatment duration was 27 nights, followed by 3 placebo-controlled withdrawal nights. Both drugs showed significant efficacy compared to placebo during the active treatment period. A trend toward tolerance was noted in the triazolam group but not in the zolpidem group. The increase in total sleep time in the zolpidem group was accompanied by an increase in the number of sleep cycles. When active treatment was discontinued, clear rebound insomnia was present in the triazolam group while it was not possible to observe any rebound in the placebo and zolpidem groups. Subjective feelings of the patients assessed by means of visual analog scale correlated well with polysomnographic data.

Comparisons of Non-benzodiazepines

zaleplon (Sonata) versus zolpidem (Ambien)

In a double-blind study, 615 patients were randomized to receive zaleplon 5 mg, zaleplon 10 mg, or zaleplon 20 mg, zolpidem 10 mg, or placebo.¹⁵⁰ The 4-phase study consisted of a pre-study washout period (1-3 weeks), a single-blind placebo run-in period (7 nights), a double-blind treatment period (28 nights), and a single-blind placebo run-out period (3 nights). In the 574 patients who completed the study, zolpidem significantly reduced SOL during weeks 1 through 3, as did zaleplon 5 mg. Zaleplon 20 mg and zolpidem 10 mg significantly increased sleep duration during all 4 weeks of the double-blind treatment. No significant differences were observed in number of awakenings between the placebo and active treatment groups during the double-blind treatment periods. Scores for sleep quality were significantly better than placebo during week 1 with zaleplon 10 and 20 mg and for all weeks with zolpidem 10 mg. On the first night after treatment discontinuation, significantly more patients who received zolpidem experienced longer SOL relative to baseline and reported withdrawal effects (depressed mood, pain in muscles, peculiar taste, loss of memory, olfactory sensitivity). The most common adverse event in all treatment groups was headache. There were no significant differences in the frequency of treatment-emergent adverse events among the active treatment groups and the placebo group.

A randomized, double-blind, placebo-controlled, 3-period, crossover design was used to study 37 adults with insomnia who received treatment during an experimental awakening 4 hours after bedtime.¹⁵¹ The study objective was to assess the efficacy of zaleplon 10 mg and zolpidem 10 mg administered during experimental middle-of-the-night awakenings in patients with sleep maintenance insomnia using objective polysomnographic measures and to assess daytime residual sedation 4 to 7 hours after dosing

using sleep latency testing. Latency to persistent sleep and total sleep time before and after awakening were recorded. Compared with placebo, latency to persistent sleep after both zaleplon and zolpidem was shorter and total sleep time after administration of the drugs was longer. Significant differences from placebo were not found with zaleplon in daytime sedation measures. At 4, 5, and 7 hours after zolpidem, sleep onset, measured by sleep latency testing, was shorter than after placebo. Self-reported measures of concentration and alertness and Digit Symbol Substitution Test scores after zolpidem were also lower than placebo. Zaleplon 10 mg and zolpidem 10 mg effectively shorten sleep latency and lengthen sleep duration after dosing when administered during experimental nocturnal awakening. Residual sedation was not detected as little as 4 hours after zaleplon 10 mg, but was detected with zolpidem 10 mg up to 7 hours after treatment.

Placebo-controlled Trials of Non-benzodiazepines

doxepin (Silenor) versus placebo

A randomized, double-blind, parallel-group, placebo-controlled study was conducted in healthy adults with transient insomnia.¹⁵² Subjects received a single nighttime dose of placebo (n = 282) or doxepin 6 mg (n = 283) in a sleep laboratory. Efficacy was evaluated objectively by polysomnography and subjectively by morning questionnaire. The primary endpoint was latency to persistent sleep (LPS). Secondary polysomnography endpoints included wake after sleep onset (WASO), total sleep time (TST), wake time after sleep (WTAS), and sleep efficiency (SE). Doxepin demonstrated statistically significant improvements in LPS (13-minute decrease; $p < 0.0001$), WASO (39 minutes less; $p < 0.0001$), TST (51 minutes more; $p < 0.0001$), WTAS ($p < 0.0001$), overall SE ($p < 0.0001$), and SE in each quarter of the night ($p < 0.0001$), all versus placebo. There was no consistent evidence of residual sedation or minor sleep stage alterations. The incidence of doxepin adverse events was comparable to placebo.

A randomized, double-blind, parallel-group, 5-week placebo-controlled trial was conducted in adults diagnosed with primary insomnia and reported difficulty maintaining sleep (n = 229).¹⁵³ Wake time after sleep onset (WASO) on the first night was identified as the primary outcome measure, and both active doses (3 mg and 6 mg) resulted in statistically significant reductions in WASO ($p < 0.0001$). Subsequent WASO measurements were also statistically significant including on night 15 (3 mg $p = 0.0025$; 6 mg $p = 0.0009$), and night 29 (3 mg $p = 0.00248$; 6 mg $p = 0.0009$). Other secondary endpoints with significant improvement compared to placebo included latency to persistent sleep (3 mg $p = 0.0047$; 6 mg $p = 0.0007$) on night 1 only, and Total Sleep Time on nights 1 (both doses $p < 0.0001$), and 29 (3 mg $p = 0.0261$; 6 mg $p < 0.0001$). Although most measures were statistically significant, 2 measures did not reach significance for the 3 mg capsule. Both Sleep efficiency (SE) on the 29th night and total sleep time on night 15 did not reach statistical significance. In contrast to benzodiazepine and other non-benzodiazepine hypnotics, there was no evidence of rebound insomnia when doxepin was discontinued.

eszopiclone (Lunesta) versus placebo

A double-blind study enrolled 308 patients, 21 to 64 years of age, with primary chronic insomnia.^{154,155} Patients were randomized to receive eszopiclone 2 mg, eszopiclone 3 mg, or placebo for 44 consecutive nights followed by 2 nights of single-blind placebo. Treatment with either dose of eszopiclone resulted in an approximate 45-minute improvement in the primary endpoint of SOL (placebo 58 minutes; $p < 0.001$ for both doses). Eszopiclone also significantly improved the secondary endpoint of sleep efficiency ($p < 0.0001$ for both doses compared to placebo). Another secondary

endpoint, wake time after sleep onset (WASO), was reduced only by the higher dose of eszopiclone (41.2 minutes) compared to placebo (49.1 minutes; $p=0.02$). There was no evidence of tolerance or rebound insomnia after therapy discontinuation. There was no decrement in psychomotor performance relative to baseline, nor was there a difference between eszopiclone and placebo. The most common adverse event related to eszopiclone was unpleasant taste.

A double-blind study randomized 231 patients, 65 to 85 years of age, with chronic insomnia to receive either eszopiclone 1 mg, eszopiclone 2 mg, or placebo nightly for 2 weeks.^{156,157,158} In the study, the higher dose of eszopiclone improved sleep maintenance ($p<0.05$), total sleep time (by 40 minutes; $p<0.001$), quality of sleep ($p<0.001$), depth of sleep ($p<0.002$), and reduced the number of naps (median 0 versus 2, $p<0.05$) compared to placebo. Patients receiving the higher dose also reported significant improvements in daytime alertness, daytime ability to function, sense of well-being, and reduced morning sleepiness ($p<0.05$ for all comparisons to placebo). Both doses of eszopiclone were effective at decreasing SOL ($p<0.004$ compared to placebo) and reducing total nap time ($p<0.05$ compared to placebo).

In a double-blind study, 264 patients, 65 to 85 years of age, with a diagnosis of primary insomnia were randomized to receive eszopiclone 2 mg or placebo nightly for 2 weeks.¹⁵⁹ Compared with placebo, eszopiclone 2 mg significantly reduced objective (polysomnographic) and subjective SOL ([$p<0.0001$ for both measurements] and [$p<0.05$ and $p=0.0019$, respectively]). Subjective improvement was also noted in sleep efficiency ($p<0.04$), total sleep time ($p<0.0001$), and the cumulative number and duration of naps among patients who napped ($p=0.03$) compared to placebo. Eszopiclone also produced improvements in the quality of sleep and in physical functioning. There was no rebound insomnia after treatment withdrawal, and the most common adverse event was unpleasant taste.

In a double-blind study, investigators randomized 545 patients with insomnia and major depressive disorder to receive, in addition to daily fluoxetine, eszopiclone 3 mg or placebo nightly.¹⁶⁰ In the 8-week study, patients treated with eszopiclone showed improvements in the primary endpoint, wake time after sleep onset ($p\leq 0.002$), compared to those receiving placebo. The active treatment was also more effective than placebo in improving the secondary endpoints of SOL ($p\leq 0.0001$) and TST ($p\leq 0.0004$). Patients in the eszopiclone group reported superior subjective improvements in sleep quality ($p\leq 0.0002$), depth of sleep ($p\leq 0.0007$), daytime alertness ($p=0.03$), clarity of thought and concentration ($p=0.02$), and ability to function ($p=0.007$). Patients in the eszopiclone group demonstrated significantly greater improvement in symptoms of depression, as measured by HAM-D17 (Hamilton Depression Rating Scale), at weeks 4 ($p=0.01$) and 8 ($p=0.002$). HAM-D17 response were noted in 59% of patients in the eszopiclone group compared to 48% of patients in the placebo group; remission rates were 42% and 33%, respectively ($p=0.03$). Study completion rates and treatment tolerability were similar between groups.

A multicenter, randomized, double-blind, placebo-controlled trial evaluating eszopiclone treatment upon patient-reported sleep, fatigue and sleepiness, insomnia severity, quality of life, and work limitations for 6 months.¹⁶¹ A total of 830 patients with primary insomnia, who reported mean nightly total sleep time ≤ 6.5 hours/night and/or mean nightly sleep latency > 30 minutes, were randomized to eszopiclone 3 mg or matching placebo for 6 months. Patient-reported sleep and daytime function, Insomnia Severity Index, Physical Functioning, Vitality, and Social Functioning, and Work Limitations Questionnaire domain scores were improved with eszopiclone versus placebo (all $p<0.05$).

ramelteon (Rozerem) versus placebo

In a double-blind study, investigators randomized 829 elderly patients (mean age 72.4 years) with chronic primary insomnia to either ramelteon 4 mg, ramelteon 8 mg, or placebo nightly for 5 weeks.¹⁶² Administration of ramelteon resulted in a reduction in subjective SOL and an increase in TST at weeks 1, 3, and 5 of the study. Ramelteon did not change subjective sleep quality, the number of night time awakenings, or the ease of falling back to sleep. Withdrawal effects, including rebound insomnia, were not observed.

In a double-blind study, 405 patients (mean age 39.3 years) with primary insomnia were randomized to receive ramelteon 8 mg, ramelteon 16 mg, or placebo nightly for 35 nights.¹⁶³ Polysomnography indicated that both doses of ramelteon were associated with a reduction in SOL at each assessment starting on nights 1 and 2. Ramelteon was also associated with an improvement in TST and sleep efficiency on nights 1 and 2.

A 6-month, randomized, double-blind, placebo-controlled, multicenter study evaluated the long-term efficacy of ramelteon for insomnia in 451 adults (age ≥ 18 years) with chronic primary insomnia.¹⁶⁴ Patients were randomized to receive either ramelteon 8 mg or placebo 30 minutes before bedtime nightly for 6 months. Sleep was evaluated by polysomnography and morning questionnaires on the first 2 nights of week 1; the last 2 nights of months 1, 3, 5, and 6; and nights 1 and 2 of the placebo run-out. Next-morning residual effects, as well as adverse effects and vital signs, were recorded at each visit. Rebound insomnia and withdrawal effects were evaluated during placebo run-out. During the 6 months of treatment, ramelteon consistently reduced latency to persistent sleep compared with baseline and with placebo; significant decreases were observed at week 1 and months 1, 3, 5, and 6 ($p < 0.05$). Ramelteon significantly reduced subjective sleep latency relative to placebo at week 1, month 1, and month 5 ($p < 0.05$), with reductions nearing statistical significance at months 3 and 6 ($p \leq 0.08$). No significant next-morning residual effects were detected during ramelteon treatment. No withdrawal symptoms or rebound insomnia were detected after ramelteon discontinuation. Most adverse events were mild or moderate in severity.

suvorexant (Belsomra) versus placebo

In 3 clinical trials, suvorexant was evaluated in patients with insomnia characterized by sleep onset and sleep maintenance difficulties.¹⁶⁵ Non-elderly patients (age 18 to 64) and elderly patients (age ≥ 65) were randomized separately in 2 similarly designed, 3-month, randomized, double-blind, placebo-controlled, parallel-group studies. For the studies together, there were 465 female and 275 male non-elderly adults with a mean age of 46 years treated with 20 mg suvorexant ($n = 291$) or placebo ($n = 449$). Elderly patients with a mean age of 71 years including 346 female and 174 male patients were treated with 15 mg suvorexant ($n = 202$) or placebo ($n = 318$). In both studies, both suvorexant 15 mg or 20 mg was superior to placebo for sleep latency assessed objectively by polysomnography, as well as subjectively by patient-estimated sleep latency. Efficacy was similar for males and females and for Caucasians and non-Caucasians, based on the limited data. These doses were also superior to placebo for sleep maintenance as assessed objectively by polysomnography, as well as subjectively by patient-estimated total sleep time. In these 2 studies, suvorexant was also evaluated at doses of 30 mg and 40 mg. The higher doses had similar efficacy to the lower doses; however, there were more side effects reported at the higher doses.

The third study was a 1-month crossover study, that involved non-elderly adults, ages 18 to 64 years, with a mean age of 44 years, treated with placebo (n = 249) and suvorexant at a dose of 10 mg (n = 62), 20 mg (n = 61) or up to 80 mg. Both 10 mg and 20 mg were superior to placebo for sleep latency and sleep maintenance, both assessed objectively by polysomnography.

There were also special safety studies for suvorexant to evaluate effects on driving, next-day memory and balance in elderly and non-elderly patients, middle of the night safety in elderly patients, rebound effects, withdrawal effects, and respiratory safety.

tasimelteon (Hetlioz) versus placebo

The effectiveness of tasimelteon was evaluated in 104 patients in 2 randomized, double-masked, placebo-controlled, clinical trials of totally blind individuals with non-24-hour sleep-wake disorder (non-24) disorder.^{166,167} Patients were randomized to receive tasimelteon or placebo 1 hour prior to bedtime, at the same time every night. In Study 1 (Safety and Efficacy of Tasimelteon [SET]), 84 patients with non-24 were randomized to receive tasimelteon 20 mg or placebo, 1 hour prior to bedtime, at the same time every night for up to 6 months. Study 2 (Randomized Withdrawal Study of the Efficacy and Safety of Tasimelteon [RESET]) was a randomized withdrawal trial in that evaluated tasimelteon (n = 10) compared to placebo (n = 10), in patients with non-24 to evaluate the maintenance of efficacy. Patients were treated initially for approximately 12 weeks. Patients in whom the calculated time of peak melatonin level occurred at approximately the same time of day (in contrast to the expected daily delay) during the run-in phase were randomized to receive placebo or continue treatment for 8 weeks. Study 1 and Study 2 evaluated the duration and timing of night time sleep and daytime naps via patient-recorded diaries. Because symptoms of night time sleep disruption and daytime sleepiness are cyclical in patients with non-24, with severity varying, efficacy endpoints for night time total sleep time and daytime nap duration were based on the 25% of nights with the least night time sleep, and the 25% of days with the most daytime nap time. In Study 1, patients had, at baseline, an average 195 minutes of night time sleep and 137 minutes of daytime nap time on the 25% of most symptomatic nights and days, respectively. Treatment with tasimelteon resulted in a significant improvement, compared with placebo. In Study 1, mean total night time sleep was 28 minutes longer and daytime nap time was 27 minutes shorter in the tasimelteon group compared to placebo. In Study 2, synchronization was maintained in 90% of the tasimelteon group versus 20% of placebo. Mean total night time sleep was 67 minutes longer and daytime nap time was 59 minutes shorter in the tasimelteon group compared to placebo. A responder analysis of patients with both ≥ 45 minutes increase in night time sleep and ≥ 45 minutes decrease in daytime nap time was conducted in Study 1: 29% (n = 12) of patients treated with tasimelteon, compared with 12% (n = 5) of patients treated with placebo met the responder criteria.

zolpidem (Intermezzo) versus placebo

A randomized, double-blind, placebo-controlled, 3-way crossover study evaluated the efficacy and safety of low-dose, sublingual zolpidem tartrate when taken during a scheduled middle-of-the-night (MOTN) awakening in subjects with insomnia characterized by difficulty returning to sleep following MOTN awakenings.¹⁶⁸ The study was performed at 5 sleep laboratories and enrolled adults (24 males, 58 females, mean age 45.9 years) with a diagnosis of DSM-IV primary insomnia and a history of prolonged MOTN awakenings. Baseline difficulties with MOTN awakenings were confirmed by a 10-day screening sleep diary and polysomnography (PSG) screening. Each treatment period consisted of 2 consecutive nights of dosing separated by a washout period of 5 to 12 days. Subjects were awakened 4

hours after lights out, dosed with sublingual zolpidem 3.5 mg, zolpidem 1.75 mg, or placebo, kept awake for 30 minutes, and then returned to bed for an additional 4 hours. Sleep parameters were assessed by PSG and post-sleep questionnaires. Results demonstrated that low-dose sublingual zolpidem tartrate demonstrated significant dose-related decreases in latency to persistent sleep and total sleep time ($p < 0.001$) compared to placebo after MOTN dosing. All subject reports paralleled PSG observations. Neither dose showed next-morning impairment on the digit symbol substitution test (DSST) or ratings of sleepiness. The 3.5-mg dose produced improvements in reports of sleep quality ($p < 0.001$), ability to function, and level of refreshed sleep ($p < 0.05$ for both dosages) compared to placebo. Sublingual zolpidem tartrate lozenges were generally safe and well tolerated.

In a randomized, double-blind, placebo-controlled outpatient as-needed study, adults aged 18 to 64 years ($n = 295$; 201 female; 94 male) with a history of problems returning to sleep after MOTN awakenings were evaluated in a 4-week study using zolpidem tartrate.^{169,170} Patients used either 3.5 mg zolpidem tartrate or placebo (as needed) following episodes of MOTN waking with difficulties in returning to sleep. Patients were required to have at least 4 hours of time to remain in bed following zolpidem dosing. The results were patient evaluations of the time to return to sleep following as needed dosing were significantly shorter for zolpidem 3.5 mg than placebo.

zolpidem (Zolpimist) versus placebo

In a double-blind, parallel-group, single-night trial in adults experiencing transient insomnia ($n = 462$), zolpidem 7.5 mg or 10 mg or placebo were compared.¹⁷¹ Both zolpidem doses were superior to placebo by measure of sleep latency, sleep duration, and number of awakenings.

zolpidem ER (Ambien CR) versus placebo

Two similar 3-week studies of zolpidem ER were conducted in patients with primary insomnia.^{172,173} One study randomized 205 elderly patients (mean age 70.2 years) to zolpidem ER 6.25 mg or placebo while the other randomized 212 adults (mean age 44.3 years) to zolpidem ER 12.5 mg or placebo. In each study, zolpidem ER was found to lead to significant improvement compared to placebo in polysomnographic WASO in the first 6 hours of the night, as well as improvement in SOL and TST. Subjects did not report any residual impairment or sedation.

META-ANALYSES

A meta-analysis of clinical trials submitted to the FDA assessed the efficacy of eszopiclone, zaleplon, and zolpidem compared to placebo for the treatment of insomnia in adults (13 trials; $n = 4,378$).¹⁷⁴ Overall, these agents had significant but minimal improvements in the following primary outcomes: effect size for polysomnographic sleep latency (weighted mean difference, -0.36; 95% confidence interval [CI], -0.57 to -0.16), effect size for subjective sleep latency (weighted mean difference, -0.33; 95% CI, -0.62 to -0.04), and weighted mean raw differences in sleep latency (-22 minutes; 95% CI, -33 to -11). No significant differences were found in other outcomes.

Other meta-analyses also have found minimal differences with regard to efficacy of benzodiazepines and the non-benzodiazepine hypnotics, but are limited by treatment duration (e.g., zaleplon, zolpidem, eszopiclone).^{175,176,177}

SUMMARY

The selection of a specific hypnotic is based in large part on whether the patient has problems with initiation or maintenance of sleep, co-morbid conditions, side effect tolerance, and availability. Sedative hypnotics should be prescribed at the lowest dose that treats the patients' symptoms.

The assumed increased risk with benzodiazepine medications over non-benzodiazepines is based on indirect comparisons, and there is evidence of publication bias as both groups have increased incidence of adverse risks in patients over 60 years of age.

Benzodiazepines are Scheduled IV controlled substances. In general, the benzodiazepines decrease the time for sleep onset and prolong the duration of sleep, although dependence, tolerance, and abuse may occur. Among the benzodiazepines, the duration of action is the primary variable that may make 1 preferable to another in a given patient. Triazolam (Halcion) has the shortest duration of action, while temazepam (Restoril) and estazolam have intermediate durations. Flurazepam and quazepam (Doral) have long durations of effect and, as a result, should be avoided in the elderly or others in whom daytime sedation may be a concern.

Eszopiclone (Lunesta), suvorexant (Belsomra), zaleplon (Sonata), zolpidem (Ambien, Ambien CR, Edluar, Intermezzo, Zolpimist) are also Scheduled IV controlled substances. Patients with a history of addiction to, or abuse of, drugs or alcohol are at increased risk for misuse, abuse and addiction these agents; patients should be monitored carefully when taking these agents. Doxepin (Silenor), ramelteon (Rozerem), and tasimelteon (Hetlioz) are not controlled substances and are not associated with abuse or physical dependence.

Rebound insomnia may develop when benzodiazepines are abruptly withdrawn and is more likely to occur with the short-acting benzodiazepines. Rebound insomnia can be minimized by using smaller doses and tapering the dosage. Some studies have highlighted concerns with increased falls and hip fractures in the elderly following benzodiazepine use; however, others have found that untreated insomnia itself increases the risk of falls.

Similar to the benzodiazepines, the BZ-1 selective agents decrease sleep latency with duration of action again being the primary difference among these agents. Although dependence, tolerance, and abuse may occur with these agents, next day sedation, rebound insomnia, and drug interactions are generally lessened. For the BZ-1 selective agents, zaleplon (Sonata) is more rapid acting with a shorter duration than zolpidem tablets (Ambien, Ambien CR). Eszopiclone (Lunesta) has a longer half-life than either zaleplon or zolpidem. Specialized formulations of zolpidem (Edluar, Intermezzo, Zolpimist) do not have a significant clinical advantage over tablets. Additionally, due to gender differences in zolpidem clearance, women require lower doses of zolpidem.

Doxepin (Silenor) appears to have some success in treating insomnia, particularly in cases where sleep latency and not sleep initiation is the issue. Patients experiencing problems with sleep latency who have not achieved success with more common sedative/hypnotic agents may benefit from doxepin. However, in general, no data are present to suggest doxepin is superior to other agents in this class.

The melatonin receptor agonist, ramelteon (Rozerem), has demonstrated reduction in sleep latency, but not in sleep maintenance. Patient evaluations are inconsistent, and there are no direct comparative studies. There is a low likelihood of dependence or abuse, and adverse effects are rare.

Tasimelteon (Hetlioz), a melatonin receptor agonist pharmacologically similar to ramelteon (Rozerem), is approved for non-24 in totally blind patients. Due to differences in circadian rhythms, it can take weeks or months of daily use of tasimelteon before the patient experiences any benefit. Comparisons to ramelteon or melatonin are lacking.

Suvorexant (Belsomra), an orexin receptor antagonist, has shown better sleep latency and sleep maintenance compared to placebo. There is no evidence that suvorexant offers significant advantages over existing drugs.

Current treatment guidelines for insomnia do not recommend one agent within this class over another; rather, they suggest that treatment should be individualized.

REFERENCES

- 1 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 2 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 3 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 4 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 5 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 6 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 7 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 8 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 9 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 10 Halcion [package insert]. New York, NY; December 2016.
- 11 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 12 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 13 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 14 Intermezzo [package insert]. Stamford, CT; Purdue; September 2015.
- 15 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 16 Ambien CR [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 17 Diagnostic Classification Commission. International Classification of Sleep Disorders: Diagnostic and Coding Manual (ICSD). Rochester, MN: American Sleep Disorders Association; 1990.
- 18 Morgenthaler T, Kramer M, Alessi, et al. Practice Parameters for the Psychological and Behavioral treatment of Insomnia: An Update. An American Academy of Sleep Medicine Report. *Sleep*. 2006 (11) 1415-1419.
- 19 Sateia MJ, Buysse D, Krystal AD, et al. Clinical Practice Guideline for the Pharmacologic Treatment of Chronic Insomnia in Adults: An American Academy of Sleep Medicine Clinical Practice Guideline <http://www.aasmnet.org/practiceparameters.aspx?cid=109> *J Clin Sleep Med*. 2017; 13(2):307-342. Available at: Accessed December 17, 2018.
- 20 Management of Chronic Insomnia Disorder in Adults: A Clinical Practice Guideline From the American College of Physicians 2016. Available at: <https://www.acponline.org/clinical-information/guidelines>. Accessed December 17, 2018.
- 21 Sadeh A, Raviv A, Gruber R. Sleep patterns and sleep disruptions in school-age children. *Dev Psychol*. 2000; 36:291-301.
- 22 Lindblom N, Heiskala H, Kaski M, et al. Neurological impairments and sleep-wake behaviour among the mentally retarded. *J Sleep Res*. 2001; 10:309-18.
- 23 Quine L. Sleep problems in primary school children: comparison between mainstream and special school children. *Child Care Health Dev*. 2001; 27:201-21.
- 24 Mindell JA, Emslie G, Blumer J, et al. Pharmacologic Management of Insomnia in Children and Adolescents: Consensus Statement. *Pediatrics*. 2006; 117:1223-32.
- 25 Owens JA, Babcock D, Blumer J, et al. The Use of Pharmacotherapy in the Treatment of Pediatric Insomnia in Primary Care: Rational Approaches. A Consensus Meeting Summary. *J Clin Sleep Med*. 2005; 1:49-59.
- 26 Mindell JA, Emslie G, Blumer J, et al. Pharmacologic Management of Insomnia in Children and Adolescents: Consensus Statement. *Pediatrics*. 2006; 117:1223-32. Available at: <http://pediatrics.aappublications.org/content/117/6/e1223>. Accessed December 17, 2018.
- 27 Non-24 Hour Sleep-wake Disorder. Available at: <http://www.rarediseases.org/rare-disease-information/rare-diseases/byID/1275/viewAbstract>. Accessed December 17, 2018.
- 28 Auger RR, Burgess HJ, Emens JS, et al. Clinical Practice Guideline for the Treatment of Intrinsic Circadian Rhythm Sleep-Wake Disorders: Advanced Sleep-Wake Phase Disorder (ASWPD), Delayed Sleep-Wake Phase Disorder (DSWPD), Non-24-Hour Sleep-Wake Rhythm Disorder (N24SWD), and Irregular Sleep-Wake Rhythm Disorder (ISWRD). An Update for 2015. Available at: <https://aasm.org/clinical-resources/practice-standards/practice-guidelines/>. Accessed December 17, 2018.
- 29 The Medical Letter Treatment Guidelines. Drugs for Insomnia. March 2009; 7: 79.
- 30 Drover DR. Comparative pharmacokinetics and pharmacodynamics of short-acting hypnotics: zaleplon, zolpidem and zopiclone. *Clin Pharmacokinet*. 2004; 43:227-38.
- 31 Kato K, Hirai K, Nishiyama K, et al. Neurochemical properties of ramelteon (TAK-375), a selective MT1/MT2 receptor agonist. *Neuropharmacology*. 2005; 48:301-10.
- 32 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 33 Ramakrishnan K, Scheid D. Treatment options for Insomnia. *American Family Physician*. 2007; 76:517-26, 527-8.

-
- 34 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 35 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 36 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 37 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 38 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 39 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 40 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 41 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 42 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 43 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 44 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 45 Halcion [package insert]. New York, NY; December 2016.
- 46 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 47 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 48 Ambien CR [package insert]. New York, NY; Sanofi-Aventis; September 2018.
- 49 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 50 Intermezzo [package insert]. Stamford, CT; Purdue; September 2015
- 51 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 52 Gustavson LE, Carrigan PJ. The clinical pharmacokinetics of single doses of estazolam. *Am J Med.* 1990; 88(3A):2S-5S.
- 53 Gary M, Rubens R, Amato D. Pharmacokinetic (PK) and pharmacodynamic (PD) effects of eszopiclone: a comparison of healthy non-elderly and elderly adults [abstract]. *Sleep.* 2004; 27(suppl):A56.
- 54 Dopheide JA, Stimmel GL. Sleep disorder. In: Koda-Kimble MA, Young LY, eds. *Applied Therapeutics: The Clinical Use of Drugs*, 8th ed. Philadelphia: Lippincott Williams and Wilkins, 2004:77.1-77.2.
- 55 *The Medical Letter.* Low-dose sublingual zolpidem (intermezzo) for insomnia due to middle-of-the-night awakening. April 2, 2012;54:25-26.
- 56 Greenblatt DJ, Legangneux E, Harmatz JHS, et al. Dynamics and kinetics of a modified-release formulation of zolpidem: comparison with immediate-release standard zolpidem and placebo. *J Clin Pharmacol.* 2006; 46:1469-80.
- 57 Available at: <http://www.clinicalpharmacology.com>. Accessed December 17, 2018.
- 58 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 59 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 60 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 61 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 62 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 63 Halcion [package insert]. New York, NY; December 2016.
- 64 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 65 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 66 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 67 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 68 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 69 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 70 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 71 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 72 FDA. FDA Drug Safety Communication: Risk of next-morning impairment after use of insomnia drugs; FDA requires lower recommended doses for certain drugs containing zolpidem (Ambien, Ambien CR, Edluar, and Zolpimist). <https://www.fda.gov/downloads/Drugs/DrugSafety/UCM335007.pdf>. Accessed December 17, 2018.
- 73 FDA Safety Communication. FDA warns about serious risks and death when combining opioid pain or cough medicines with benzodiazepines; requires its strongest warning. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm518473.htm>. Accessed December 17, 2018.
- 74 FDA Safety Communication. FDA urges caution about withholding opioid addiction medications from patients taking benzodiazepines or CNS depressants: careful medication management can reduce risks Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm575307.htm>. Accessed December 17, 2018.
- 75 Donnelly K, Bracchi R, Hewitt J, et al. Benzodiazepines, Z-drugs, and the risk of hip fracture: A systematic review and a meta-analysis. *PLOS one.* 2017;12(4).
- 76 Available at: <http://www.clinicalpharmacology.com>. Accessed December 17, 2018.
- 77 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 78 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 79 Flurazepam [package insert]. Morgantown, WV; Mylan Pharmaceuticals; September 2016.
- 80 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 81 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 82 Halcion [package insert]. New York, NY; December 2016.
- 83 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 84 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 85 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 86 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 87 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 88 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 89 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 90 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 91 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
-

-
- 92 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 93 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 94 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 95 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 96 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 97 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 98 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 99 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 100 Halcion [package insert]. New York, NY; December 2016.
- 101 Sonata [package insert]. New York, NY; Pfizer; August 2014.
- 102 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 103 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 104 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 105 Intermezzo [package insert]. Stamford, CT; Purdue; September 2015
- 106 Ambien CR [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 107 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 108 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 109 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 110 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 111 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 112 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 113 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 114 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 115 Halcion [package insert]. New York, NY; December 2016.
- 116 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 117 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 118 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 119 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 120 Owens JA, Rosen CL, Mindell JA. Medication Use in the Treatment of Pediatric Insomnia: Results of a Survey of Community-Based Pediatricians. *Pediatrics*. 2003; 111:e628-35.
- 121 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 122 Pariente A, et al. Benzodiazepines and Injurious Falls in Community Dwelling Elders. *Drugs & Aging*. 2008; 25:61.
- 123 Avidan AY, et al. Insomnia and Hypnotic Use, Recorded in the Minimum Data Set as Predictors of Falls and Hip Fractures in Michigan Nursing Homes. *Journal of American Geriatrics Society*. 2005; 53: 955.
- 124 The Medical Letter Treatment Guidelines. *Drugs for Insomnia*. March 2009; 7: 79.
- 125 Available at: <http://www.clinicalpharmacology.com>. Accessed December 17, 2018.
- 126 Estazolam [package insert]. Greenville, NC; Mayne; October 2016.
- 127 Lunesta [package insert]. Marlborough, MA; Sunovion; November 2018.
- 128 Flurazepam [package insert]. Morgantown, WV; Mylan; September 2016.
- 129 Doral [package insert]. Marietta, GA; Galt; August 2017.
- 130 Restoril [package insert]. Dublin, Ireland; Mallinckrodt; September 2017.
- 131 Halcion [package insert]. New York, NY; December 2016.
- 132 Sonata [package insert]. New York, NY; Pfizer; November 2016.
- 133 Ambien [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 134 Rozerem [package insert]. Deerfield, IL; Takeda; December 2018.
- 135 Ambien CR [package insert]. Bridgewater, NJ; Sanofi-Aventis; September 2018.
- 136 Edluar [package insert]. Somerset, NJ; Meda; October 2018.
- 137 Silenor [package insert]. Morristown, NJ. Pernix; March 2010.
- 138 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 139 Intermezzo [package insert]. Stamford, CT; Purdue; September 2015
- 140 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 141 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 142 Cohn JB, Wilcox CS, Bremner J, et al. Hypnotic efficacy of estazolam compared with flurazepam in outpatients with insomnia. *J Clin Pharmacol*. 1991; 31:747-50.
- 143 Scharf MB, Roth PB, Dominguez RA, et al. Estazolam and flurazepam: a multicenter, placebo-controlled comparative study in outpatients with insomnia. *J Clin Pharmacol*. 1990; 30:461-7.
- 144 Dement WC. Objective measurements of daytime sleepiness and performance comparing quazepam with flurazepam in two adult populations using the Multiple Sleep Latency Test. *J Clin Psychiatry*. 1991; 52 Suppl:31-7.
- 145 Saletu B, Anderer P, Brandstatter N, et al. Insomnia in generalized anxiety disorder: polysomnographic, psychometric and clinical investigations before, during and after therapy with a long- versus a short-half-life benzodiazepine (quazepam versus triazolam). *Neuropsychobiology*. 1994; 29:69-90.
- 146 Scharf MB. Feasibility of an every-other-night regimen in insomniac patients: subjective hypnotic effectiveness of quazepam, triazolam, and placebo. *J Clin Psychiatry*. 1993; 54:33-8.
- 147 Voshaar RC, van Balkom AJ, Zitman FG. Zolpidem is not superior to temazepam with respect to rebound insomnia: a controlled study. *Eur Neuropsychopharmacol*. 2004; 14:301-6.
- 148 Walsh JK, Fry J, Erwin CW, et al. Efficacy and tolerability of 14-day administration of zaleplon 5 mg and 10 mg for the treatment of primary insomnia. *Clin Drug Invest*. 1998; 16:347-54.
-

- 149 Monti JM, Attali P, Monti D, et al. Zolpidem and rebound insomnia—a double-blind, controlled polysomnographic study in chronic insomniac. *Pharmacopsychiatry*. 1994; 27:166-75.
- 150 Elie R, Ruther E, Farr I, et al. Sleep latency is shortened during 4 weeks of treatment with zaleplon, a novel nonbenzodiazepine hypnotic. *J Clin Psychiatry*. 1999; 60:536-44.
- 151 Zammit GK, Corser B, Doghramji K, et al. Sleep and residual sedation after administration of zaleplon, zolpidem, and placebo during experimental middle-of-the-night awakening. *J Clin Sleep Med*. 2006; 2(4):417-23.
- 152 Roth T, Heith Durrence H, Jochelson P, et al. Efficacy and safety of doxepin 6 mg in a model of transient insomnia. *Sleep Med*. 2010;11(9):843-7.
- 153 Krystal A, Lankford A, Durrence H, et al. Efficacy and safety of doxepine 3 and 6 mg in a 35—day sleep laboratory trial in adults with chronic primary insomnia. *Sleep*. 2011; 34(10):1433-1442.
- 154 Zammit GK, McNabb LJ, Caron J, et al. Efficacy and safety of eszopiclone across 6-weeks of treatment for primary insomnia. *Curr Med Res Opin*. 2004; 20:1979-91.
- 155 Rubens R, Wessel T, Zammit G. A six-week efficacy and safety study of eszopiclone in adults with chronic insomnia [abstract]. *Neurology*. 2004; 62(suppl 5):A59-60.
- 156 Wessel T, Rubens R, McCall V. A study of eszopiclone 2 mg in elderly patients with chronic insomnia [abstract]. *Neurology*. 2004; 62(suppl 5):A60.
- 157 Scharf M, Seiden D, Erman M, et al. Eszopiclone rapidly induced sleep and provided sleep maintenance in elderly patients with chronic insomnia [abstract]. *Int Psychogeriatrics*. 2003; 15(suppl 2):200-201.
- 158 Scharf M, McCall W, Erman M, et al. Patient-reported efficacy of eszopiclone (ESZ) in elderly patients with chronic insomnia [abstract]. *J Am Geriatrics Soc*. 2004; 52(suppl):S14.
- 159 Erman M, Rosenberg R, Caron J. Polysomnographic and patient-reported evaluation of the efficacy and safety of eszopiclone in elderly subjects with chronic insomnia [abstract]. *Sleep*. 2004; 27(suppl):A257.
- 160 Fava M, McCall MW, Krystal A, et al. Eszopiclone Co-Administered With Fluoxetine in Patients With Insomnia Coexisting With Major Depressive Disorder. *Biological Psychiatry*. 2006; 59(11):1052-60.
- 161 Walsh JK, Krystal AD, Amato DA, et al. Nightly treatment of primary insomnia with eszopiclone for six months: effect on sleep, quality of life, and work limitations. *Sleep*. 2007; 30(8):959-68.
- 162 Seiden D, Zee P, Weigand S, et al. Double-blind, placebo-controlled outpatient clinical trial of ramelteon for the treatment of chronic insomnia in an elderly population. Program and abstracts of the Associated Professional Sleep Societies 19th Annual Meeting; June 18-23, 2005; Denver, Colorado. Abstract 0679.
- 163 Zammit G, Roth T, Erman M, et al. Double-blind, placebo-controlled polysomnography and outpatient trial to evaluate the efficacy and safety of ramelteon in adult patients with chronic insomnia. Program and abstracts of the Associated Professional Sleep Societies 19th Annual Meeting; June 18-23, 2005; Denver, Colorado. Abstract 0680.
- 164 Mayer G, Wang-Weigand S, Roth-Schechter B, et al. Efficacy and safety of 6-month nightly ramelteon administration in adults with chronic primary insomnia. *Sleep*. 2009; 32(3):351-360.
- 165 Belsomra [package insert]. Whitehouse Station, NJ; Merck; July 2018.
- 166 Hetlioz [package insert]. Washington, D.C.; Vanda; December 2014.
- 167 Lockley SW, Dressman MA, Licamele L, et al. Tasimelteon for non-24-hour sleep-wake disorder in totally blind people (SET and RESET): two multicentre, randomised, double-masked, placebo-controlled phase 3 trials. *Lancet*. 2015; 386(10005): 1754-1764. DOI: 10.1016/S0140-6736(15)60031-9.
- 168 Roth T, Hull SG, Lankford DA, et al. Low-dose sublingual zolpidem tartrate is associated with dose-related improvement in sleep onset and duration in insomnia characterized by middle-of-the-night (MOTN) awakenings. *Sleep*. 2008; 31(9):1277-1284.
- 169 Intermezzo [package insert]. Stamford, CT; Purdue; September 2015.
- 170 Roth T, Krystal A, Steinberg FJ, et al. Novel sublingual low-dose zolpidem tablet reduces latency to sleep onset following spontaneous middle-of-the-night awakening in insomnia in a randomized, double-blind, placebo-controlled, outpatient study. *Sleep*. 2013;36(2):189-96. DOI: 10.5665/sleep.2370.
- 171 Zolpimist [package insert]. Louisville, KY; Magna; January 2019.
- 172 Rohers TA, Soubrane C, Walsh J, et al. Efficacy and safety of 6.25 mg of zolpidem modified-release formulation in elderly patients with primary insomnia. Program and abstracts of the American Psychiatric Association 2005 Annual Meeting; May 21-26, 2005; Atlanta, GA. Abstract NR586.
- 173 Soubrane C, Walsh JK, Roth T. Efficacy and safety of a modified release formulation of zolpidem in adults. Program and abstracts of the American Psychiatric Association 2005 Annual Meeting; May 21-26, 2005; Atlanta, GA. Abstract NR582.
- 174 Huedo-Medina TB, Kirsch I, Middlemass J, et al. Effectiveness of non-benzodiazepine hypnotics in treatment of adult insomnia: meta-analysis of data submitted to the Food and Drug Administration. *BMJ*. 2012; 345: e8343. DOI: 10.1136/bmj.e8343.
- 175 Dunder Y, Dodd S, Strobl J, et al. Comparative efficacy of newer hypnotic drugs for the short-term management of insomnia: a systematic review and meta-analysis. *Hum Psychopharmacol*. 2004; 19(5): 305-322.
- 176 Smith MT, Perlis ML, Park A, et al. Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *Am J Psychiatry*. 2002; 159(1): 5-11.
- 177 Nowell PD, Mazumdar S, Buysse DJ, et al. Benzodiazepines and zolpidem for chronic insomnia: a meta-analysis of treatment efficacy. *JAMA*. 1997; 278(24): 2170-2177.