NUTRITION RISK CRITERIA

CODE	NUTRITION RISK CRITERIA
101★	UNDERWEIGHT WOMAN
103★	UNDERWEIGHT / AT RISK FOR UNDERWEIGHT—INFANTS/CHILDREN
111★	OVERWEIGHT WOMAN
113★	OBESE—CHILDREN (2-5)
114★	OVERWEIGHT---CHILDREN (2-5)
115★	HIGH WEIGHT-FOR-LENGTH---INFANTS/CHILDREN (< 24 MONTHS)
121★	SHORT STATURE/AT RISK OF SHORT STATURE—INFANTS/CHILDREN
131★	LOW MATERNAL WEIGHT GAIN
132★	MATERNAL WEIGHT LOSS IN PREGNANCY
133★	HIGH MATERNAL WEIGHT GAIN
134	FAILURE TO THRIVE
135★	SLOWED/FALTERING GROWTH PATTERN
141★	LOW BIRTH WEIGHT
142★	PRETERM OR EARLY TERM DELIVERY
153★	LARGE FOR GESTATIONAL AGE
201★	LOW HEMATOCRIT/LOW HEMOGLOBIN
301	HYPEREMESIS GRAVIDARUM
302	GESTATIONAL DIABETES
303	HX GESTATIONAL DIABETES
304	HISTORY OF PREECLAMPSIA
311	HX PRETERM OR EARLY TERM DELIVERY
312	HX LOW BIRTH WEIGHT
321	HX FETAL OR NEONATAL LOSS
331★	PREGNANCY—YOUNG AGE
332★	SHORT INTERPREGNANCY INTERVAL
334	LACK OF OR INADEQUATE PRENATAL CARE

CODE	NUTRITION RISK CRITERIA
335★	MULTIFETAL GESTATION
336	FETAL GROWTH RESTRICTION
337	HX BIRTH LGA INFANT
338	PREGNANT AND BREASTFEEDING
339	HX BIRTH—CONGENITAL DEFECT
341	NUTRIENT DEFICIENCY DISEASES
342	GASTRO-INTESTINAL DISORDERS
343	DIABETES MELLITUS
344	THYROID DISORDERS
345	HYPERTENSION AND PREHYPERTENSION
346	RENAL DISEASE
347	CANCER
348	CENTRAL NERVOUS SYSTEM DISORDERS
349	GENETIC AND CONGENITAL DISORDERS
351	INBORN ERRORS OF METABOLISM
352.1	INFECTIOUS DISEASES - ACTUE
352.2	INFECTIOUS DISEASES - CHRONIC
353	FOOD ALLERGY
354	CELIAC DISEASE
355	LACTOSE INTOLERANCE
356	HYPOGLYCEMIA
357	DRUG NUTRIENT INTERACTIONS
358	EATING DISORDERS
359	RECENT MAJOR SURGERY, TRAUMA, BURNS
360	OTHER MEDICAL CONDITIONS
361	DEPRESSION
362	DEVELOPMENTAL, SENSORY, OR MOTOR DELAYS INTERFERING WITH ABILITY TO EAT

CODE	NUTRITION RISK CRITERIA
363	PRE-DIABETES
371	MATERNAL SMOKING
372	ALCOHOL OR ILLEGAL DRUG USE
381	ORAL HEALTH CONDITIONS
382	FETAL ALCOHOL SYNDROME
383	NEONATAL ABSTINENCE SYNDROME
401	FAILURE TO MEET DIETARY GUIDELINES FOR AMERICANCS
411	INAPPROPRIATE NUTRITION PRACTICES—INFANT
425	INAPPROPRIATE NUTRITION PRACTICES—CHILD
427	INAPPROPRIATE NUTRITION PRACTICES—WOMAN
428	DIETARY RISK ASSOCIATED WITH COMPLEMENTARY FEEDING PRACTICES
501	POSSIBILITY OF REGRESSION
502★	TRANSFER OF CERTIFICATION (VOC)
601	BF MOM OF INFANT AT NUTRITIONAL RISK
602	BF COMPLICATIONS OR POTENTIAL COMPLICATIONS—WOMAN
603	BF COMPLICATION—INFANT
701	BORN TO WIC MOM / BORN TO POTENTIAL WIC MOM
702	BF INFANT OF MOM AT NUTRITIONAL RISK
801★	HOMELESSNESS
802★	MIGRANCY
902	FEEDING SKILLS LIMITATION
903★	FOSTER CARE
904	EXPOSURE TO ENVIRONMENTAL TOBACCO SMOKE (ETS)

Nutrition Risk Criteria **101★ UNDERWEIGHT WOMAN**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	-

Definition

Pregnant: Body Mass Index (BMI) less than (<) 18.5.

- Use pre-pregnancy weight in BMI calculation.

Breastfeeding/Non-breastfeeding: BMI less than (<) 18.5.

- Use pre-pregnancy weight in BMI calculation ***or*** current weight in BMI calculation for all women who are less than 6 months postpartum.
- Use current weight for breastfeeding women who are equal or greater than 6 months postpartum.

Note: A BMI table is attached to assist in determining weight classifications. Until research supports the use of different BMI cut-offs for adolescent pregnancies, the same BMI cut-offs will be used for all women, regardless of age, when determining WIC eligibility (1).

Required Documentation

Pregnant

- Pre-pregnancy weight
- Height
- BMI

Breastfeeding/Non-breastfeeding

- Pre-pregnancy weight
- Current weight
- Height
- BMI

Justification

Underweight women who become pregnant are at a higher risk for delivery of low birth weight (LBW) infants, retarded fetal growth, and perinatal mortality. Pre-pregnancy underweight is also associated with a higher incidence of various pregnancy complications, such as antepartum hemorrhage, premature rupture of membranes, anemia, endometriosis, and cesarean delivery.

The goal in prenatal nutritional counseling provided by WIC is to achieve recommended weight gain by emphasizing food choices of high nutritional quality; and for the underweight woman, by encouraging increased consumption and/or the inclusion of some calorically dense foods.

The 2009 Institute of Medicine (IOM) report: *Weight Gain During Pregnancy: Reexamining the Guidelines* (1) updated the pregnancy weight categories to conform to the categories developed by the World Health Organization and adopted by the National Heart Lung and Blood Institute in 1998 (3). The reexamination of the guidelines consisted of a review of the determinants of a wide range of short-and long-term consequences of variation in weight gain during pregnancy for both the mother and her infant. The IOM prenatal weight

gain recommendations based on pre-pregnancy weight status categories are associated with improved maternal and child health outcomes (1).

Included in the 2009 IOM guidelines is the recommendation that the BMI weight categories used for adult women be used for pregnant adolescents as well. More research is needed to determine whether special categories are needed for adolescents. It is recognized that the IOM BMI cut-offs for defining weight categories will classify some adolescents differently than the CDC BMI-for-age charts. For the purpose of WIC eligibility determination, the IOM BMI cut-off will be used for all women regardless of age. However, due to the lack of research on relevant BMI cut-offs for pregnant and postpartum adolescents, professionals should use all of the tools available to them to assess these applicants' anthropometric status and tailor nutrition counseling accordingly.

Weight during the early postpartum period, when most WIC certifications occur, is very unstable. During the first 4-6 weeks fluid shifts and tissue changes cause fluctuations in weight. After 6 weeks, weight loss varies among women. Pre-pregnancy weight, amount of weight gain during pregnancy, race, age, parity and lactation all influence the rate of postpartum weight loss. By 6 months postpartum, body weight is more stable and should be close to the pre-pregnancy weight. In most cases therefore, pre-pregnancy weight is a better indicator of weight status than postpartum weight in the first 6 months after delivery. The one exception is the woman with a BMI of <18.5 during the immediate 6 months after delivery. Underweight at this stage may indicate inadequate weight gain during pregnancy, depression, an eating disorder or disease; any of which need to be addressed (4).

While being on the lean side of normal weight is generally considered healthy, being underweight can be indicative of poor nutritional status, inadequate food consumption, and/or an underlying medical condition. Underweight women who are breastfeeding may be further impacting their own nutritional status. Should she become pregnant again, an underweight woman is at a higher risk for delivery of low birth weight (LBW) infants, retarded fetal growth, and perinatal mortality. The role of the WIC Program is to assist underweight women in the achievement of a healthy dietary intake and body mass index.

References

1. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines (Prepublication Copy). National Academy Press, Washington, D.C.; 2009. www.nap.edu. Accessed June 2009.
2. Institute of Medicine. WIC Nutrition Risk Criteria: A Scientific Assessment. National Academy Press, Washington, D.C.; 1996.
3. National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health (NIH). Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. NIH Publication No. 98-4083,1998. www.nih.gov. Accessed June 2009.
4. Crowel DT. Weight changes in the postpartum period: a review of the literature. Journal of Nurse-Midwifery. Vol. 40, No. 5, September/October 1995; 418-423.

Additional Related References

1. Parker JD, Abrams B. Prenatal weight gain advice: an examination of the recent prenatal weight gain recommendations of the Institute of Medicine. Obstet Gynecology, 1992; 79:664-9.
2. Siega-Riz AM, Adair LS, Hobel CJ. Institute of Medicine maternal weight

gain recommendations and pregnancy outcomes in a predominately Hispanic population. Obstet Gynecol, 1994; 84:565-73.

3. Suitor CW, editor. Maternal weight gain: A report of an expert work group. Arlington, Virginia: National Center for Education in Maternal and Child Health; 1997. Sponsored by Maternal and Child Health Bureau, Health Resources and Services Administration, Public Health Service, U.S. Department of Health and Human Services.

Nutrition Risk Criteria **103★ UNDERWEIGHT OR AT RISK OF UNDERWEIGHT – INFANTS/CHILDREN**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD
CHILDREN	3	RD

Definition

UNDERWEIGHT:
BIRTH TO AGE 2: Weight-for-length less than or equal to (≤) 2.3rd percentile.*

CHILDREN AGE 2 TO 5 YEARS: Body Mass Index-for-age less than or equal to (≤) 5th percentile.**

AT RISK OF UNDERWEIGHT:
BIRTH TO AGE 2: greater than (>) 2.3rd percentile through less than or equal to (≤) 5th percentile weight-for-length.*

CHILDREN AGE 2 TO 5 YEARS: greater than (>) 5th percentile through less than or equal to (≤) 10th percentile Body Mass Index (BMI)-for-age.**

***Based on the Centers for Disease Control and Prevention (CDC) Birth to 24 months gender specific growth charts (1) based on 2006 World Health Organization (WHO) international growth standards (3). For the Birth to <24 months "underweight" definition, CDC labels the 2.3rd percentile as the 2nd percentile on the Birth to 24 months gender specific growth charts. For more information about the percentile cut-off, please see Clarification.

**Based on National Center for Health Statistics/Centers for Disease Control and Prevention age/sex specific growth charts (2000).

Required Documentation

DOB
Weight
Height
BMI
CDC/WHO growth charts, CDC growth charts

Justification

The CDC uses the 2.3rd percentile weight-for-length (for birth to 24 months of age) and the 5th percentile BMI-for-age (for 2-5 years of age), as the cut-offs to define underweight in its Pediatric Nutrition Surveillance System (1,2). However, CDC does not have a position regarding the cut-off percentile, which should be used to determine at risk of underweight as a nutrition risk in the WIC Program. At risk of underweight is included in this criterion to reflect the preventive emphasis of the WIC program.

A review of literature on weight-for-length or stature cut-off percentiles indicates that: a) many children at or below the 5th percentile for weight are in need of nutritional intervention, and b) those at or below the 10th percentile may be at nutritional risk and in need of preventive nutritional intervention, or at least further evaluation (4).

Weight-for-length/stature describes body proportionality and is sensitive to acute under nutrition but can also reflect long-term status (5). Physical growth delay is used as a proxy for the deleterious effects under nutrition can have on immune function, organ development, hormonal function and brain development (6).

Implications for WIC Nutrition Services

Participation in WIC has been associated with improved growth in both weight and height in children (7). An infant or child determined to be underweight at WIC certification should be monitored at regular intervals during the certification period, as appropriate. Through client-centered counseling, WIC staff can assist families in making nutritionally balanced food choices to promote adequate weight gain. Also, the foods provided by the WIC program are scientifically-based and intended to address the supplemental nutritional needs of the Program's target population and can be tailored to meet the needs of individual participants.

In addition, WIC staff can greatly assist families by providing referrals to medical providers and other services, if available, in their community. Such resources may provide the recommended medical assessments, in order to rule out or confirm medical conditions, and offer treatment when necessary and/or in cases where growth improvement is slow to respond to dietary interventions.

Clarification

The cut-off for underweight for infants and children <24 months is 2.3; however, for ease of use, CDC labels it as the 2nd percentile on the hard copy Birth to 24 months growth charts. Electronic charts should use the 2.3rd percentile as the cut-off.

References

1. Centers for Disease Control and Prevention. Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United States. MMWR 2010; 59 (No. RR-9). Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5909a1.htm?s_cid=rr5909a1_w. Accessed September 2010.
2. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics; no. 314. Hyattsville, Maryland: National Center for Health Statistics. 2000.
3. World Health Organization. WHO child growth standards: Length/height-for-age, weight-for-age, weight-for-height and body mass index-for-age: Methods and development. Geneva, Switzerland: World Health Organization; 2006. Available at http://www.who.int/childgrowth/publications/technical report pub/en/index.html. Accessed September 2010.
4. Food and Nutrition Information Center, National Agriculture Library. Update of analysis of literature regarding cut-off percentiles for low weight for length in infants. Washington, D.C.; February 5, 1991.
5. Sherry B. Epidemiology of inadequate growth. In: Kessler DB, Dawson P, Editors. Failure to thrive and pediatric under nutrition: A transdisciplinary approach. Baltimore: Paul H. Brooks Publishing Company, Inc.; 1999.
6. Metallinos-Katsaras E, Gorman KS. Effects of under nutrition on growth and development. Kessler DB, Dawson P, editors. Failure to thrive and pediatric under nutrition: A transdisciplinary approach. Baltimore: Paul H. Brooks Publishing Company, Inc.; 1999. p. 38.
7. Disbrow DD. The costs and benefits of nutrition services: a literature review. J Am Diet Assoc. 1989; 89:S3-66.

Nutrition Risk Criteria **111 ★ OVERWEIGHT WOMAN**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Pregnant: Body Mass Index (BMI) greater than or equal to (≥) 25.

- Use pre-pregnancy weight in BMI calculation.

Breastfeeding/Non-breastfeeding: BMI greater than or equal to (≥) 25.0.

- Use pre-pregnancy weight in BMI calculation for all women who are less than 6 months postpartum.
- Use current weight for breastfeeding women who are equal or greater than 6 months postpartum.

Note: A BMI table is attached to assist in determining weight classifications. Until research supports the use of different BMI cut-offs for adolescent pregnancies, the same BMI cut-offs will be used for all women, regardless of age, when determining WIC eligibility (1).

Required Documentation

Pregnant

- Pre-pregnancy weight
- Height
- BMI

Breastfeeding/Non-breastfeeding

- Pre-pregnancy weight
- Current weight
- Height
- BMI

Justification

Maternal overweight and obesity are associated with higher rates of cesarean delivery, gestational diabetes mellitus, preeclampsia and other pregnancy-induced hypertensive disorders, as well as postpartum anemia (2). Several studies have established an association between obesity and an increased risk for hypertension, dyslipidemia, diabetes mellitus, cholelithiasis, coronary heart disease, osteoarthritis, sleep apnea, stroke and certain cancers (1).

One goal of prenatal nutritional counseling is to achieve recommended weight gain. For the overweight woman, emphasis should be on selecting food choices of high nutritional quality and avoiding calorie rich foods, thereby minimizing further risks associated with increased overweight and obesity.

The 2009 Institute of Medicine (IOM) report: *Weight Gain During Pregnancy: Reexamining the Guidelines* (1) updated the pregnancy weight categories to conform to the categories developed by the World Health Organization and adopted by the National Heart, Lung and Blood Institute in 1998 (3). The reexamination of the guidelines consisted of a review of the determinants of a wide range of short-and long-term consequences of variation in weight gain during pregnancy for both the mother and her infant. The IOM prenatal weight

gain recommendations based on pre-pregnancy weight status categories are associated with improved maternal and child health outcomes (1).

Included in the 2009 IOM guidelines is the recommendation that the BMI weight categories used for adult women be used for pregnant adolescents as well. More research is needed to determine whether special categories are needed for adolescents. It is recognized that the IOM BMI cut-offs for defining weight categories will classify some adolescents differently than the CDC BMI-for-age charts. For the purpose of WIC eligibility determination, the IOM BMI cut-offs will be used for all women regardless of age. However, due to the lack of research on relevant BMI cut-offs for pregnant and postpartum adolescents, professionals should use all of the tools available to them to assess these applicants' anthropometric status and tailor nutrition counseling accordingly.

Weight during the early postpartum period, when most WIC certifications occur, is very unstable. During the first 4-6 weeks fluid shifts and tissue changes cause fluctuations in weight. After 6 weeks, weight loss varies among women. Pre-pregnancy weight, amount of weight gain during pregnancy, race, age, parity and lactation all influence the rate of postpartum weight loss. By 6 months postpartum, body weight is more stable and should be close to the pre-pregnancy weight. In most cases therefore, pre-pregnancy weight is a better indicator of weight status than postpartum weight in the first 6 months after delivery.

The percentage of adolescents who are overweight is increasing rapidly and more than 60% of adults in the U.S. are overweight. Due to the significant impact that overweight and obesity have on morbidity and mortality, it is imperative that every effort be made to identify individuals who are overweight and to assist them in achieving a more healthful weight. The WIC Program is in a position to play an important role in helping to reduce the prevalence of overweight not only by working with postpartum women on improving their own weight status, but also by helping them to see their role in assisting their children to learn healthful eating and physical activity behaviors.

References

1. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines (Prepublication Copy). National Academy Press; Washington, D.C.; 2009. www.nap.edu. Accessed June 2009.
2. Bodnar LM, Catov JM, Klbanoff MA, Ness RB, Roberts JM. Prepregnancy body mass index and the occurrence of severe hypertensive disorders of pregnancy. Epidemiology 2007;18(2):234-239.
3. National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health (NIH). Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. NIH Publication No. 98-4083, 1998. www.nhlbi.nih.gov. Accessed June 2009.
4. Crowell DT. Weight changes in the postpartum period: a review of the literature. Journal of Nurse-Midwifery. Vol. 40, No. 5, September/October 1995; 418-423.

Additional Related References

1. Naye, R.L.: Maternal body weight and pregnancy outcome. American Journal Clin. Nutr.; 1990; 52:273-279.
2. Parker JD, Abrams B. Prenatal weight gain advice: an examination of the recent prenatal weight gain recommendations of the Institute of Medicine. Obstet Gynecol, 1992; 79:664-9.

3. Siega-Riz AM, Adair LS, Hobel CJ. Institute of Medicine maternal weight gain recommendations and pregnancy outcomes in a predominately Hispanic population. Obstet Gynecol, 1994; 84:565-73.
4. Suitor CW, editor. Maternal weight gain: A report of an expert work group. Arlington, Virginia: National Center for Education in Maternal and Child Health; 1997. Sponsored by Maternal and Child Health Bureau, Health Resources and Services Administration, Public Health Service, U.S. Department of Health and Human Services.

Nutrition Risk Criteria **113★ OBESE – CHILDREN (2-5)**

	Category	Priority	Referral
Category, Priority and Referral	CHILDREN	3	RD

Definition

Children Age 2 and Older: Body mass index-for-age greater than or equal to (≥) 95th percentile. Based on Centers for Disease Control and Prevention (CDC) age and sex specific growth charts. *

*The cut-off is based on standing height measurements. Therefore, recumbent length measurements may not be used for this risk. See Clarification for more information.

Required Documentation

DOB
Weight
Height
BMI
CDC growth charts

Justification

The rapid rise in the prevalence of obesity in children and adolescents is one of the most important public health issues in the United States today. The National Health and Nutrition Examination Survey (NHANES) from the mid-1960s to the early 2000s document a significant increase in obesity among children from preschool age through adolescence. These trends parallel a concurrent increase in obesity among adults, suggesting that fundamental shifts occurring in dietary and/or physical activity behaviors are having an adverse effect on overall energy balance (3).

The causes of increased obesity rates in the United States are complex. Both genetic make-up and environmental factors contribute to the obesity risk. Important contributors include a large and growing abundance of calorically dense foods and an increased sedentary lifestyle for all ages. Although obesity tends to run in families, a genetic predisposition does not inevitably result in obesity. Environmental and behavioral factors can influence the development of obesity in genetically at-risk people (3).

BMI is a measure of body weight adjusted for height. While not a direct measure of body fatness, BMI is a useful screening tool to assess adiposity (3). Children ≥ 2 years of age, with a BMI-for-age ≥ 85th and < 95th percentile are considered *overweight* and those at or above the 95th percentile, *obese* (4). Research on BMI and body fatness shows that the majority of children with BMI-for-age at or above the 95th percentile have high adiposity and less than one-half of the children in the 85th to < 95th percentiles have high adiposity (4). Although an imperfect tool, elevated BMI among children most often indicates increased risk for future adverse health outcomes and/or development of diseases (5). BMI should serve as the initial screen and as the starting point for classification of health risks (3).

Use of the 95th percentile to define obesity identifies those children with a greater likelihood of being obese as adolescents and adults, with increased risk of obesity-related disease and mortality. It is recommended that an obese child (≥ 95th percentile) undergo a medical assessment and careful evaluation to identify any underlying health risks or secondary complications (3). Obesity can result from excessive energy intake, decreased energy expenditure, or a medical condition that impairs the regulation of energy metabolism. In addition, obesity in early childhood may signify problematic feeding practices or evolving family behaviors that, if continued, may contribute to health risks in adulthood related to diet and inactivity.

Implications for WIC Nutrition Services

The WIC Program plays an important role in public health efforts to reduce the prevalence of obesity by actively identifying and enrolling young children who may be obese or at risk of overweight/obesity in later childhood or adolescence. When identifying this risk, it is important to communicate with parents/caregivers in a way that is supportive and nonjudgmental, and with a careful choice of words that convey an empathetic attitude and minimize embarrassment or harm to a child's self-esteem (4). In recognition of the importance of language, the 2007 American Medical Association Expert Committee Report recommends the use of the terms *overweight* and *obese* for documentation and risk assessment **only** and the use of more neutral terms (e.g., weight disproportional to height, BMI) when discussing a child's weight with a parent/caregiver (3).

BMI is calculated and plotted on growth charts at each WIC certification. However, growth charts are meant to be used as a screening tool and comprise only one aspect of the overall growth assessment. A clinical assessment to determine if a child is at a healthy weight is more complex. Weight classification (derived from the growth chart) should be integrated with the growth pattern, familial obesity, medical risks and dietary and physical activity habits to determine the child's obesity risk (1, 5).

The goal in WIC nutrition counseling is to help the child achieve recommended rates of growth and development. WIC staff can frame the discussion to make achieving normal growth a shared goal of the WIC Program and the parent/caregiver and make clear that obesity is a medical condition that can be addressed (4). Parents/caregivers of children may need education on recognition of satiety dues and other physiological needs that lead to crying, and ways to comfort a child (holding, reading, rocking) other than by feeding. The foods provided by the WIC Program are scientifically-based and intended to address the supplemental nutritional needs of the program's target population and can be tailored to meet the needs of individual participants. Emphasis can be placed on promoting food choices of high nutritional quality while avoiding unnecessary or excessive amounts of calorie rich foods and beverages, and reducing inactivity (like decreasing sedentary TV viewing).

Beliefs about what is an attractive or healthy weight, the importance of physical activity, what foods are desirable or appropriate for parents to provide to children, family mealtime routines, and many other lifestyle habits are influenced by different cultures, and should be considered during the nutrition assessment and counseling (6). The following resources for obesity prevention can be found:

- Fit WIC Materials: http://www.nal.usda.gov/wicworks/Sharing_Center/gallery/foodfunfamilies.htm.
- MyPyramid for Preschoolers:

http://www.mypyramid.gov/preschoolers/index.html

In addition, WIC staff can greatly assist families by providing referrals to medical providers and other services, if available, in their community. Such resources may provide the recommended medical assessments, in order to rule out or confirm medical conditions, and offer treatment when necessary and/or in cases where growth improvement is slow to respond to dietary interventions.

Clarification

The 2000 CDC Birth to 36 Months growth charts cannot be used as a screening tool for the purpose of assigning this risk because these charts are based on recumbent length rather than standing height data. However, these charts may be used as an assessment tool for evaluating growth in children aged 24-36 months who are not able to be measured for the standing height required for the 2000 CDC 2-20 Years growth charts.

References

1. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics; no. 314. Hyattsville, Maryland: National Center for Health Statistics. 2000.
2. Grummer-Strawn LM, Reinold C, Krebs NF. Use of World Health Organization and CDC growth charts for children aged 0-59 Months in the United States. CDC Morbidity and Mortality Weekly Report (September 2010); no 59 (rr09); 1-15. Available at: http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5909a1.htm?s_cid=rr5909a1_w. Accessed September 2010.
3. Barlow, SE, Expert Committee. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. Pediatrics 2007; 120; S164-S192.
4. Ogden CL, Flegal KM. Changes in Terminology for childhood overweight and obesity. National health statistics reports; no. 25. Hyattsville (MD): National Center for Health Statistics. 2010.
5. U.S. Department of Health and Human Services. The Surgeon General's vision for a healthy and fit nation. Rockville (MD): U.S. Department of Health and Human Services, Office of the Surgeon General. 2010.
6. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. Pediatrics 2007; 120 Suppl 4: S103-S228.

Nutrition Risk Criteria **114★ OVERWEIGHT – CHILDREN (2-5)**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	CHILDREN	3	-

Definition

Overweight:

Children age 2 and older: Being ≥ 24 months of age and ≥ 85th and < 95th percentile Body Mass Index (BMI)*-for-age as plotted on the 2000 Centers for Disease Control and Prevention (CDC) 2-20 Years gender specific growth charts (1, 2).*

*DO NOT USE if NRC 113 Obese Children applies

*The cut-off is based on standing height measurements. Therefore, recumbent length measurements may not be used to determine this risk. See Clarification for more information.

Required Documentation

DOB
Weight
Height
BMI
CDC growth charts

Justification

The rise in the prevalence of overweight in children and adolescents is one of the most important public health issues in the United States today. The National Health and Nutrition Examination Survey (NHANES) from the mid-1960s to the early 2000s document a significant increase in overweight among children from preschool age through adolescence. These trends parallel a concurrent increase in overweight among children from preschool age through adolescence. These trends parallel a concurrent increase in obesity among adults, suggesting that fundamental shifts in dietary and/or physical activity behaviors are having an adverse effect on overall energy balance (3).

BMI is a measure of body weight adjusted for height. While not a direct measure of body fatness, BMI is a useful screening tool to assess adiposity (3). Children ≥ 2 years of age, with a BMI-for-age ≥ 85th and < 95th percentile are considered overweight and those at or above the 95th percentile, *obese* (4). Research on BMI and body fatness shows that the majority of children with BMI-for-age at or above the 95th percentile have high adiposity and less than one-half the children in the 85th to < 95th percentiles have high adiposity (4). Although an imperfect tool, elevated BMI among children most often indicates increased risk for future adverse health outcomes and/or development of diseases (5). BMI should serve as the initial screen and as the starting point for classification of health risks (3).

Increasingly, attention is being focused on the need for comprehensive strategies that focus on preventing overweight/obesity and a sedentary lifestyle for all ages. Scientific evidence suggests that the presence of obesity in a parent greatly increases the risk of overweight in preschoolers, even when no other

overt signs of increasing body mass are present (6). The presence of parental obesity should lead to greater efforts by nutrition services staff to assist families in establishing or improving healthy behaviors (3).

Implications for WIC Nutrition Services

The WIC Program plays an important role in public health efforts to reduce the prevalence of obesity by actively identifying and enrolling infants and children who may be overweight or at risk of overweight in childhood or adolescence. When identifying this risk, it is important to communicate it in a way that is supportive, nonjudgmental, and with a careful choice of words to convey an empathetic attitude and to minimize embarrassment or harm to a child's self-esteem (4). In recognition of the importance of language, the 2007 American Medical Association expert committee report recommends the use of the terms *overweight* and *obese* for documentation and risk assessment **only** and the use of more neutral terms (e.g., *weight disproportional to height, excess weight, BMI*) when discussing a child's weight with a parent/caregiver (3).

BMI is calculated and plotted on growth charts at each WIC certification. However, growth charts are meant to be used as a screening tool and comprise only one aspect of the overall growth assessment. A clinical assessment to determine if a child is at a healthy weight is more complex. Weight classification (derived from the growth chart) should be integrated with the growth pattern, familial obesity, medical risks, and dietary and physical activity habits to determine the child's obesity risk (1, 5).

The goal in WIC nutrition counseling is to help the child achieve recommended rates of growth and development. WIC staff can frame the discussion to make achieving normal growth a shared goal of the WIC Program and the parent/caregiver. Studies have shown that the early childhood eating environment provides a great opportunity for preventive intervention (7). Parents/caregivers of infants and toddlers may need recognition of satiety cues and other physiological needs that lead to crying, and ways to comfort a child (holding, reading, rocking) other than by feeding. Young children look upon their parents as role models for eating behaviors. Through client-centered counseling, WIC staff can emphasize the importance of prevention and can assist families in making changes that improve parenting skills that promote healthy eating, and physical activity behaviors and a healthy weight in children. Also, the foods provided by the WIC Program are scientifically-based and intended to address the supplemental nutritional needs of the Program's target population and can be tailored to meet the needs of individual participants.

Beliefs about what is an attractive or healthy weight, the importance of physical activity, what foods are desirable or appropriate for parents to provide children, family mealtime routines, and many other lifestyle habits are influenced by different cultures and should be considered during the nutrition assessment and counseling (8). The following resources for obesity prevention can be found at:

- Fit WIC Materials
  http://www.nal.usda.gov/wicworks/Sharing_Center/gallery/foodfunfamilies.htm
- MyPyramid for Preschoolers:
  http://www.mypyramid.gov/preschoolers/index.html

In addition, WIC staff can greatly assist families by providing referrals to medical providers and other services, if available, in their community. Such resources may provide the recommended medical assessments, in order to rule out or

confirm medical conditions, and offer treatment when necessary and/or in cases where growth improvement is slow to respond to dietary interventions.

Clarification

The 2000 CDC Birth to 36 months growth charts cannot be used as a screening tool for the purpose of assigning this risk because these charts are based on recumbent length rather than standing height data. However, these charts may be used as an assessment tool for evaluating growth in children aged 24-36 months who are not able to be measured for the standing height required for the 2000 CDC 2-20 years growth charts.

References

1. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics; no. 314. Hyattsville, Maryland: National Center for Health Statistics. 2000.
2. Grummer-Strawn LM, Reinold C, Krebs NF. Use of World Health Organization and CDC growth charts for children aged 0-59 Months in the United States. CDC Morbidity and Mortality Weekly Report (September 2010); no 59 (rr09); 1-15. Available at: http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5909a1.htm?s_cid=rr5909a1_w. Accessed September 2010.
3. Barlow SE, Expert Committee. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. Pediatrics. 2007; 120; S164-S192.
4. Ogden CL, Flegal KM. Changes in Terminology for childhood overweight and obesity. National health statistics reports; no. 25. Hyattsville (MD): National Center for Health Statistics. 2010.
5. U.S. Department of Health and Human Services. The Surgeon General's vision for a healthy and fit nation. Rockville (MD): U.S. Department of Health and Human Services, Office of the Surgeon General. 2010.
6. Whitaker RC, Wright JA, Pepe MS, Seidel KD, Dietz WH. Predicting obesity in young adulthood from childhood and parental obesity. N.Eng.J.Med. September 25,1997; Vol 337; No 13: pgs 869-873.
7. Anzman SL, Rolllins BY, Birch LL. Parental influence on children's early eating environments and obesity risk: implications for prevention. International Journal of Obesity 34, 1116-1124 (July 2010).
8. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. Pediatrics 2007; 120 Suppl 4: S103-S228.

Nutrition Risk Criteria **115** **HIGH WEIGHT-FOR-LENGTH (INFANTS AND CHILDREN <24 MONTHS OF AGE)**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	-	-
CHILDREN	3	-

Definition

High weight-for-length for infants and children < 24 months of age is defined as follows:

Age	Cut-Off Value
Birth to less than (<) 24 months	Greater than or equal to (≥) 97.7th percentile weight-for-length as plotted on the Centers for Disease Control and Prevention (CDC) Birth to 24 months gender specific growth charts (1) (available at: www.cdc.gov/growthcharts).*
**Based on the 2006 World Health Organization (WHO) international growth standards (2). CDC labels the 97.7th percentile as the 98th percentile on the Birth to 24 months gender specific growth charts. For more information about the percentile cut-off, please see Clarification.*	

Required Documentation

DOB
Weight
Length
CDC Growth Charts (based on the WHO growth standards)

Justification

The WHO and CDC growth charts are similar in that both describe weight-for-age, length (or stature)-for-age, weight-for-length (or stature) and body mass index (BMI) for age. However, they differ in the approach taken to create the growth charts. The WHO growth charts are growth standards that describe how healthy children grow under optimal environmental and health conditions. The 2000 CDC charts are a growth reference, not a standard, and describe how certain children grew in a particular place and time (2).

The WHO growth standards for children < 24 months are based on data collected from 1997-2003 in 6 countries (including the U.S.), from children who were born between 37 and 42 weeks gestation, breastfed for at least 12 months, and introduced to complementary food by at least 6 months but not before 4 months. Infants and children of low-income mothers and/or mothers who smoked were not included in the data sample (2).

The 2000 CDC charts for infants and children < 36 months are based on birth weight (from 1968 to 1980 and from 1985 to 1994) and birth length data (from 1989 to 1994) obtained from U.S. birth certificates, National Health and Nutrition Examination Survey (NHANES) data, and measurements from infants who had been breastfed and formula fed (approximately 50% ever breastfed and approximately 33% who were still breastfeeding at 3 months). Very low birth

weight infants were not included in the sample population. This was the only exclusion criterion applied to the sample population (2, 3).

Prior to making its recommendation, CDC convened an Expert Panel with the National Institutes of Health and the American Academy of Pediatrics to review the scientific evidence and discuss the potential use of the WHO growth standards in the U.S. The recommendation to use WHO growth standards for infants and children < 24 months was made on the basis of input from the Expert Panel. In addition, CDC concluded that the WHO growth standards are based on a high quality study and, since breastfeeding is the recommended infant feeding practice, it is appropriate to use the breastfed infant as the standard against which all other infants are compared (2).

The WHO growth standards use values of 2 standard deviations away from the median to identify children whose growth might be indicative of adverse health conditions (1). The CDC Birth to 24 months growth charts (based on the WHO growth standards) labels 2 standard deviations above the median as the 97.7th percentile. Thus, an infant or child (< 24 months) is categorized as high weight-for-length when plotted at or above the 97.7th percentile, labeled as the 98th percentile on the CDC Birth to 24 months growth charts. The CDC recommends that all infants and children < 24 months be assessed using the CDC Birth to 24 months growth charts regardless of type of feeding (formula or breastfed) (2). (See clarification for information about standard deviations and the cut-off used to determine high weight-for-length.)

In 2006, WHO released international growth standards for infants and children aged 0-59 months (2), similar to the 2000 CDC growth references. Since then, the CDC has developed Birth to 24 months growth charts, based on the WHO growth standards, and recommends their use in the United States (1). For persons 2-20 years, the 2000 CDC growth charts will continue to be used (1).

Implications for WIC Nutrition Services

The WIC Program plays an important role in public health efforts to reduce the prevalence of obesity by actively identifying and enrolling infants and young children who may be at risk of overweight/obesity in later childhood or adolescence. When identifying this risk, it is important to communicate with parents/caregivers in a way that is supportive and nonjudgmental, and with a careful choice of words that convey an empathetic attitude and minimize embarrassment or harm to a child's self-esteem (4). In recognition of the importance of language, the 2007 American Medical Association Expert Committee Report recommends the use of more neutral terms such as *weight disproportional to height*, *excess weight*, and *high weight-for-length* when communicating with a parent/caregiver (5).

Height and weight measurements are plotted on growth charts at each WIC certification. However, growth charts are meant to be used as a screening tool and comprise only one aspect of the overall growth assessment. A clinical assessment to determine if a child is at a healthy weight is more complex. Weight classification (derived from the growth chart) should be integrated with the growth pattern, familial obesity, medical risks, and dietary and physical activity habits to determine the child's obesity risk (3, 6).

The goal in WIC nutrition counseling is to help the child achieve recommended rates of growth and development. WIC staff can frame the discussion to make achieving normal growth a shared goal of the WIC Program and the parent/caregiver. Studies have shown that the early childhood eating

environment provides a great opportunity for preventive intervention (7). Parents/caregivers of infants and toddlers may need education on recognition of satiety cues and other physiological needs that lead to crying, and ways to comfort a child (holding, reading, rocking) other than by feeding. Young children look upon their parents as role models for eating behaviors. Through client-centered counseling, WIC staff can emphasize the importance of prevention and can assist families in making changes that improve parenting skills that promote healthy eating, physical activity behaviors and a healthy weight in children. Also, the foods provided by the WIC Program are scientifically-based and intended to address the supplemental nutrition needs of the program's target population and can be tailored to meet the needs of individual participants.

Beliefs about what is an attractive or healthy weight, the importance of physical activity, what foods are desirable or appropriate for parents to provide to children, family mealtime routines, and many other lifestyle habits are influenced by different cultures, and should be considered during the nutrition assessment and counseling (8). The following resources for obesity prevention can be found:

- Fit WIC Materials: http://www.nal.usda.gov/wicworks/Sharing_Center/gallery/foodfunfamilies.htm.
- MyPyramid for Preschoolers: http://www.mypyramid.gov/preschoolers/index.html.

In addition, WIC staff can greatly assist families by providing referrals to medical providers and other services, if available, in their community. Such resources may provide the recommended medical assessments, in order to rule out or confirm medical conditions, and offer treatment when necessary and/or in cases where growth improvement is slow to respond to dietary interventions.

Clarification

Standard deviation is a measurement widely used in statistical analysis. It shows how much variation there is from the median. The WHO growth charts use standard deviations to illustrate the proximity of a given child's growth from that of the average child of the same age and gender. For infants and children < 24 months of age, 2 standard deviations above the median indicates high weight-for-length. A measurement of 2 standard deviations below the median indicates underweight. Since most health care providers in the U.S. are more familiar with percentiles, the CDC developed growth charts based on the WHO growth standards but converted standard deviations into percentile readings. Two standard deviations above the median is the 97.7th percentile; however, for ease of use, CDC labels it as the 98th percentile on the hard copy Birth to 24 months growth charts. Electronic charts should use the 97.7th percentile as the cut-off.

References

1. Centers for Disease Control and Prevention. Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United States. MMWR 2010; 59 (No. RR-9). Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5909a1.htm?s_cid=rr5909a1_w. Accessed September 2010.
2. World Health Organization. WHO child growth standards: Length/height-for-age, weight-for-age, weight for height and body mass index-for-age: Methods and development. Geneva, Switzerland: World Health Organization; 2006. Available at

http://www.who.int/childgrowth/publications/technical_report_pub/en/index.html. Accessed September 2010.
3. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics; no. 314. Hyattsville, Maryland: National Center for Health Statistics, 2000.
4. Ogden CL, Flegal KM. Changes in Terminology for childhood overweight and obesity. National health statistics reports; no. 25. Hyattsville (MD): National Center for Health Statistics. 2010.
5. Barlow SE, Expert Committee. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. Pediatrics. 2007; 120; S164-S192.
6. U.S. Department of Health and Human Services. The Surgeon General's vision for a healthy and fit nation. Rockville (MD): U.S. Department of Health and Human Services, Office of the Surgeon General. 2010.
7. Anzman SL, Rolllins BY, Birch LL. Parental influence on children's early eating environments and obesity risk: implications for prevention. International Journal of Obesity 34, 1116-1124 (July 2010).
8. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. Pediatrics 2007; 120 Suppl 4: S103-S228.

Nutrition Risk Criteria **121★ SHORT STATURE OR AT RISK OF SHORT STATURE-INFANTS/CHILDREN**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	-
CHILDREN	3	-

Definition

Short Stature
Birth up to 2 years: Less than or equal (≤) to 2.3rd percentile length-for-age as plotted on the Centers for Disease Control and Prevention (CDC) Birth to 24 months gender specific growth charts (1).*

2 to 5 years: Less than or equal (≤) to 5th percentile stature (height)-for-age as plotted on the 2000 CDC age/gender specific growth charts (2).

At Risk of Short Stature
Birth up to 2 years: Greater than (>) 2.3rd percentile and less than or equal (≤) 5th percentile length-for-age as plotted on the CDC Birth to 24 months gender specific growth charts (1).*

2 to 5 years: Greater than (>) 5th percentile and less than or equal (≤) 10th percentile stature-for-age as plotted on the 2000 CDC age/gender specific growth charts (2).

*Based on 2006 World Health Organization international growth standards (3). CDC labels the 2.3rd percentile as the 2nd percentile on the Birth to 24 months gender specific growth charts. For more information about the percentile cut-off, please see Clarification.

Notes:

1. The Birth to 24 months and the 2000 CDC growth charts are available at: www.cdc.gov/growthcharts.
2. For premature infants and children (with a history of prematurity) up to 2 years of age, assignment of this risk criterion will be based on adjusted gestational age. For information about adjusting for gestational age see: Guidelines for Growth Charts and Gestational Age Adjustment for Low Birth Weight and Very Low Birth Weight Infants.

Required Documentation

DOB
Height
CDC growth charts, CDC growth charts based on the WHO growth standards

Justification

The CDC uses the 2.3rd percentile (for birth to 24 months of age) and the 5th percentile (for 2-5 years of age) stature-for-age, as the cut-offs to define short stature in its Pediatric Nutrition Surveillance System (1, 2). However, CDC does not have a position regarding the cut-off percentile which should be used to determine *at risk of short stature* as a nutritional risk in the WIC Program. *At risk of short stature* is included in this criterion to reflect the preventive emphasis of the WIC Program.

Abnormally short stature in infants and children is widely recognized as a response to an inadequate nutrient supply at the cellular level (4). This indicator can help identify children whose growth is stunted due to prolonged under nutrition or repeated illness (3). Short stature is related to the lack of total dietary energy and to a poor dietary quality that provides inadequate protein, particularly animal protein, and inadequate amounts of micronutrients such as zinc, vitamin A, iron, copper, iodine, calcium, and phosphorus (4). In these circumstances, maintenance of basic metabolic functions takes precedence, and thus resources are diverted from linear growth.

Demonstrable differences in stature exist among children of different ethnic and racial groups. However, racial and ethnic differences are relatively minor compared with environmental factors (1). Growth patterns of children of racial groups whose short stature has traditionally been attributed to genetics have been observed to increase in rate and in final height under conditions of improved nutrition (5, 6).

Short stature may also result from disease conditions such as endocrine disturbances, inborn errors of metabolism, intrinsic bone diseases, chromosomal defects, fetal alcohol syndrome, and chronic systemic diseases (4).

Implications for WIC Nutrition Services

Participation in WIC has been associated with improved growth in both weight and height in children (7). A more in-depth dietary assessment and/or referral to a health care provider may be necessary to determine if short stature is a result of dietary inadequacy or a disease condition. Also, more frequent follow-up to monitor growth is appropriate for children in these categories. Through client-centered counseling WIC staff can assist families in improving dietary intake to promote healthy growth and development. In addition, the foods provided by the WIC Program are scientifically-based and intended to address the supplemental nutritional needs of the Program's target population and can be tailored to meet the needs of the individual participants.

In addition, WIC staff can greatly assist families by providing referrals to medical providers and other services, if available, in their community. Such resources may provide the recommended medical assessments, in order to rule out or confirm medical conditions, and offer treatment when necessary and/or in cases where growth improvement is slow to respond to dietary interventions.

Clarification

The cut-off for short stature for infants and children > 24 months is 2.3; however, for ease of use, CDC labels it as the 2nd percentile on the Birth to 24 months hard copy growth charts. Electronic charts should use the 2.3rd percentile as the cut-off.

References

1. Centers for Disease Control and Prevention. Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United States. MMWR 2010; 59 (No. RR-9). Available at: http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5909a1.htm?s_cid=rr5909a1_w. Accessed September 2010.
2. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al. CDC growth charts: United States. Advance data from vital and health statistics; no. 314. Hyattsville, Maryland: National Center for Health Statistics. 2000.

3. World Health Organization. WHO child growth standards: Length/height-for-age, weight-for-age, weight for height and body mass index-for-age: Methods and development. Geneva, Switzerland: World Health Organization; 2006. Available at: http://www.who.int/childgrowth/publications/technical_report_pub/en/index.html. Accessed September 2010.
4. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. Washington (DC): National Academy Press; 1996. p. 104-109.
5. Pipes PL, Trahms CM. Nutrition in infancy and childhood, 6th edidition. Seattle (WA): WCB/McGraw-Hill; 1997. p. 2.
6. Berhane R, Dietz WH. Clinical assessment of growth. In: Kessler DB, Dawson P., editors. Failure to thrive and pediatric under nutrition: A transdisciplinary approach. Baltimore (MD): Paul H. Brooks Publishing Company, Inc.; 1999. p. 199.
7. Disbrow DD. The costs and benefits of nutrition services: a literature review. J Am Diet Assoc. 1989; 89:S3-66.

Nutrition Risk Criteria **131★ LOW MATERNAL WEIGHT GAIN**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT	1	RD

Definition

Low maternal weight gain is defined as:

1. A low rate of weight gain, such that in the 2nd and 3rd trimesters, for singleton pregnancies (1):
   - Underweight women gain less than an average of 1 lb per week
   - Normal weight women gain less than an average of .8 lbs per week
   - Overweight women gain less than an average of .5 lbs per week
   - Obese women gain less than an average of .4 lbs per week

OR

2. Low weight gain at any point in pregnancy, such that using an Institute of Medicine (IOM)-based weight gain grid, a pregnant woman's weight plots at any point beneath the bottom line of the appropriate weight gain range for her respective prepregnancy weight category (1), as follows:

Prepregnancy Weight Groups	Definition (BMI)	Total Weight Gain Range (lbs)
Underweight	< 18.5	28-40
Normal Weight	18.5 to 24.9	25-35
Overweight	25.0 to 29.9	15-25
Obese	≥ 30.0	11-20

Multi-fetal Pregnancies: See Justification for information.

Note: A BMI table is attached to assist in determining weight classifications. Also, until research supports the use of different BMI cut-offs to determine weight categories for adolescent pregnancies, the same BMI cut-offs to determine weight categories for adolescent pregnancies, the same BMI cut-offs will be used for all women, regardless of age, when determining WIC eligibility.

Required Documentation

Self-reported weight loss is acceptable as long as it meets the criteria.

Pre-pregnancy weight
Height
Weeks gestation
Amount of weight gained to date

Justification

Maternal weight gain during the 2nd and 3rd trimesters is an important determinant of fetal growth. Low maternal weight gain is associated with an

increased risk of small for gestational age (SGA) infants, especially in underweight and normal-weight women (1). In addition, low maternal weight gain is associated with failure to initiate breastfeeding and preterm birth among underweight and to a lesser extent normal weight women (1).

The 2009 Institute of Medicine (IOM) report: Weight Gain During Pregnancy: Reexamining the Guidelines (1) updated the pregnancy weight categories to conform to the categories developed by the World Health Organization and adopted by the National Heart, Lung and Blood Institute in 1998 (2). The reexamination of the guidelines consisted of a review of the determinants of a wide range of short and long term consequences of variation in weight gain during pregnancy for both the mother and her infant. The IOM prenatal weight gain recommendations based on prepregnancy weight status categories are associated with improved maternal and child health outcomes (1).

Included in the 2009 IOM guidelines is the recommendation that the BMI weight categories used for adult women be used for pregnant adolescents as well. More research is needed to determine whether special categories are needed for adolescents.

It is recognized that the IOM BMI cut-offs for defining weight categories will classify some adolescents differently than the CDC BMI-for-age charts. For the purpose of WIC eligibility determination, the IOM BMI cut-offs will be used for all women regardless of age. However, due to the lack of research on relevant BMI cut-offs for pregnant and postpartum adolescents, professionals should use all of the tools available to them to assess these applicants' anthropometric status and tailor nutrition counseling accordingly.

For twin gestations, the 2009 IOM recommendations provide provisional guidelines; normal-weight women should gain 37-54 lbs; overweight women, 31-50 lbs; and obese women, 25-42 lbs. There was insufficient information for the IOM committee to develop even provisional guidelines for underweight women with multiple fetuses (1). A consistent rate of weight gain is advisable. A gain of 1.5 lbs per week during the second and third trimesters has been associated with a reduced risk of preterm and low-birth weight delivery in twin pregnancy (3). In triplet pregnancies the overall gain should be around 50 lbs with a steady rate of gain of approximately 1.5 lbs per week throughout the pregnancy (3). For WIC eligibility determinations, multi-fetal pregnancies are considered a nutrition risk in and of themselves (ID NRC 61, USDA Code 335, Multi-Fetal Gestation), aside from the weight gain issue.

The supplemental foods, nutrition education, and counseling related to the weight gain guidelines provided by the WIC Program may improve maternal weight status and infant outcomes (4).

Clarification

The Centers for Disease Control and Prevention (CDC) defines a trimester as a term of three months in the prenatal gestation period with the specific trimesters defined as follows in weeks:

First Trimester: 0-13 weeks
Second Trimester: 14-26 weeks
Third Trimester: 27-40 weeks

Further, CDC begins the calculation of weeks starting with the first day of the last menstrual period. If that date is not available, CDC estimates that date from the

estimated date of confinement (EDC). This definition is used in interpreting CDC's Prenatal Nutrition Surveillance System data, comprised primarily of data on pregnant women participating in the WIC Program

References

1. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines (Prepublication Copy). National Academy Press, Washington, D.C.; 2009. www.nap.edu. Accessed June 2009.
2. National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health (NIH). Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults, NIH Publication No.: 98-4083, 1998. www.nhlbi.nih.gov, Accessed June 2009.
3. Brown JE and Carlson M. Nutrition and multifetal pregnancy. J Am Diet Assoc. 2000; 100:343-348.
4. Institute of Medicine. (1996) *WIC Nutrition Risk Criteria: A Scientific Assessment.* National. Academy Press, Washington, D.C.

Additional Related References

1. Brown JE, Schloesser PT. Pregnancy weight status, prenatal weight gain, and the outcome of term twin gestation. Am. J. Obstet. Gynecol. 1990; 162:182-6.
2. Parker JD, Abrams B. Prenatal weight gain advice: an examination of the recent prenatal weight gain recommendations of the Institute of Medicine. Obstet Gynecol, 1992; 79:664-9.
3. Siega-Riz AM, Adair LS, Hobel CJ. Institute of Medicine maternal weight gain recommendations and pregnancy outcomes in a predominately Hispanic population. Obstet Gynecol, 1994; 84:565-73.
4. Suitor, CW. (1997) Maternal Weight Gain: A Report of an Expert Work Group. Arlington, VA: National Center for Education in Maternal and Child Health. Sponsored by Maternal and Child Health Bureau. Health Resources and Services Administration, Public Health Service, U.S. Department of Health and Human Services.
5. Williams RL, Creasy RK, Cunningham GC< Hawes WE, Norris FD, Tashiro M. Fetal growth and perinatal viability in California. Obstet. Gynecol. 1982; 59:624-32.

Nutrition Risk Criteria **132★ MATERNAL WEIGHT LOSS DURING PREGNANCY**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	PREGNANT WOMEN	1	RD

Definition

FIRST TRIMESTER (weeks 1-13) Any weight loss below pre-pregnancy weight.
SECOND TRIMESTER (weeks 14-26) Weight loss of 2 lbs or greater.
THIRD TRIMESTER (weeks 27-40+) Weight loss 2 lbs or greater.

Required Documentation

Self-reported weight loss is acceptable as long as it meets the criteria
Pre-pregnancy
Height
Weeks gestation
Amount of weight gained to date

Justification

Weight loss during pregnancy may indicate underlying dietary or health practices or health or social conditions associated with poor pregnancy outcomes. These outcomes could be improved by the supplemental food, nutrition education, and referrals provided by the WIC Program.

References

1. Brown JE. Prenatal weight gain considerations for WIC. Final report. Commissioned by the Risk Identification and Selection Collaborative. 1998.
2. Centers for Disease Control and Prevention. Prenatal Nutrition Surveillance System User's Manual. Atlanta: CDC, 1994.
3. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
4. Metropolitan Life Insurance Company. New weight standards for men and women. Stat.Bull.Metrop.Life Insur.Co., 1959.

Nutrition Risk Criteria **133★ HIGH MATERNAL WEIGHT GAIN**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Pregnant Women

1. A high rate of weight gain, such that in the 2nd and 3rd trimesters, for singleton pregnancies (1):
   - Underweight women gain more than 1.3 lbs per week
   - Normal weight women gain more than 1 lb per week
   - Overweight women gain more than .7 lbs per week
   - Obese women gain more than .6 lbs per week

OR

2. High weight gain at any point in pregnancy, such that using an Institute of Medicine (IOM)-based weight gain grid, a pregnant woman's weight plots at any point above the top line of the appropriate weight gain range for her respective prepregnancy weight category.

Breastfeeding or Non-breastfeeding Women (most recent pregnancy only): total gestational weight gain exceeding the upper limit of the IOM's recommended range (2) based on Body Mass Index (BMI) for singleton pregnancies, as follows (1):

Prepregnancy Weight Groups	Definition (BMI)	Cut-off Value
Underweight	< 18.5	> 40 lbs
Normal Weight	18.5 to 24.9	> 35 lbs
Overweight	25.0 to 29.9	> 25 lbs
Obese	≥ 30.0	> 20 lbs

Multifetal Pregnancies: See Justification for information.

Note: A BMI table is attached to assist in determining weight classification. Also, until research supports the use of different BMI cut-offs to determine weight categories for adolescent pregnancies, the same BMI cut-offs will be used for all women, regardless of age, when determining WIC eligibility. (See Justification for a more detailed explanation.)

Required Documentation

Self-reported weight gain is acceptable as long as it meets the criteria.

Pre-pregnancy weight
Height
Weeks gestation

Amount of weight gained to date

Justification

Women with excessive gestational weight gains are at increased risk for cesarean delivery and delivering large for gestational age infants that can secondarily lead to complications during labor and delivery. There is a strong association between higher maternal weight gain and both postpartum weight retention and subsequent maternal obesity. High maternal weight gain may be associated with glucose abnormalities and gestational hypertension disorders, but the evidence is inconclusive. (1)

Childhood obesity is one of the most important long-term child outcomes related to high maternal weight gain. A number of epidemiologic studies show that high maternal weight gain is associated with childhood obesity as measured by BMI. (1)

The 2009 Institute of Medicine (IOM) report: *Weight Gain During Pregnancy: Reexamining the Guidelines* (1) updated the pregnancy weight categories to conform to the categories developed by the World Health Organization and adopted by the National Heart, Lung and Blood Institute in 1998 (2). The reexamination of the guidelines consisted of a review of the determinants of a wide range of short and long term consequences of variation in weight gain during pregnancy for both the mother and her infant. The IOM prenatal weight gain recommendations based on prepregnancy weight status categories are associated with improved maternal and child health outcomes (1).

Included in the 2009 IOM guidelines is the recommendation that the BMI weight categories used for adult women be used for pregnant adolescents as well. More research is needed to determine whether special categories are needed for adolescents.

It is recognized that both the IOM BMI cut-offs for defining weight categories will classify some adolescents differently than the CDC BMI-for-age charts. For the purpose of WIC eligibility determination, the IOM BMI cut-offs will be used for all women regardless of age. However, due to the lack of research on relevant BMI cut-offs for pregnant and postpartum adolescents, professionals should use all of the tools available to them to assess these applicants' anthropometric status and tailor nutrition counseling accordingly.

For twin gestations, the 2009 IOM recommendations provide provisional guidelines: normal weight women should gain 37-54 lbs; overweight women, 31-50 lbs; and obese women, 25-42 lbs. There was insufficient information for the IOM committee to develop even provisional guidelines for underweight women with multiple fetuses (1). A consistent rate of weight gain is advisable. A gain of 1.5 lbs per week during the second and third trimesters has been associated with a reduced risk of preterm and low-birth weight delivery in twin pregnancy (3). In triplet pregnancies the overall gain should be around 50 lbs with a steady rate of gain of approximately 1.5 lbs per week throughout pregnancy (3). Education by the WIC nutritionist should address a steady rate of weight gain that is higher than for singleton pregnancies. For WIC eligibility determinations, multi-fetal pregnancies are considered a nutrition risk in and of themselves (ID NRC 61, USDA Code 335, Multi-Fetal Gestation), aside from the weight gain issue.

The supplemental foods, nutrition education, and counseling related to the weight gain guidelines provided by the WIC Program may improve maternal weight status and infant outcomes (1). In addition, WIC nutritionists can play an important role, through nutrition education and physical activity promotion, in assisting postpartum women achieve and maintain a healthy weight.

Clarification

The Centers for Disease Control and Prevention (CDC) defines a trimester as a term of three months in the prenatal gestation period with the specific trimesters defined as follows in weeks:

First Trimester: 0-13 weeks
Second Trimester: 14-26 weeks
Third Trimester: 27-40 weeks

Further, CDC begins the calculation of weeks starting with the first day of the last menstrual period. If that date is not available, CDC estimates that date from the estimated date of confinement (EDC). This definition is used in interpreting CDC's Prenatal Nutrition Surveillance System data, comprised primarily of data on pregnant women participating in the WIC Program.

References

1. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines (Prepublication Copy), National Academy Press, Washington, D.C.; 2009. www.nap.edu. Accessed June 2009.
2. National Heart, Lung, and Blood Institute (NHLBI), National Institutes of Health (NIH), Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. National Institutes of Health Publication No.: 98-4083, 1998. www.nhibi.nih.gov. Accessed June 2009.
3. Brown JE and Carlson M. Nutrition and multifetal pregnancy. J Am Diet Assoc. 2000; 100:343-348.
4. Institute of Medicine. WIC nutrition risk criteria: a scientific assessment. National Academy Press, Washington, D.C., 1996.

Additional Related References

1. Carmichael S, Abrams B, Selvin S. The pattern of maternal weight gain in women with good pregnancy outcomes. Am. J. Pub. Hlth. 1997; 87; 12:1984-1988.
2. Brown JE, Schlosser PT. Pregnancy weight status, prenatal weight gain, and the outcome of term twin gestation. Am. J. Obstet. Gynecol. 1990; 162:182-6.
3. Parker JD, Abrams B. Prenatal weight gain advice: an examination of the recent prenatal weight gain recommendations of the Institute of Medicine. Obstet Gynecol 1992; 79:664-9.
4. Siega-Riz AM, Adair LS, Hobel CJ. Institute of Medicine maternal weight gain recommendations and pregnancy outcomes in a predominately Hispanic population. Obstet Gynecol, 1994; 84:565-73.
5. Suitor CW, editor. Maternal weight gain: A report of an expert work group. Arlington, Virginia: National Center for Education in Maternal and Child Health; 1997. Sponsored by Maternal and Child Health Bureau, Health Resources and Services Administration, Public Health Service, U.S. Department of Health and Human Services.
6. Waller K. Why neural tube defects are increased in obese women. Contemporary OB/GYN 1997; p. 25-32.

Nutrition Risk Criteria **134** **FAILURE TO THRIVE**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD
CHILDREN	3	RD

Definition

Presence of failure to thrive (FTT) diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Note: For premature infants with a diagnosis of FTT also see: "Guidelines for Growth Charts and Gestational Age Adjustment for Low Birth Weight and Very Low Birth Weight Infants" (FNS Policy Memorandum 98-9, Revision 7, April 2004).

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Failure to thrive (FTT) is a serious growth problem with an often complex etiology. Some of the indicators that a physician might use to diagnose FTT include:

- weight consistently below the 3rd percentile for age;
- weight < 80% of ideal weight for height/age;
- progressive fall-off in weight to below the 3rd percentile; or
- a decrease in expected rate of growth along the child's previously defined growth curve irrespective of its relationship to the 3rd percentile.

FTT may be a mild form of Protein Energy Malnutrition (PEM) that is manifested by a reduction in rate of somatic growth. Regardless of the etiology of FTT, there is inadequate nutrition to support weight gain.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Berkow R, Fletcher AJ. The Merck manual of diagnosis and therapy. Rahway, N.J.: Merck Sharp & Dohme Research Laboratories, 1992.
2. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Nutrition Risk Criteria **135★** SLOWED/FALTERING GROWTH PATTERN (FORMERLY INADEQUATE GROWTH)

Category, Priority and Referral

Category	Priority	Referral
INFANTS ≤ 6 months age	1	RD

Definition

Slowed/Faltering Growth is defined as:

Age	Cut-Off Value
Infants Birth to 2 Weeks	Excessive weight loss after birth, defined as ≥ 7% of birth weight (1, 2).
Infants 2 Weeks to 6 Months of Age	Any weight loss. ***Use two separate weight measurements taken at least eight weeks apart (3).***

Required Documentation

DOB
Weight
Height
CDC WHO growth chart

Justification

Growth faltering is defined as a growth rate below that which is appropriate for an infant's age and sex. It can effect length, weight, and head circumferences resulting in values lower than expected. Growth faltering may include weight faltering (a drop in weight-for-age) or slowed growth where both weight and length growth are slower than expected. An example of weight faltering is a drop in weight after a minor illness or a measurement/plotting error (4).

Growth in infants is steady and predictable. It is a reflection of health and nutritional status and the overwhelming majority of infants have no growth problems (5, 6). Normal growth is also pulsatile, with periods of rapid growth or growth spurts followed by periods of slower or no measurable growth (5-8). Catch-up and catch-down growth during early childhood are normal phenomena that affect large numbers of children, particularly during infancy, and may merely be an adjustment to the genetic potential for growth (9). Growth is also seasonal, with length velocities (the change in growth over time) increased during the spring and summer months and stagnant other months (10). Weight may vary depending on the time of day and infant feeding schedule. Growth may be increased or slowed by a variety of conditions, with changes in growth as the first sign of a pathological condition. Such conditions include: under nutrition, hypothyroidism, iron deficiency, human immunodeficiency virus (HIV), inborn errors of metabolism, lead toxicity, zinc deficiency, immune deficiency, failure of a major organ system such as the gastrointestinal digestive system, renal, cardiovascular, and pulmonary (11). Infants that do not follow a steady predictable pattern, such as those with short stature or decreased growth rate, should be the focus of concern (11).

The timely detection of poor growth in early life is a way to identify infants who may be at risk for growth faltering, and intervene before under-nutrition has detrimental health outcomes, such as growth retardation, when incurred early are irreversible (12). It can help prevent short stature and adverse functional and deleterious long term consequences, such as poor cognition and educational performance, low adult wages, lost productivity, and when accompanied by weight gain later in childhood, an increased risk of nutrition-related chronic diseases (13, 14).

Excessive Weight Loss After Birth

Infant weight loss in the early postpartum period is physiologically normal, and nearly universal but the amount of weight loss varies (15). Weight loss of 5% and 7% of birth weight is not unusual for formula-fed or breastfed infants, respectively (16). Healthy infants are expected to regain their birth weight within 8-10 days after birth (17). However, if a breastfed infant loses 7% of birth weight in the first 72 hours after birth, an evaluation and review of the mother-infant dyad is needed and any problems resolved immediately. Risk of dehydration and failure to thrive in breastfed newborns can be mitigated by early screening and providing lactation support in the early postpartum period (18).

A weight loss of up to 10% or birth weight is the maximum acceptable weight loss for newborn infants, with any additional loss a potential emergency (17, 19). Contributing factors include (2, 16, 17, 20):

- Hospital practices like epidurals, pacifier use, low or non-nutritive feedings, or strict-feeding schedules.
- Maternal factors such as retained placenta, parity, anxiety, and poor maternal knowledge.
- Infant factors such as birth weight, gestational age, gender, and feeding method.
- Breastfed infants with poor positioning, latch and/or milk transfer.

WIC staff should identify and address any potential underlying feeding issues causing newborn weight loss (21). An infant with a weight loss of greater or equal to seven percent signals the need for careful evaluation and intervention, infants with a weight loss of ten percent or more is a marker for a medical referral (22).

Any Weight Loss 2 Weeks to 6 Months

While the 2006 CDC/WHO growth charts show slower growth from 3 – 18 months of age as a normal growth pattern, weight loss is not expected beyond the first two weeks of life and requires follow-up (23). After birth, growth faltering is caused by inadequate caloric intake, normal caloric intake in an environment of excessive loss or malabsorption; or increased metabolic needs. In cases of dehydration or acute illnesses like gastroenteritis, fluid loss that exceeds fluid intake may also lead to significant weight loss. Weight loss in young infants is commonly caused by acute infections, feeding problems, allergy to milk protein, lead poisoning, HIV, malnutrition, pyloric stenosis, gastrointestinal reflux, celiac disease, cystic fibrosis, neglect, growth failure, congenital heart disease, and inborn errors of metabolism.

The primary goal of the intervention is to enhance infant health outcomes by addressing causes of slowed growth and keeping vulnerable infants tracking along growth percentiles established in infancy. In some cases, it may be

important to intervene quickly, while in other cases a period of frequent growth monitoring would be more appropriate to prevent too rapid refeeding and subsequent increased risk of type 2 diabetes, obesity, and cardiovascular disease later in life (24, 25).

If faltering growth is suspected, maternal neglect and inadequate caloric intake due to inappropriate formula mixing, breastfeeding problems, early introduction of solid food, maternal depression, and emotional deprivation, must be ruled out and addressed (6). Growth monitoring should occur on a monthly basis – utilizing two separate weight measurements taken at least eight weeks apart as data markers. It is imperative that WIC staff involved in measuring infant growth use standardized equipment and receive adequate training prior to conducting infant measurements to increase the reliability between measures (26). If the participant does not respond to nutritional management (i.e. weight continues to falter) or if other markers falter (such as length for age), then the infant should be referred to their health care provider for assessment.

Normal Growth Patterns

Understanding normal growth patterns in infants is important. The pattern of weight gain during infancy varies depending on the method of feeding. Compared to formula-fed infants, breastfed infants gain weight rapidly in the first three to four months of life and relatively slowly thereafter. Although the weights of formula-fed breastfed infants are similar by one to two years of age, the typical pattern of slowed weight gain after three to four months among breastfed infants may lead to unnecessary early introduction of solid foods or cessation of breastfeeding if the slowed weight gain is perceived as lactational inadequacy. (27, 28, 29)

The table below shows the average mean values for weight gain for healthy exclusively breastfed infants:

Average of Mean Values for Gains in Weight for Healthy Exclusively Breastfed Infants (30)

Interval (mo)	Girls (g/day)	Boys (g/day)
0-1	30	33
1-2	28	34
2-3	22	23
3-4	19	20
4-5	15	16
5-6	13	14
6-7	12	11

Screening for Slow of Faltering Growth Patterns

Screening for slow or faltering growth patterns is a preventive health measure

which requires careful growth monitoring and critical thinking skills. And while a single measure of weight-for-age may be cause for concern, it cannot be interpreted to show growth faltering. No single measurement on its own is adequate for identifying nutritional growth delay (31). As stated earlier, it is imperative that WIC staff involved in measuring infant growth use standardized equipment and receive adequate training prior to conducting infant measurements to increase reliability between measures (26).

Growth faltering is a reflection of two weight measures, preferable eight weeks (two months) apart, to calculate an increment in growth. It is possible to use four week (one month) intervals for the assessment of slow growth patterns, but since there may be errors in clinical measurement, it is more prudent to use eight weeks as the minimum time interval between measurements. Infant weight will fluctuate over the course of the day and length growth may occur in discrete eriods lasting no more than 24 hours separated by growth-free intervals lasting as long as two months. Thus, growth that seems abnormal may be nothing more than a growth-free period in a child's life (10).

Screening for early growth failure should be done using multiple growth indicators, including risks for underweight (Risk #103), short stature (Risk #121) and failure to thrive (Risk #134) to allow for timely remedial interventions and prevention of further growth failure.

In summary, a three-step approach should be considered for evaluation of infants with suspected abnormal growth. First, growth data should be assessed for accuracy. Second, feeding problems, improper formula preparation, etc. should be assessed to determine if calorie intake is insufficient for growth and development. Third, the infant should be assessed for other medical conditions or developmental delay.

Implications for WIC Services

In most situations, growth may not simply be a factor of under nutrition, but rather a combination of environmental and other factors which will require a broad intervention strategy for successful health outcomes (32). In general, intervention strategies may include screening for environmental health factors such as (25, 32):

- Adequate nutrition and nutrient dense foods, including a history of breast milk or formula feeding.
- Appropriate introduction of complementary foods.
- Maternal conditions that impact lactation performance: mastitis, prolonged labor, C-Section, hypo or hyperthyroidism, Diabetes, low birth weight infant, pregravida BMI >27, pregnancy-induced hypertension, flat/inverted nipples, vitamin B12 deficiency.
- Meal time routine and eating/feeding behavior.
- Growth faltering in light of familial growth patterns.
- Neglect.
- Lack of social support.
- Adverse social and psychological environment.
- Depressed or poor mental abilities of parent/caregiver. It may manifest as dressing inappropriately for the weather; looking disheveled and lacking in hygiene; or making inappropriate faces or reactions like laughing.
- Lack of parental education and nutrition knowledge.

Nutrition counseling for this risk would ideally be provided by staff with specialized education and training to assess growth parameters and identify causative factors accurately. Intervention strategies to address this criterion include:

- Appropriate timing and type of participant intervention.
- Effective participant-centered nutrition counseling.
- Early postpartum breastfeeding support to minimize risk of dehydration and/or failure to thrive.
- Review of baby behavior hunger and satiety cues. (For more information see WIC Baby Behavior Basics, WIC Online Learning Module available on the WIC Works Resource System:
- https://wicworks.fns.usda.gov/wic-learning-online .)
- Review/adjustment to breastfeeding/formula feeding schedule.
- Review adjustment of formula mixing technique.
- Referral to lactation specialist for latch and position assistance.
- Tailored food package prescription.
- Review accuracy of weight and length measurements.

Referral to allied health professionals such as: physician, early childhood intervention, social services, and home visiting program.

A variety of intervention strategies can help infants establish and maintain individual growth patterns. The desired outcome is one where the infant's own growth curve tracks within the channel established in early infancy. Also, because growth monitoring is an intervention that happens largely after the fact, there may be benefit to anticipatory guidance that provides prevention rather than crisis management of this problem (33). It is suggested that when feeding is going well, the baby will eat as much as she needs and grow in the way that is right for her if parents maintain a division of responsibility in feeding (34).

References

1. American Academy of Pediatrics. Policy Statement. Breastfeeding and the use of human milk. Ped. 2005; 115(2):496-506.
2. Academy of Breastfeeding Medicine. Clinical Protocal #3. Hospital guidelines for the use for supplementary feedings in the healthy term breastfed neonate. 2009;(4):175-82.
3. Grummer-Strawn L, Reinhold C, Krebs N. Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United Staes. CDC Recommendations and Reports September 10, 2010;59(rr09):1-15.
4. Wacogne, S. Weight faltering and failure to thrive in infancy and childhood. BMJ, 345:e5931.
5. Beker J. Principles of growth assessment. Pediatrics in Review 2006:27:196-198.
6. Kessler D. Growth assessment and growth failure: Overview and the role of nutrition. Pediatric Perspectives 2007;4(1); 1-4.
7. Lampl, M, Velhius JD, Johnson ML. Saltation and stasis: A model of human growth. Science 1992: 258:801-803.
8. Lamp, M, Johnson ML Emmett, PM, et al. Mixed distribution analysis identifies saltation and stasis growth. Ann Hum Biol 2001;28:403.
9. Mei, Z. Shifts in percentiles of growth during early childhood: analysis of longitudinal data from the California Child Health and Development Study. Pediatrics 2004;133(6):617-27.

10. Gelander L, Karlberg J, Albertsson-Wikland K. Seasonality in lower leg length velocity in prepubertal children. Acta Paediatr 1994;83:1249.
11. Legler J, Rose L. Assessment of abnormal growth curves. Am Fam Phys 1998: (1)153-58.
12. Cameron N, Preece MA. Catch-up growth or regression to the mean? Recovery from stunting revisited. Am J Hum Biol 2005;17:412-7.
13. Victoria CG, Adair L, Fall C, et al.Maternal and child undernutrition: consequences for adult health and human capital. Lancet. 2008:371:340-357.
14. De Onis, Mercedes, et al. Worldwide implementation of the WHO Child Organization growth standards. Public Health Nutrition 2012;1-8.
15. Flaherman VJ, Schafer EW, Kuzniewicz MW, Li SX, Walsh EM, Paul IM. Key weight loss nomograms for exclusively breastfeed newborns. Ped. 2015;135(1)e16-23.
16. Martens PJ, Romph, L. Factors associated with newborn in-hospital weight loss: comparisons by feeding method, demographics, and birthing procedures. J Hum Lact 2007;23:233.
17. Wright CM, Parkinson KN. Postnatal weight loss in term infants: what is normal and do growth charts allow for it? Arch Dis Child Fetal Neonatal Ed. 2004;89:f254-257.
18. Gahagan, S. Failure to thrive: a consequence of undernutrition. Pediatrics in Review 2006;27:1-11.
19. Kirkland R, Motil K. Failure to thrive (undernutrition) in children younger than two years. Up To Date 2014 (August).
20. Reilev M, Borch K, Pryds OA. Neonatal hypernatramic dehydration – why increasing incidence? Ugeskr Laiger. 2007:169:1227-31.
21. Livingston, V. Failure to thrive while breastfeeding. Breastfeeding Medicine Vol 1, No. 2, 2006.
22. Lawrence RA, Lawrence RM. Breastfeeding: a guide for the medical profession. 6th Edition, St. Louis, MO: Elseveier Mosby. 2005.
23. Grummer-Strawn L. Response letter to Patricia Daniels 2008 risk criterion #135 questions. Centers for Disease Control and Prevention. October 11, 2008.
24. Eriksson JG, et al. Catch-up growth in childhood and death from coronary heart disease: longitudinal study. BMJ 1999; 318:427-31. Later 2001 reference entitled, early growth and coronary heart disease in later life: longitudinal study by JG Ericsson et al, BMJ, Apr 21, 2001; 322(7292);949-953.
25. Hales CN; Ozanne SE. The dangerous road of catch-up growth. J Physiol 2003;547:5-10.
26. World Health Organization. Physical status: The use and interpretation of anthropometry. Report of a WHO Expert Committee. WHO Technical Report Series no. 854. Geneva: WHO (1995).
27. Mei Z, et al. Comparison of prevalence of shortness, underweight, and overweight among US children aged 0-59 months by using the CDC 2000 and the WHO 2006 growth charts. J Pediatr. 2008:153:622.
28. Van Dijk, et al. Growth-curve standards and the assessment of early excess weight gain in infancy. Pediatrics 2009; 123:102.
29. Whitehead RG. The importance of diet-specific growth charts. Acta Paediatr 2003; 92:137.
30. The American College of Obstetricians and Gynocologists, American Academy of Pediatrics. Breastfeeding Handbook for Physicians. 2nd e. 2014. ACOG, Washington DC; AAP Elk Grove, IL.
31. Kessler D, et al. Failure to thrive: the prevalence and concurrence of anthropometric criteria in a general infant population. Archives fo Disease in Childhood. 2007 Feb; 92(2):109-114.

32. Barker DJ. The developmental origins of adult disease. Eur J Epidemiol 2003;18:733-6.
33. Shrimpton R, et al. Worldwide timing of growth faltering: implications for nutritional interventions. Pediatrics 2001:107:e75.
34. American Academy of Pediatrics. Bright Futures Nutrition. 3rd ed. 2011.

Nutrition Risk Criteria **141★ LOW BIRTH WEIGHT**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD
CHILDREN	3	RD

Definition

Low Birth Weight (LBW)
Birth weight defined as less than or equal to (≤) 5 lbs 8 oz (≤ 2500g) for infants and children less than 24 months.

Required Documentation

DOB
Birth weight

Justification

Low Birth Weight (LBW) is one of the most important biologic predictors of infant death and deficiencies in physical and mental development during childhood among those babies who survive and continues to be a strong predictor of growth in early childhood. Infants and children born with LBW, particularly if caused by fetal growth restriction, need an optimal nutrient intake to survive, meet the needs of an extended period of relatively rapid postnatal growth, and complete their growth and development.

References

1. Groh-Wargo S, Thompson M, Cox JH, Hartline JV. Nutritional care for high-risk newborns. editors Sharon Groh-Wargo, Melody Thompson, Janice Hovasi Cox; consulting editor John V. Hartline. Chicago: Precept Press, Inc, 1994.
2. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Nutrition Risk Criteria **142★ PRETERM OR EARLY TERM DELIVERY**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD
CHILDREN (less than 24 months of age)	3	RD

Definition

Preterm and early term delivery are defined as follows (1, 2):
Preterm: Delivery of an infant born less than or equal to (≤)36 6/7 weeks gestation.
Early term: Delivery of an infant born greater than or equal to (≥)37 0/7 weeks gestation and less than or equal to (≤)38 6/7 weeks gestation.

Note: See Clarification section for information on plotting growth measurements for preterm infants.

Required Documentation

EDD
DOB

Justification

Preterm birth is a significant cause of newborn morbidity and mortality. Preterm and early term deliveries strain society's healthcare resources due to the longer hospital stays for the infant and the long-term effects on the health of the newborn (3, 4).

Typically, a pregnancy lasts about 40 weeks. Premature or preterm birth, however, is defined as a birth that occurs between 20 and 37 weeks of pregnancy, according to the American College of Obstetricians and Gynecologists (ACOG) (5). In the past, the period from 3 weeks before until 2 weeks after the estimated date of delivery was considered a "term" pregnancy, with the expectation that a baby would have similar health outcomes if they were born any time during this interval. In 2013, ACOG released a committee opinion that the label "term" should be replace with the designations *early term* (≥ 37 0/7 weeks and ≤38 6/7 weeks gestation) and *full term* (≥39 0/7 weeks and ≤40 6/7 weeks gestation) to more accurately describe these groups of infants (1).

Preterm Delivery

Prematurity affects about 12% of all live births in the U.S., and about 50% of these preterm births were preceded by preterm labor (6). In 2011, the annual rate of premature births in the United States reached 11.7%, nearly two times the rate in European nations (6). Preterm births also account for approximately 70% of newborn deaths and 36% of infant deaths (5).

Several factors have been found to increase the risk of preterm delivery. Epidemiological studies have consistently reported low socioeconomic status, nonwhite race, maternal age of ≤ 18 years or ≥ 40 years, and low pre-pregnancy weight as risk factors. A history of one previous preterm birth is associated with recurrent risk of 17-37%; the risk increases with the number of prior preterm births and decreases with the number of term deliveries. Other maternal factors associated with a risk of preterm birth may include low weight gain during pregnancy, maternal obesity, hypertension, diabetes, or sexually transmitted

diseases (7). (See risk 311 *History of Preterm or Early Term Delivery* for more details.)

Despite advances in neonatal care, preterm birth remains a leading cause of infant death in the United States (8). Preterm infants may have health problems because their organs did not have enough time to develop in the womb. Babies that are born too early may have a number of health conditions, including:

- Low or very low birth weight (9)
- Increased caloric needs (9)
- Feeding difficulties due to a lack of reflexes for sucking and swallowing (9)
- Immature digestion and impaired absorption of carbohydrates and lipids (10, 11)
- Breathing problems like chronic lung disease/bronchopulmonary dysplasia and apnea (9, 12, 13)
- Cerebral palsy, an impairment of the brain that controls movement and muscle tone (10, 14)
- Developmental delay and poorer cognitive function (12, 15, 16, 17)
- Vision problems like retinopathy of prematurity (ROP), which may cause blindness (12, 15)
- Hearing problems
- Behavioral problems and psychiatric disorders (16, 17)
- Increased risk for necrotizing enterocolitis (NEC) due to their immature gastrointestinal systems (10, 12)
- Increased risk for Sudden Infant Death Syndrome (SIDS) (10)
- Temperature control problems (9, 10)
- Heart problems like patent ductus arteriosus and low blood pressure (hypotension) (10, 12)
- Blood problems like anemia and jaundice (10, 13)
- Hypoglycemia (9, 10)
- Immature immune systems, which may result in infections (9)

Preterm infants often need special medical care in a neonatal intensive care unit (NICU) and may need to stay there for days or even months. Breastfeeding is recommended as the normative standard for infant feeding and nutrition for all infants, especially preterm babies. Breastfeeding preterm infants has been associated with positive health outcomes for these infants, including:

- Improved motor maturity and cognitive ability (18, 19, 20)
- Reduced risk of NEC (21, 22)
- Reduced risk of ROP and retinal detachment (23)

Additionally, mothers of preterm infants produce milk that is designed to meet the baby's particular nutritional needs during the first few weeks of life. It is higher in protein and minerals, such as salt, and contains different types of fat that are easier to digest and absorb compared to fats in the milk of mothers of full term babies. The fat in human milk also helps to enhance the development of the baby's brain and neurologic tissues, which is especially important for premature infants. Human milk is also easier for babies to digest than infant formula and avoids exposing the baby's immature intestinal lining to the cow's milk proteins found in premature infant formula. Preterm infants who are breastfed are less likely to develop intestinal infections than babies who are formula fed, and the colostrum produced in the first few days contains high concentrations of antibodies that will help the baby fight infection (22).

Breastfeeding preterm infants, especially if they are in the NICU, may present unique challenges for breastfeeding dyads. These mothers will benefit from extra breastfeeding support due to the delay of direct breastfeeding, reliance on breast pumps, and the stress of having a sick newborn. Even if the baby cannot breastfeed directly from the breast at first, the mother can be encouraged to express her milk to ensure that her supply is maintained. Supportive care for infants in the NICU may include the use of a feeding tube. Expressed human milk can be passed through the tube, therefore, it is important for the mother to discuss her feeding decisions with her baby's doctor. Preterm infants sometimes need additional calories and nutrients to facilitate adequate growth and in such cases a human milk fortifies may be prescribed by a health care provider (22).

Preterm infants who are not breastfed may require the use of a formula higher in calories and nutrients to support their growth. According to the American Academy of Pediatrics (AAP), soy formulas are typically not recommended for low birth weight preterm infants, as their use may result in less weight gain and lower serum albumin and phosphorus levels than cow's milk-based formulas (24).

In addition to breastfeeding, skin-to-skin care or kangaroo care (holding your baby naked or in just a diaper on your bare chest), can help preterm infants breathe better, gain weight, keep their body at the right temperature, and prepare them for breastfeeding (25). All caregivers can provide skin-to-skin care, not just the mother.

Infants born at 34 0/7 through 36 6/7 weeks gestation, called late preterm infants, are sometimes mistaken for term infants since their actual size and weight may be similar (10). However, caregiver, healthcare providers, nutritionists, and lactation consultants must be aware that these babies are physiologically and metabolically immature (9). In addition to the health conditions previously mentioned for preterm infants, it is important to be aware that lare preterm babies have an increased risk of morbidity and mortality which is often related to feeding problems. Due to their immaturity, late preterm infants may have more challenges with breastfeeding because they tire easily and have less stamina, which results in greater difficulty with latching, sucking, and swallowing. Mothers of late preterm infants will benefit greatly from timely lactation assessment and support since feeding difficulties, slow weight gain, failure to thrive, hypoglycemia, and jaundice are very common in these babies (26).

Preterm infants have different patterns of growth compared to term infants. Plotting the growth of preterm infants using their adjusted gestational age is an essential component of care until they reach 24 to 36 months of age (27). (See the *Clarification* section for more information on how to determine adjusted gestational age.) Most preterm infants, however, show catch-up growth in weight, length, and head circumference after their initial postnatal growth failure. If catch-up growth occurs, it usually starts early in the first months of life and is often achieved within the first years of life (28).

The effects of preterm birth can continue beyond infancy. Children who were born prematurely are at an increased risk for the following:

- Neurodevelopmental problems (29)
- Intellectual/cognitive impairments, which can lead to learning disabilities and the need for special education services (29, 30, 31)
- Motor problems (31)

- Feeding difficulties such as problems with chewing and swallowing, late development of feeding skills, food refusal, eating behavior problems, and poor appetite (32)
- Emotional problems such as anxiety and depression (31)
- Behavioral concerns such as attention problems and hyperactivity (31)

Early Term Delivery

Up to 10% of babies in the United States are scheduled for early term deliveries via labor-inducing medication or cesarean section before 39 weeks of gestation despite neither the mother nor the baby being at risk if the pregnancy continues (4). Elective deliveries like this are sometimes requested for reasons such as wanting to schedule the date of the infant's birth, physician preference, or for relief of symptoms at the end of pregnancy (4).

Research shows that a fetus will experience a significant amount of development and growth of the lungs, brain, and liver between 37 and 39 weeks of gestation. The brain develops at its fastest rate at the end of the pregnancy, at a rate of up to one third between weeks 35 and 39. Additionally, layers of fat are added under the infant's skin during the last few weeks of pregnancy which helps them keep warm after birth. According to ACOG, non-medically warranted deliveries prior to 39 weeks should be avoided (33). Early term delivery puts an additional strain on society as the early term infant will likely require a longer hospital stay and may have long term healthcare needs (4).

Implications for WIC Nutrition Services

WIC services can directly support preterm and early term infants and their caregivers, as these babies may have unique feeding difficulties. Preterm delivery is often unexpected and a mother may not have made decisions about how to feed her baby yet. These infants may require additional calories, extra breastfeeding support, and/or the use of a human milk fortifier or special infant formula.

WIC can support preterm and early term infants and their caregivers through:

- Promoting and supporting breastfeeding as the normative standard for infant nutrition and providing early and frequent breastfeeding support.
- Recommending the use of a hospital grade electric breast pump for expressing milk if the baby is in the NICU or the baby is unable to breastfeed directly from the breast.
- Providing anticipatory guidance about potential feeding challenges.
- Encouraging caregivers to provide skin-to-skin contact.
- Providing education on safe preparation, handling, and storage of breast milk and/or formula.
- Educating pregnant women about the importance of carrying a baby to term, unless medically contraindicated.
- Monitoring the child's growth to ensure healthy weight gain.
- Providing nutrition education for mothers/caregivers and appropriate referrals as necessary for growth, feeding, health, and/or infant developmental issues.

Clarification

All preterm infants and children (up to 2 years of age) who have reached the equivalent age of 40 weeks gestation, shall be assessed for growth using the

Centers for Disease Control and Prevention (CDC) Birth to 24 Months gender specific growth charts adjusting for gestational age as follows:

1. Document the infant/child's gestational age (at delivery) in weeks. (Mother/caregiver can self-report, or referral information from the medical provider may be used.)
2. Subtract the child's gestational age in weeks from 40 weeks (gestational age of term infant) to determine the adjustment for prematurity in weeks.
3. Subtract the adjustment for prematurity in weeks from the child's chronological postnatal age in weeks to determine the child's gestation-adjusted age.

Example:

Randy was born prematurely on March 19. His gestational age at birth was determined to be 30 weeks based on ultrasonographic examination. At the time of the June 11 clinic visit, his chronological postnatal age is 12 weeks. What is his gestation-adjusted age?

- 30 = gestational age in weeks
- 40 – 30 = 10 weeks adjustment for prematurity
- 12 – 10 = 2 weeks gestation-adjusted age

His measurements would be plotted on a growth chart as a 2-week-old infant.

Note: Preterm infants (less than or equal to ≤ 36 6/7 weeks gestation) who have not reached the equivalent age of 40 weeks gestation may be assessed for growth using a growth chart for low birth weight (LBW) or very low birth weight (VLBW) infants (e.g., Infant Health and Development Program [IHDP]) consistent with the protocols of the local medical community in which the WIC clinic operates. The CDC does not recommend the use of the CDC Growth Charts for preterm infants who have not reached the equivalent age of 40 weeks gestation.

Reference

1. American College of Obstetricians and Gynecologists. Definition of term pregnancy. Committee Opinion No.579. Obstet Gynecol. 2013 Nov;122:1139-40
2. Ob-Gyns redefine meaning of "term pregnancy" [Internet]. Washington, DC: American College of Obstetricians and Gynecologists; c2013 [updated 2013 Oct 22; cited 2016 Dec 6]. Available from: http://www.acog.org/About-ACOG/News-Room/News-Releases/2013/Ob-Gyns-Redefine-Meaning-of-Term-Pregnancy .
3. Wang P, Liou S, Cheng C. Prediction of maternal quality of life on preterm birth and low birthweight: a longitudinal study. BMC Pregnancy Childbirth. 2013 June 2;13:124.
4. National Institute for Health Care Management. Born too early-improving maternal and child health by reducing early elective deliveries. NIHCM Issue Brief, March 2014. [cited 2016 Dec 6]. Available from: http://www.nihcm.org/pdf/Early_Elective_Delivery_Prevention_Brief_2014.pdf .
5. ACOG.org [Internet]. Washington, DC: The American College of Obstetricians and Gynecologists; c2016 [updated 2016 Nov; cited 2016 Dec 6]. Available from: http://www.acog.org/Patients/FAQs/Preterm-Premature-Labor-and-Birth .

6. Dag M, Lie TR, Markestad T. Long-term medical and social consequences of preterm birth. N Engl J Med. 2008 Jul 17. Web. 07 Apr. 2014.
7. Hoffman HJ, Bakketeig LS. Risk factors associated with the occurrence of preterm birth. Clin Obstet Gynecol. 1984 Sep; 27:539-52.
8. Iams JD. Prevention of preterm parturition. N Engl J of Med. 2014 Jan;370(3):254-261.
9. Engle WA, Tomashek KM, Wallman C. “Late-preterm” infants: a population at risk. Pediatrics. 2007 Dec;120(6):1390-1401.
10. Mayoclinic.org [Internet]. Rochester: Mayo Foundation for Medical Education and Research; c1998-2016 [updated 2014 Nov 27; cited 2016 Dec 1]. Available from: http://www.mayoclinic.org/diseases-conditions/premature-birth/basics/definition/con-20020050 .
11. Barlow SM, Finan DS, Lee J, Chu S. Synthetic orocutaneous stimulation entrains preterm infants with feeding difficulties to suck. Journal of Perinatology. 2008 June 12;28:541-548.
12. Ward RM, Beachy JC. Neonatal complications following preterm birth. BJOG. 2003 April;110(s20):8-16.
13. Baraldi E, Filippone M. Chronic lung disease after premature birth. N Engl J Med. 2007 Nov 8;357:1946-55.
14. Vincer MJ, Allen AC, Joseph KS, Stinson DA, Scott H, Wood E. Increasing prevalence of cerebral palsy among very preterm infants: a population-based study. Pediatrics. 2006 Dec;18(6)e1621-26.
15. American Academy of Pediatrics Policy Statement. Screening examination of premature infants for retinopathy of prematurity. Pediatrics. 2013 Jan;131(1):189-95.
16. Vohr B. Long-term outcomes of moderately preterm, late preterm, and early term infants. Clin. Perinatol. 2013 Dec;40(4):739-51.
17. De Jong M, Verhoeven M, van Baar AL. School outcome, cognitive functioning, and behavior problems in moderate and late preterm children and adults: a review. Semin Fetal Neonatal Med. 2012 Jun;17(3):163-9.
18. Feldman R, Eidelman A. Direct and indirect effects of breast milk on the neurobehavioral and cognitive development of premature infants. Developmental Psychobiology. 2003 Sept;43(2):109-19.
19. Vohr B, Poindexter B, Dusick A, McKinley LT, Higgins RD, Langer JC, Poole KW. Persistent beneficial effects of breast milk ingested in the neonatal intensive care unit on outcomes of extremely low birth weight infants at 30 months of age. Pediatrics. 2007 Oct;120(4):e953-59.
20. Blaymore Bier J, Oliver T, Ferguson AE, Vohr B. Human Milk Improves Cognitive and Motor Development of premature infants during infancy. J Hum Lact. 2002 Nov;18(4)361-67.
21. Boyd CA, Quigley MA, Brocklehurst P. Donor breast milk versus infant formula for preterm infants: systematic review and meta-analysis. Arch Dis Child Fetal Neonatal Ed. 2007 May;92:F169-75.
22. Healthychildren.org [Internet]. Elk Grove Village: American Academy of Pediatrics; c2011 [updated 2015 Nov 21; cited 2016 Dec 6]. Available from: https://www.healthychildren.org/English/ages-stages/baby/breastfeeding/Pages/Providing-Breastmilk-for-Premature-and-Ill-Newborns.aspx .
23. Okamoto T, Shirai M, Kokubo M, Takahashi S, Kajino M, Takase M, Sakata H, Oki J. Human milk reduces the risk of retinal detachment in extremely low-birthweight infants. Pediatr Int. 2007 Oct;49(6):894-897.

24. American Academy of Pediatrics Committee on Nutrition. Soy protein-based formulas: recommendations for use in infant feeding. Pediatrics. 1998 Jan;101(1):148-53.
25. Kuhn KS, Kuhn MJ. Kangaroo care for your premature or sick baby. J Hum Lact. 2011 Feb;27(1):66-67.
26. Academy of Breastfeeding Medicine. ABM Clinical Protocol #10: Breastfeeding the late term infant (34 0/7 to 36 6/7 weeks gestation). Breastfeeding Medicine. 2011 June;6(3):151-56.
27. Canadian Paediatric Society. A health professional's guide for using the new WHO growth charts. Paediatr Child Health. 2010 Feb;15(2): 84-90.
28. De Wit CC, Sas TCJ, Wit JM, Cutfield WS. Patterns of catch-up growth. The Journal of Pediatrics. 2013 Feb;162(2):415-20.
29. Johnson S, Strauss V, Gilmore C, Jaekel J, Marlow N, Wolke D. Learning disabilities among extremely preterm children without neurosensory impairment: Comorbidity, neuropsychological profiles and scholastic outcomes. Early Hum Dev. 2016 Aug 9;103:69-75.
30. Breeman LD, Jaekel J, Baumann N, Bartmann P, Wolke D. Preterm cognitive function into adulthood. Pediatrics. 2015 Sept;136(3):415-23.
31. Hornman J, de Winter AF, Kerstjens JM, Bos AF, Reijneveld SA. Emotional and behavioral problems of preterm and fully-term children at school entry. Pediatrics. 2016 May;137(5).
32. Johnson S, Matthews R, Draper ES, Field DJ, Manktelow BN, Marlow N, Smith LK, Boyle EM. Eating difficulties in children born late and moderately preterm at 2 years of age: A prospective population-based cohort study. Am J Clin Nutr. 2016 Feb;103(2):406-14.
33. Elective delivery before 39 weeks [Internet]. Washington, DC: American College of Obstetricians and Gynecologists; c2013 [updated 2013 June; cited 2016 Dec 6]. Available from: http://www.acog.org/Patients/FAQs/Elective-Delivery-Before-39-Weeks .

Nutrition Risk Criteria **153★ LARGE FOR GESTATIONAL AGE**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	INFANTS	1	-

Definition

Large for Gestational Age
Birth weight greater than or equal to (≥) 9 lbs (4000g);

OR

Presence of large for gestational age diagnosed by a physician or a self-reported diagnosis by a physician.

Required Documentation

DOB
Birth weight
Documentation of source of information (i.e. self-report of diagnosis by a physician or written documentation by physician).

Justification

Infant mortality rates are higher among full-term infants who weigh > 4,000 g (> 9 lbs) than for infants weighing between 3,000 and 4,000 g (6.6 and 8.8 lbs). Oversized infants are usually born at term; however, preterm infants with weights high for gestational age also have significantly higher mortality rates than infants with comparable weights born at term. Large for Gestational Age may be a result of maternal diabetes (which may or may not have been diagnosed before or during pregnancy) and may result in obesity in childhood that may extend into adult life.

Very large infants regardless of their gestational age, have a higher incidence of birth injuries and congenital anomalies (especially congenital heart disease) and developmental and intellectual retardation. When large for gestational occurs with pre-term birth, the mortality risk is higher than when either condition exists alone.

Weight should be plotted weekly on an intrauterine or postnatal growth chart and monitored for progress. Intrauterine growth charts have been developed by compiling birth weight, length, and head circumference of infants at varying gestational ages and reflect intrauterine growth, while postnatal growth charts were developed from extrauterine growth data. Both types of growth charts have limitations that should be considered. Choice and use of growth charts are typically facility-specific and determined by the medical team. The Oregon Intrauterine Growth Chart (Babson/Benda), Denver Intrauterine Growth Chart (Lubchenco), and the Hall/Shaffer and Wright Postnatal Growth Charts are generally available to monitor growth by gestational age. Copies of the Babson/Benda and Lubchenco charts are provided in the Reference Material Section.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Babson SG, Benda GI. Growth graphs for the clinical assessment of infants of varying gestational age. J.Pediatr. 1976; 89:814-20.
2. Behrman RE, Kliegman R, Jenson HB. Nelson textbook of pediatrics. Philadelphia: Saunders, 2000.
3. Lubchenco LO, Hansman C, Boyd E. Intrauterine growth in length and head circumference as estimated from live births at gestational ages from 26 to 42 weeks. Pediatrics 1966; 37:403-8.

Nutrition Risk Criteria **201★ LOW HEMATOCRIT/LOW HEMOGLOBIN**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD if Hct ≤ 31.0, Hb ≤ 10.0, or no rise on recheck
BREASTFEEDING WOMEN	1	
NON-BREASTFEEDING WOMEN	6	
INFANTS	1	
CHILDREN	3	

Definition

Hemoglobin or hematocrit concentration below 95 percent confidence interval (i.e., below the .025 percentile) for healthy, well-nourished individuals of the same age, sex, and stage of pregnancy.

Cut-off values are provided in the attached Tables for each clinic site (with adjustments for smoking and altitude), based on the levels established by the Centers for Disease Control and Prevention (CDC).

Required Documentation

ALL: Hematocrit or Hemoglobin value and date, clinic altitude (tables)
INFANTS/CHILDREN: DOB
PREGNANT: Weeks gestation
WOMEN: Smoking status

Justification

Hemoglobin (Hb) and hematocrit (Hct) are the most commonly used tests to screen for iron deficiency anemia. Measurements of Hb and Hct reflect the amount of functional iron in the body. Changes in Hb concentration and Hct occur at the late stages of iron deficiency. While neither an Hb or Hct test are direct measures of iron status and do not distinguish among different types of anemia, these tests are useful indicators of iron deficiency anemia.

Iron deficiency is by far the most common cause of anemia in children and women of childbearing age. It may be caused by a diet low in iron, insufficient assimilation of iron from the diet, increased iron requirements due to growth or pregnancy, or blood loss. Anemia can impair energy metabolism, temperature regulation, immune function, and work performance. Anemia during pregnancy may increase the risk of prematurity, poor maternal weight gain, low birth weight, and infant mortality. In infants and children, even mild anemia may delay mental and motor development. The risk increases with the duration and severity of anemia, and early damages are unlikely to be reversed through later therapy.

Clarification

1. **Basis for bloodwork assessment:** For pregnant women being assessed for iron deficiency anemia, bloodwork must be evaluated using trimester values established by CDC. Thus, the blood test result for a pregnant woman would be assessed based on the trimester in which her bloodwork was taken.

2. **Definition of Trimester:** CDC defines a trimester as a term of three months in the prenatal gestation period with the specific trimesters defined as follows in weeks:

First Trimester: 0-13 weeks
Second Trimester: 14-26 weeks
Third Trimester: 27-40 weeks.

3. **Adjustments for smoking:** A State agency may elect to use only one cutoff for all smokers rather than making specific adjustments based on the individual applicant's smoking frequency. If the State chooses to use only one category for this issue, the "up to < 1 pack/day" cutoff values category as shown on Tables 201-A and 201-B is the only one that may be used.

Further, CDC begins the calculation of weeks starting with the first day of the last menstrual period. If that date is not available, CDC estimates that date from the estimated date of confinement (EDC). This definition is used in interpreting CDC's Prenatal Nutrition Surveillance System data, comprised primarily of data on pregnant women participating in the WIC Program.

References

1. Centers for Disease Control and Prevention. Criteria for anemia in children and childbearing-aged women. MMWR 1998:47: RR-3.
2. Centers for Disease Control and Prevention. Prenatal Nutrition Surveillance System User's Manual. Atlanta: CDC, 1994.
3. Institute of Medicine. Iron deficiency anemia: recommended guidelines for the prevention, detection, and management among U.S. children and women of childbearing age. National Academy Press, Washington, D.C.; 1993.
4. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
5. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 1: PANHANDLE WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Athol Bonners Ferry Hayden Kellogg Priest River Sandpoint St. Maries	NON-SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Athol Bonners Ferry Hayden Kellogg Priest River Sandpoint St. Maries	NON-SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 2: NORTH CENTRAL WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Kamiah Lewiston Moscow Orofino	NON-SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---
3000-3999 ft. Grangeville	NON-SMOKER	11.2	10.7	11.2	12.0	11.2	11.3
	SMOKER	11.5	11.0	11.5	12.3	---	---

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Kamiah Lewiston Moscow Orofino	NON-SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---
3000-3999 ft. Grangeville	NON-SMOKER	33.5	32.5	33.5	36.2	33.4	33.5
	SMOKER	34.5	33.5	34.5	37.2	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 3: SOUTHWEST WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Canyon County Canyon Springs Council Emmett Farmway Village Grand View Homedale Payette Teen Parent Weiser Wilder	NON-SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 3: SOUTHWEST WIC PROGRAM

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Canyon County Canyon Springs Council Emmett Farmway Village Grand View Homedale Payette Teen Parent Weiser Wilder	NON-SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 4: CENTRAL WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Boise Glenns Ferry Horseshoe Bend	NON- SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---
3000-3999 ft. Garden Valley Idaho City Mountain Home	NON- SMOKER	11.2	10.7	11.2	12.0	11.2	11.3
	SMOKER	11.5	11.0	11.5	12.3	---	---
4000-4999 ft. Cascade	NON-SMOKER	11.3	10.8	11.3	12.1	11.3	11.4
	SMOKER	11.6	11.1	11.6	12.4	---	---
5000-5999 ft. McCall	NON-SMOKER	11.5	11.0	11.5	12.3	11.5	11.6
	SMOKER	11.8	11.3	11.8	12.6	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 4: CENTRAL WIC PROGRAM

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Boise Glenns Ferry Horseshoe Bend	NON- SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---
3000-3999 ft. Garden Valley Idaho City Mountain Home Mtn Home AFB	NON-SMOKER	33.5	32.5	33.5	36.2	33.4	33.5
	SMOKER	34.5	33.5	34.5	37.2	---	---
4000-4999 ft. Cascade	NON-SMOKER	34.0	33.0	34.0	36.7	33.9	34.0
	SMOKER	35.0	34.0	35.0	37.7	---	---
5000-5999 ft. McCall	NON-SMOKER	34.5	33.5	34.5	37.2	34.4	34.5
	SMOKER	35.5	34.5	35.5	38.2	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 5: SOUTH CENTRAL WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
3000-3999 ft. Gooding Jerome Shoshone Twin Falls	NON-SMOKER	11.2	10.7	11.2	12.0	11.2	11.3
	SMOKER	11.5	11.0	11.5	12.3	---	---
4000-4999 ft. Burley	NON-SMOKER	11.3	10.8	11.3	12.1	11.3	11.4
	SMOKER	11.6	11.1	11.6	12.4	---	---
5000-5999 ft. Bellevue Hailey	NON-SMOKER	11.5	11.0	11.5	12.3	11.5	11.6
	SMOKER	11.8	11.3	11.8	12.6	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 5: SOUTH CENTRAL WIC PROGRAM

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
3000-3999 ft. Gooding Jerome Shoshone Twin Falls	NON-SMOKER	33.5	32.5	33.5	36.2	33.4	33.5
	SMOKER	34.5	33.5	34.5	37.2	---	---
4000-4999 ft. Burley	NON-SMOKER	34.0	33.0	34.0	36.7	33.9	34.0
	SMOKER	35.0	34.0	35.0	37.7	---	---
5000-5999 ft. Bellevue Hailey	NON-SMOKER	34.5	33.5	34.5	37.2	34.4	34.5
	SMOKER	35.5	34.5	35.5	38.2	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 6: SOUTHEASTERN WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
4000-4999 ft. Aberdeen American Falls Blackfoot Fort Hall Malad Preston Pocatello Roosevelt Teen	NON-SMOKER	11.3	10.8	11.3	12.1	11.3	11.4
	SMOKER	11.6	11.1	11.6	12.4	---	---
5000-5999 ft. Arco Montpelier Soda Springs	NON-SMOKER	11.5	11.0	11.5	12.3	11.5	11.6
	SMOKER	11.8	11.3	11.8	12.6	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 6: SOUTHEASTERN WIC PROGRAM

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
4000-4999 ft. Aberdeen American Falls Blackfoot Fort Hall Malad Preston Pocatello Roosevelt Teen	NON-SMOKER	34.0	33.0	34.0	36.7	33.9	34.0
	SMOKER	35.0	34.0	35.0	37.7	---	---
5000-5999 ft. Arco Montpelier Soda Springs	NON-SMOKER	34.5	33.5	34.5	37.2	34.4	34.5
	SMOKER	35.5	34.5	35.5	38.2	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 7: EASTERN WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
4000-4999 ft. Idaho Falls Rexburg Rigby Salmon St. Anthony Terreton	NON-SMOKER	11.3	10.8	11.3	12.1	11.3	11.4
	SMOKER	11.6	11.1	11.6	12.4	---	---
5000-5999 ft. Challis Dubois	NON-SMOKER	11.5	11.0	11.5	12.3	11.5	11.6
	SMOKER	11.8	11.3	11.8	12.6	---	---
6000-6999 ft. Driggs	NON-SMOKER	11.7	11.2	11.7	12.5	11.7	11.8
	SMOKER	12.0	11.5	12.0	12.8	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

DISTRICT 7: EASTERN WIC PROGRAM

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
4000-4999 ft. Idaho Falls Rexburg Rigby Salmon St. Anthony Terreton	NON-SMOKER	34.0	33.0	34.0	36.7	33.9	34.0
	SMOKER	35.0	34.0	35.0	37.7	---	---
5000-5999 ft. Challis Dubois	NON-SMOKER	34.5	33.5	34.5	37.2	34.4	34.5
	SMOKER	35.5	34.5	35.5	38.2	---	---
6000-6999 ft. Driggs	NON-SMOKER	35.0	34.0	35.0	37.7	34.9	35.0
	SMOKER	36.0	35.0	36.0	38.7	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

AGENCY 8: NEZ PERCE WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Kamiah Lapwai	NON- SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Kamiah Lapwai	NON- SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---

TABLE 1: HEMOGLOBIN AND HEMATOCRIT BASELINE VALUES

AGENCY 11: BENEWAH WIC PROGRAM

HEMOGLOBIN

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Plummer	NON- SMOKER	11.0	10.5	11.0	11.8	11.0	11.1
	SMOKER	11.3	10.8	11.3	12.1	---	---

HEMATOCRIT

Values less than baseline values for healthy, well-nourished individuals are considered at risk. Adjustments are made for trimester of pregnancy, smoking status, and altitude. Baseline is the 95th confidence interval (0.025th percentile).

CLINIC SITES	SMOKING STATUS	PREGNANT WOMEN			ALL POSTPARTUM WOMEN	INFANTS AND CHILDREN	
		FIRST 1-13 WKS	SECOND 14-26 WKS	THIRD 27-40+ WKS		6-23 MO	2-5 YRS
<3000 ft. Plummer	NON- SMOKER	33.0	32.0	33.0	35.7	32.9	33.0
	SMOKER	34.0	33.0	34.0	36.7	---	---

Nutrition Risk Criteria **301 HYPEREMESIS GRAVIDARUM**

	Category	Priority	Referral
Category, Priority and Referral	PREGNANT WOMEN	1	RD

Definition	Severe nausea and vomiting to the extent that the pregnant woman becomes dehydrated and acidotic. Presence of Hyperemesis Gravidarum diagnosed by physician as self-reported by applicant/participant/caregiver, or as reported or documented by a physician, or someone working under physician's orders.
Required Documentation	Document source of information (i.e. self-report of diagnosis by physician; written physician diagnosis).
Justification	Nausea and vomiting are common early in gestation; 50% or more of normal pregnant women experience some vomiting. However, pregnant women with severe vomiting during pregnancy are at risk of weight loss, dehydration, and metabolic imbalances. Nutrition risk is based on chronic conditions, not single episodes.
Clarification	Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.
Reference	Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Nutrition Risk Criteria **302 GESTATIONAL DIABETES**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	PREGNANT WOMEN	1	RD

Definition

Gestational diabetes mellitus (GDM) is defined as any degree of glucose/ carbohydrate intolerance with onset or first recognition during pregnancy (1, 2).

Presence of gestational diabetes diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Document source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The definition of GDM applies regardless of whether insulin or only diet modification is used for treatment, or whether the condition persists after pregnancy. Included in this classification are women who may have had undiagnosed diabetes prior to pregnancy but who are first diagnosed during pregnancy (1, 2). Pregnant women requiring the use of exogenous steroids, tocolytics, or other medications, or who have medical conditions that alter glucose tolerance, may develop GDM (2). GDM represents nearly 90% of all pregnancies complicated by diabetes (1). The criteria for the diagnosis of GDM (3) are shown in Table 1 (see Clarification).

Pregnancy is an insulin-resistant and diabetogenic state (2). Deterioration of glucose tolerance occurs normally during pregnancy, particularly in the 3rd trimester (1, 2). Untreated or poorly treated GDM results in a higher risk of morbidity and mortality for both the mother and the fetus (2).

Established risk factors for GDM are advanced maternal age, obesity, and family history of diabetes (4). Risk assessment for GDM should be undertaken at the first prenatal visit. Women with clinical characteristics consistent with a high risk for GDM (e.g., those with marked obesity, personal history of GDM or delivery of a previous large-for-gestation-age infant, glycosuria, polycystic ovary syndrome, or a strong family history of diabetes) should undergo glucose testing as soon as possible (5). Unquestionably, there are also ethnic differences in the prevalence of GDM. In the U.S., Native Americans, Asians, Hispanics, and African American women are at a higher risk for GDM than non-Hispanic White women. Besides obesity, there is a suggestion that physical inactivity, diets high in saturated fat and smoking are associated with increasing risk for GDM or recurrent GDM (4).

Infants of women with GDM are at an increased risk of developing obesity, impaired glucose tolerance or diabetes as children or young adults (4). GDM is associated with a higher incidence of maternal and fetal complications. Maternal complications include polycythemia, respiratory distress syndrome, and increased rate of stillbirth (6). Although rarely seen in GDM, congenital anomalies, neural tube defects, cardiac abnormalities and/or caudal regression may occur if a woman has GDM in the early first trimester (6, 7).

Since GDM is a risk factor for subsequent type 2 diabetes after delivery, lifestyle modifications aimed at reducing weight and increasing physical activity are recommended (8). The National Diabetes Education Program (NDEP) is currently promoting a GDM Prevention Initiative, targeting both providers and women with a GDM history (((). Key messages are illustrated in Table 2 (see Clarification).

Medical Nutrition Therapy (MNT) is the primary treatment for the management of GDM (7). MNT for GDM primarily involves a carbohydrate-controlled meal plan that promotes optimal nutrition for maternal and fetal health with adequate energy for appropriate gestational weight gain, achievement and maintenance of normoglycemia, and absence of ketosis (7, 8). Breastfeeding should be strongly encouraged as it is associated with maternal weight loss and reduced insulin resistance for both mother and offspring (10). WIC nutrition services can reinforce and support the medical and diet therapies (such as MNT) that participants with GDM receive form their health care providers.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Women at high risk for GDM who have tested negative at the initial screening, and women at average risk for GDM should be tested by a license medical provider, between 24 and 28 weeks of gestation. Women of average risk should be tested at 24-28 weeks of gestation. Testing should follow one of two approaches:

- **One-step approach:** perform a diagnostic 100-g OGTT (Oral Glucose Tolerance Test)

- **Two-step approach:**
  1. A screening test (glucose challenge test) that measures plasma or serum glucose is done 1 hour after a 50 g oral glucose load without regard for time of day or time of last meal. If a plasma or serum glucose level meets or exceeds the threshold (≥ 130 mg/dl [7.2 mmol/L] or ≥ 140 mg/dl [7.8 mmol/L] respectively), an OGTT is performed (3).

  2. A diagnosis of GDM is made with a 100 g oral glucose load after an overnight fast. Using a 3-hour test, if two or more plasma or serum glucose levels meet or exceed the threshold, a diagnosis of GDM is made. Alternatively, the diagnosis can be made using a 75 g oral glucose load. The glucose threshold values for both tests are listed in Table 1 (10). The 75 g glucose load test is not as well validated as the 100 g OGTT.

With either the 75 g OGTT or the 100 g OGTT, it is recommended that the test be performed after an overnight fast of at least 8 hours but no longer than 14 hours. For 3 days prior to the test the woman should consume an unrestricted diet (≥ 150 g carbohydrate per day) and maintain unrestricted physical activity. Women need to remain seated and not smoke during the test. (1, 2).

Table 1. Diagnosis of Gestational Diabetes Mellitus with a 100 g or 75 g Oral Glucose Load

Time (h)	100 g Oral Glucose Load	75 g Oral Glucose Load
Fasting	95 mg/dL (5.3 mmol/L)	95 mg/dL (5.3 mmol/L)
1	180 mg/dL (10.0 mmol/L)	180 mg/dL (10.0 mmol/L)
2	155 mg/dL (8.6 mmol/L)	155 mg/dL (8.6 mmol,/L)
3	140 mg/dL (7.8 mmol/L)	

Two or more of the venous plasma concentrations must be met or exceeded for a positive diagnosis.
Source: American Diabetes Association (3).

Table 2. Gestational Diabetes Mellitus (GDM) Prevention Initiative from the National Diabetes Education Program

- GDM imparts lifelong risk for diabetes, mostly type 2
- Modest weight loss and physical activity can delay or prevent type 2 diabetes
- Offspring can lower risk of diabetes by eating healthy foods, being active, and not becoming overweight

Conservative recommendations to patients include:

- Let health care practitioners know of any history of GDM.
- Get glucose testing at 6 to 12 weeks postpartum, then every 1-2 years.
- Reach pre-pregnancy weight 6 to 12 months postpartum.
- If still overweight, lose at least 5 to 7% of weight slowly, over time, and keep it off.

Adapted from the National Diabetes Education Program (9).

References

1. American Diabetes Association: Diagnosis and classification of diabetes mellitus. Diabetes Care, Jan 2008; 31 Suppl 1:S55-60.
2. Franz MJ, Biastre SA, Slocum J, Diabetes in the life cycle and research. In: Gestational Diabetes – A core curriculum for diabetes education, American Association of Diabetes Educators. 5th ed. 2003.
3. American Diabetes Association. Gestational diabetes mellitus (position statement), Diabetes Care. 2003; 26 Suppl 1:S103-105.
4. Ferrara, A. Increasing prevalence of gestational diabetes mellitus: a public health perspective. Proceedings of the fifth international workshop – conference on Gestational Diabetes Mellitus. Diabetes Care. Jul 2007; 30 Suppl 2:S141-46.
5. American Diabetes Association. Standards of medical care in diabetes (position statement). Diabetes Care. Jan 2007; 30 Suppl 2:S4-41.
6. Thomas AM, Gutierrez YM. American Dietetic Association guide to gestational diabetes mellitus in postpartum considerations. Eds. American Dietetic Association; 2005:101-113.
7. Brian SR, Nickless N, Thung SF, Inzucchio SE. Gestational diabetes update: screening, medical management and follow-up. Practical Diabetology. Mar 2007; 10-18.
8. American Diabetes Association. Diagnosis and classification of diabetes mellitus. Diabetes Care. 2008 Jan; 31 Suppl 1:S55-60.
9. Ratner, RE. Prevention of type 2 diabetes in women with previous gestational diabetes. Proceedings of the fifth international workshop – conference on gestational diabetes mellitus. Diabetes Care. Jul 2007; 30 Suppl 2:S242-245.

10. Evert AG, Vande Hei K. Gestational diabetes education and diabetes prevention strategies. Diabetes Spectrum. 2006; 19(3):135-139.

Nutrition Risk Criteria **303 HISTORY OF GESTATIONAL DIABETES**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD

Definition

History of diagnosed gestational diabetes mellitus (GDM).

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Document source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Women who have had a pregnancy complicated by GDM are 40-60% more likely to develop diabetes within 15-20 years (1), usually type 2 (2). This risk of subsequent diabetes is greatest in women with GDM who are diagnosed early in the pregnancy, exhibit the highest rates of hyperglycemia during the pregnancy, and are obese.

Approximately 30-50% of the women with a history of GDM will develop GDM in a subsequent pregnancy. Studies have found that the risk factors for subsequent GDM include insulin use in the index pregnancy, obesity, diet composition*, physical inactivity, failure to maintain a healthy BMI and weight gain between pregnancies (2, 3). In addition, if a woman's lipid levels are elevated, a history of GDM is also a risk factor for cardiovascular disorders (3).

There is evidence to suggest that some women with a history of GDM show relative beta-cell dysfunction during and after pregnancy (3). Most women with a history of GDM are insulin resistant. Changes in lifestyle (dietary and physical activity) may improve postpartum insulin sensitivity and could possibly preserve B-cell function to slow the progression to type 2 diabetes (2, 3).

During WIC nutrition education and counseling, obese women with a history of GDM should be encouraged to lose weight before a subsequent pregnancy. Breastfeeding has been shown to lower the blood glucose level and to decrease the incidence of type 2 diabetes in women with a history of GDM (2, 3). Exercise also has a beneficial effect on insulin action by enhancing peripheral tissue glucose uptake (3). Medical Nutrition Therapy (MNT) is an essential component in the care of women with a history of GDM.

Women with a history of GDM but without immediate subsequent postpartum diagnosis of diabetes should be advised to discuss with their medical provider the importance of having a Glucose Tolerance Testing (GTT) at 6 to 12 weeks postpartum (see Clarification, Table 1); to have a pre-pregnancy consultation before the next pregnancy, and to request early glucose screening in the next pregnancy (4). The National Diabetes Education Program (NDEP) is currently promoting a GDM Diabetes Prevention Initiative, targeting both providers and

women with a history of GDM (5). Key messages are illustrated in Table 2. (See Clarification).

WIC nutrition services can support and reinforce the MNT and physical activity recommendations that participants receive from the health care providers. In addition, WIC nutritionists can play an important role in providing women with counseling to help manage their weight after delivery. Also, children of women with a history of GDM should be encouraged to establish and maintain healthy dietary and lifestyle behaviors to avoid excess weight gain and reduce their risk for type 2 diabetes (1).

***Diet Composition**

Carbohydrate is the main nutrient that affects postprandial glucose evaluations. During pregnancy complicated with GDM, carbohydrate intake can be manipulated by controlling the total amount of carbohydrate, the distribution of carbohydrate over several meals and snacks, and the type of carbohydrate. These modifications need not affect the total caloric intake level/prescription (6).

Because there is wide inter-individual variability in the glycemic index each woman needs to determine, with the guidance of the dietitian, which foods to avoid or use in smaller portions at all meals or during specific times of the day, for the duration of her pregnancy. Practice guidelines have avoided labeling foods as “good” or “bad”

Meal plans should be culturally appropriate and individualized to take into account the patient’s body habits, weight gain and physical activity; and should be modified as needed throughout pregnancy to achieve treatment goals (6).

Clarification

Self-reporting for “History of ”conditions should be treated in the same manner as self-reporting for current conditions requiring a physicians diagnosis, i.e., the applicant may report to the CPA that she was diagnosed by a physician with a given condition at some point in the past. As with current conditions, self-diagnosis of a past condition should never be confused with self-reporting.

Table 1. Reasons for Delayed Postpartum Glucose Testing of Women with Prior Gestational Diabetes Mellitus (GDM)

1. The substantial prevalence of glucose abnormalities detected by 3 months postpartum.
2. Abnormal test results identify women at high risk of developing diabetes over the next 5 to 10 years.
3. Ample clinical trial evidence in women with glucose intolerance that type 2 diabetes can be delayed or prevented by lifestyle interventions or modest and perhaps intermittent drug therapy.
4. Women with prior GDM and impaired glucose tolerance (IGT) have cardiovascular disease (CVD) risk factors. Interventions may reduce subsequent CVD, which is the leading cause of death in both types of diabetes.
5. Identification, treatment, and planning of pregnancy in women developing diabetes after GDM should reduce subsequent early fetal loss and major congenital malformations.

Kitzmiller JL, Dang-Kilduff L, Taslimi MM

Table 2. Gestational Diabetes Mellitus (GDM) Prevention Initiative form the National Diabetes Education Program

- GDM imparts lifelong risk for diabetes, mostly type 2.
- Modest weight loss and physical activity can delay or prevent type 2 diabetes.
- Offspring can lower risk by eating healthy foods, being active, and not becoming overweight.

Conservative recommendations to patients include:

- Let health care practitioners know of any history of GDM.
- Get glucose testing at 6 to 12 weeks postpartum, then every 1-2 years.
- Reach prepregnancy weight 6 to 12 months postpartum.
- If still overweight, lose at least 5 to 7 % of weight slowly, over time, and keep it off.

Adapted from the National Diabetes Education Program.

References

1. Evert AG, Vande Hei K. Gestational diabetes education and diabetes prevention strategies. Diabetes Spectrum. 2006; 19 (3):135-139.
2. Franz MJ, Biastre SA, Slocum J. Diabetes in the life cycle and research. In: Gestational diabetes – A core curriculum for diabetes education, American Association of Diabetes Educators. 5th ed. 2003; 145-163.
3. Thomas AM, Gutierrez YM. American Dietetic Association guide to gestational diabetes mellitus in postpartum considerations. Eds. American Dietetic Association. 2005; 101-113.
4. Kitzmiller JL, Dang-Kilduff L, Taslimi MM. Gestational diabetes after delivery: short-term management and long-term risks. Proceedings of the fifth international workshop – conference on Gestational Diabetes Mellitus. Diabetes Care. Jul 2007; 30 Suppl 2:S225-231.
5. Ratner RE. Prevention of type 2 diabetes in women with previous gestational diabetes. Proceedings of the fifth international workshop—conference on Gestational Diabetes Mellitus. Diabetes Care. Jul 2007; 30 Suppl 2:S242-245.
6. Reader DM. Medical nutrition therapy and lifestyle interventions. Proceedings of the fifth international workshop – conference on Gestational Diabetes Mellitus. Diabetes Care. Jul 2007; 30 Suppl 2:S188-193.

Nutrition Risk Criteria **304 HISTORY OF PREECLAMPSIA**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

History of diagnosed preeclampsia.

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under a physician's orders.

Required Documentation

Document source of information (i.e. self-report of physician diagnosis or written documentation from physician).

Justification

Preeclampsia is defined as pregnancy-induced hypertension (> 140mm Hg systolic or 90mm diastolic) with proteinuria developing usually after the twentieth week of gestation (1, 2). Clinical symptoms of preeclampsia may include edema, renal failure, and the HELLP syndrome (Hemolysis, Elevated Liver enzymes and Low Platelets).

Preeclampsia is a leading cause of maternal death and a major contributor to maternal and perinatal morbidity (3). Women who have had preeclampsia in a prior pregnancy have an increased risk of recurrence (about 20% overall) (4). The risk is greater in women who have had preeclampsia occurring early in pregnancy or who have had preeclampsia in more than one pregnancy. Additionally, maternal pre-pregnancy obesity with BMI ≥ 30 is the most prevalent risk factor for preeclampsia (4).

Risk factors for preeclampsia include (2, 4, 5):

- Pre-pregnancy obesity BMI ≥ 30
- Preeclampsia in a prior pregnancy
- Nulliparity (no prior delivery)
- Maternal age > 35 years
- Endocrine disorders (e.g., diabetes),autoimmune disorders (e.g., lupus), renal disorders
- Multi-fetal gestation
- Genetics
- Black race

There are few established nutrient recommendations for the prevention of preeclampsia. However, vitamin D may be important because it influences vascular structure and function, and regulates blood pressure (4). Also, calcium may prevent preeclampsia among women with very low baseline calcium intake (4).

There is no treatment for preeclampsia. The condition resolves itself only when the pregnancy terminates or a placenta is delivered (4). Early prenatal care, therefore, is vital to the prevention of the onset of the disease.

WIC is well poised to provide crucial strategies during the critical interconceptual period to help reduce the risk of recurrence of preeclampsia in a subsequent pregnancy.

WIC nutrition education encourages practices shown by research to have a protective effect against developing preeclampsia (2, 4, 5). These include:

- Gaining recommended weight based on pre-pregnancy BMI, in order to help return to a healthy postpartum weight
- Scheduling early prenatal care visits
- Consuming a diet adequate in calcium and vitamin D
- Taking prenatal vitamins
- Engaging in regular physical activity
- Discontinuing smoking and alcohol consumption

Postpartum Women: Women who have had preeclampsia should be advised that they are at risk for recurrence of the disease and development of cardiovascular disease (CVD) later in life (4, 7). WIC nutrition education can emphasize measures that support the prevention of preeclampsia in a future pregnancy, such as reaching or maintaining a healthy BMI and lifestyle between pregnancies, consuming a nutritionally adequate diet consistent with the Dietary Guidelines for Americans, and engaging in regular physical activity.

Pregnant Women: The WIC Program provides supplemental foods rich in nutrients, especially calcium and vitamin D, which research has shown to have a protective effect on preeclampsia (4). During nutrition education, WIC can encourage actions or behaviors that also have been shown to have a protective effect against preeclampsia: early prenatal care, taking a prenatal vitamin, and engaging in physical activity (6). WIC can also discourage smoking and alcohol consumption (2), and counsel pregnant women to gain recommended weight based on pre-pregnancy BMI (8) and to return to pre-pregnancy weight or a healthy BMI of < 25 for the benefit of future pregnancies.

Clarification

Self-reporting for “history of...” conditions should be treated in the same manner as self-reporting for current conditions requiring a physician’s diagnosis, e.g., the applicant may report to the CPA that she was diagnosed by a physician with a given condition at some point in the past. As with current conditions, self-diagnosis of a past condition should never be confused with self-reporting.

References

1. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
2. American Dietetic Association, Pediatric Nutrition Practice Group. Pediatric manual of clinical dietetics. Chicago: Pediatric Nutrition Dietetic Practice Group, American Dietetic Association, 1998.
3. Stenson W. The esophagus and stomach. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins 1999. p. 1125-1133.
4. Beyer PL. Medical nutrition therapy for upper gastrointestinal tract disorders. In: Mahan LK, Escott-Slump S, editors. Krause’s food nutrition and diet therapy. 11th ed. Philadelphia: Saunders; 2004. p. 688-690.

5. American Dietetic Association. Nutrition Care Manual. Gastrointestinal disease; Peptic ulcers; 2006. http://www.nutritioncaremanual.org. Accessed 1/08.
6. Allied Health Sciences Section Ad Hoc Nutrition committee: Aills L, Blankenship J, Buffington C, Furtado M and Parrot J. Bariatric nutrition: suggestions for the surgical weight loss patient. Review. Surgery for Obesity and Related Diseases 2008 May 17.
7. Scolapio JS, Fleming R. Short Bowel Syndrome. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins; 1999. p. 1135-1140.
8. Farrell M. Gastrointestinal disorders of infancy and childhood (with nutrition support and probiotics) In: Ekvall SW, Ekvall VK. Editors. Pediatric nutrition in chronic diseases and developmental disorders. 2nd ed. Oxford University Press; 2005. p. 248-249.
9. Griffiths A. Inflammatory bowel disease. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins; 1999. p. 1141-1149.
10. Lieber CS. Nutrition in liver disorders. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams 7 Wilkins; 1999. p. 1177-1189.
11. Raimondo m, Dimagno EP. Nutrition in pancreatic disorders. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkinds; 1999. p. 1169-1176.
12. Hasse JM, Matarese JE. Medical nutrition therapy for liver, biliary system and exocrine pancreas disorders. In: Mahan LK, Escott-Stump S, editors. Krause's food nutrition and diet therapy. 11th ed. Philadelphia: Saunders; 2004. p. 758-760.

Nutrition Risk Criteria **311 HISTORY OF PRETERM OR EARLY TERM DELIVERY**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

History of preterm and/or early term delivery is defined as follows (1, 2):

- Preterm: Delivery of an infant born less than or equal to (≤)36 6/7 weeks.
- Early Term: Delivery of an infant born greater than or equal to (≥)37 0/7 weeks and less than or equal to (≤)38 6/7 weeks (also known as less than <39 weeks).

Pregnant Women: any history of preterm or early term delivery
Breastfeeding/Non-breastfeeding: most recent pregnancy

Required Documentation

Documentation of DOB, EDD, or history of preterm or early term delivery

Justification

Women with a history of preterm delivery have an increased risk of spontaneous preterm delivery in a subsequent pregnancy compared to women with no history of prior spontaneous preterm delivery (3). Prior spontaneous preterm delivery is highly associated with recurrence in subsequent pregnancies. A history of one previous preterm birth is associated with a recurrent risk of 17-37%; the risk increases with the number of prior preterm births and decreases with the number of term deliveries (4).

Typically a pregnancy lasts about 40 weeks. Premature or preterm birth, however, is defined as a birth that occurs between 20 and 37 weeks of pregnancy, according to the American College of Obstetricians and Gynecologists (ACOG) (5). In the past, the period from 3 weeks before until 2 weeks after the estimated date of delivery was considered a “term” pregnancy, with the expectation that a baby would have similar health outcomes if they were born any time during this interval. In 2013, ACOG released a committee opinion that the label “term” should be replaced with the designations *early term* (≥37 0/7 weeks and ≤38 6/7 weeks gestation) and *full term* (≥39 0/7 weeks and ≤40 6/7 weeks gestation) to more accurately describe these groups of infants (1).

Preterm Delivery

Prematurity affects about 12% of all live births in the U. S., and about 50% of these preterm births were preceded by preterm labor (6). In 2011, the annual rate of premature births in the United States reached 11.7%, nearly two times the rate in European nations (6). Preterm births also account for approximately 70% of newborn deaths and 36% of infant deaths (5).

Despite advances in neonatal care, preterm birth remains a leading cause of infant death in the United States (7). More infants die from pre-term related problems than any other single cause (6). Preterm birth strains society’s healthcare resources due to its long-term effects on the health of the newborn

(6). Premature infants may have physical problems that have nutritional implications, including immature sucking, swallowing and immature digestion and absorption of carbohydrates and lipids (7). Preterm infants are at risk for a number of illnesses/health conditions that range from minor to severe complications depending on the circumstances. (See risk 142 *Preterm or Early Term Delivery* for more details.)

Several factors have been found to increase the risk of preterm delivery. Epidemiologic studies have consistently reported low socioeconomic status, nonwhite race, maternal age of ≤18 years or ≥40 years, and low pre-pregnancy underweight as risk factors (4). Studies suggest even modest restrictions in maternal nutrition around the time of conception can lead to premature births and long-term adverse health effects for offspring (8). Other factors associated with a risk of preterm birth may be identified before pregnancy, at conception, or during pregnancy include (8, 9):

- Low maternal weight gain during pregnancy
- Maternal infections
- Maternal hypertension
- Gestational diabetes
- Smoking
- Indoor pollution
- Maternal stress
- Poor housing quality
- Teen pregnancy
- Sexually transmitted diseases
- Low psychosocial health status
- Previous or present pregnancy complications
- Multiple fetuses
- Lack of perceived social support

A recent study indicated that maternal obesity is also an independent risk factor for preterm delivery (10). Complications associated with obesity (BMI ≥30) prior to conception that increase the risk for preterm delivery include (11):

- Gestational Diabetes Mellitus
- Hypertension
- Preeclampsia
- Cesarean Delivery
- Postpartum weight retention

Additional concerns related to obesity include potential intrapartum, operative, and postoperative complications and difficulties related to anesthesia management. Obese women are also less likely to initiate and sustain breastfeeding (11).

Breastfeeding is recommended as the normative standard for infant feeding and nutrition for all infants, especially preterm babies. Breastfeeding preterm infants has been associated with positive health outcomes for these infants, including:

- Improved motor maturity and cognitive ability (12, 13, 14)
- Reduced risk of necrotizing enterocolitis (15, 16)
- Reduced risk of retinopathy of prematurity and retinal detachment (17)

Additionally, mothers of preterm infants produce milk that is designed to meet the baby's particular needs during the first few weeks of breastfeeding. It is higher in protein and minerals, such as salt, and contains different types of fat that the baby will be able to digest and absorb more easily compared to the milk of mothers of full term babies. The fat in human milk also helps to enhance the development of the baby's brain and neurologic tissues, which is especially important for premature infants. Human milk is also easier for babies to digest than formula and avoids exposing the baby's immature intestinal lining to the cow's milk proteins found in premature infant formula. Preterm infants who are breastfed are less likely to develop intestinal infections than babies who are formula fed, and the colostrum produced in the first few days contains high concentrations of antibodies that will also help the baby fight infection. (16)

Breastfeeding preterm infants, especially if they are in the NICU, may present unique challenges for breastfeeding dyads. These mothers will benefit from extra breastfeeding support due to the delay of direct breastfeeding, reliance on breast pumps, and the stress of having a sick newborn. Even if the baby cannot breastfeed directly from the breast at first, the mother can be encouraged to express her milk to ensure that her supply is maintained. Supportive care for infants in the NICU may include the use of a feeding tube. Expressed human milk can be passed through the tube, so it is important for the mother to discuss her feeding decisions with her baby's doctor.

Early Term

Up to 10% of babies in the United States are scheduled for early term deliveries via labor-inducing medication or cesarean section before 39 weeks of gestation despite neither the mother nor the baby being at risk if the pregnancy continues (18). Elective deliveries like this are sometimes requested for reasons such as wanting to schedule the date of the infant's birth, physician preference, or for relief of symptoms at the end of the pregnancy (18).

Research shows that a fetus will experience a significant amount of development and growth of the lungs, brain, and liver between 37 and 39 weeks of gestation. The brain develops at its fastest rate at the end of the pregnancy, at a rate of up to one third between weeks 35 and 39. Additionally, layers of fat are added under the infant's skin during the last few weeks of pregnancy which helps them keep warm after birth. According to ACOG, non-medically warranted deliveries prior to 39 weeks should be avoided (19). Early term delivery puts an additional strain on society as the early term infant will likely require a longer hospital stay and may have long term healthcare needs (18). Factors that can increase the risk of woman delivering an early term infant are the same and are stated above for preterm birth.

When a woman delivers an early term infant or chooses an early elective delivery, she is at increased risk for postpartum depression, cesarean delivery, and other complications requiring longer hospital stays (18). Steps pregnant women can take in order to decrease the prevalence of pre-term births include (18):

- Seek regular prenatal care throughout pregnancy.
- Maintain a healthy diet, including daily prenatal vitamins.
- Cease consumption of alcohol, drugs, or other dangerous toxins during pregnancy.
- Avoid stress.

- Contact their health care provider with all questions or concerns.

Pregnant women who come from low or inadequate income households are at a greater risk for poor physical and mental health due to poor eating habits. WIC services may assist women at risk of preterm and early term births by providing them with proper nutrition.

Implication for WIC Nutrition Services

Early prevention is the primary way to stop preterm labors. WIC can assist in reducing preterm deliveries by increasing prevention strategies. WIC can improve outcomes through:

- Recommending healthy maternal weight gain and providing nutrition education that addresses the WIC food package and other healthy foods that contribute to a balanced diet.
- Promoting early and regular prenatal care.
- Encouraging use of prenatal vitamins, as prescribed by the health care provider.
- Recommending adherence to Dietary Guidelines for Americans.

WIC staff may find the below listed resources helpful in providing nutrition counseling:

- *Additional Considerations for Some Adults-Physical Activity for Women During Pregnancy and the Postpartum Period*: http://health.gov/paguidelines/guidelines/chapter7.aspx .
- *Women, Infants, and Children-About WIC, How WIC Helps*: http://www.fns.usda.gov/wic/about-wic-how-wic-helps .
- *WIC Works Resource Systems*: https://wicworks.fns.usda.gov/ .

References

1. American College of Obstetricians and Gynecologists. Definition of term pregnancy. Committee Opinion No.579. Obstet Bynecol. 2013 Nov;122:1139-40.
2. Ob-Gyns redefine meaning of “term pregnancy” [Internet]. Washington, DC: American College of Obstetricians and Gynecologists; c2013 [updated 2013 Oct 22; cited 2016 Dec 6]. Available from: http:www.acog.org/About-ACOG/News-Room/News-Releases/2013/Ob-Gyns-Redefine-Meaning-of-Term-Pregnancy.
3. Mercer M. Moawad A, Meis P, Iams JD, Das A, Caritis SN. The preterm prediction study: effect of gestational age and cause of the preterm on subsequent obstetric outcome. Am J Obstet Gynecol. 1999 Nov;181:1219-21.
4. Hoffman HJ, Bakketeig LS. Risk factors associated with the occurrence of preterm birth. Clin.Obstet.Gynecol. 1984; 27:539-52.
5. ACOG.org [Internet]. Washington DC: The American College of Obstetricians and Gynecologists: c2016 [updated 2016 Nov; cited 2016 Dec 6]. Available from: http://www.acog.org/Patients/FAQs/Preterm-Premature-Labor-and-Birth
6. Dag M, Lie TR, Markestad T. Long-term medical and social consequences of preterm birth. N Engl J Med. 2008 Jul 17. Web. 07 Apr. 2014.

7. Iams JD. Prevention of preterm parturition. N Engl of Med. 2014;370(3):254-261.

8. Wang P, Liou S, Cheng C. Prediction of maternal quality of life on preterm birth and low birthweight: a longitudinal study. BMC Pregnancy and Childbirth [serial online]. June 2, 2013;13:124. Available from: MEDLINE, Ipswich, MA. Accessed April 28, 2014.

9. Ludwig JD, Miller M. Interpreting the WIC cebate. J Pol Anal Manage. 2005; 24(4):691-701.

10. Cnattinqius S, Villamor E, Johansson S. Edstedt, Bonamy AK, Persson M, et al. Maternal obesity and risk of preterm delivery. JAMA 2013 Jun 12:309(22).

11. American College of Obstetricians and Gynecologists. Practice Bulletin. OPbesity in pregnancy. Dec. 2015.

12. Feldman R, Eidelman A. Direct and indirect effects of breast milk on the neurobehavioral and cognitive development of premature infants. Developmental Psychobiology. 2003 Sept;43(2):109-19.

13. Vohr B, Poindexter B, Dusick A, McKinely LT, Higgins RD, Langer JC, Poole KW. Persistent beneficial effects of breast milk ingested in the neonatal internsive care unit on outcomes of extremely low birth weight infants at 30 months of age. Pediatrics. 2007 Oct;120(4):e953-59.

14. Blaymore Bier J, Oliver T, Ferguson AE, Vohr B. Human Milk Improves Cognitive and Motor Development of premature infants during infancy. J Hum Lact. 2002 Nov;18(4)361-67.

15. Boyd CA, Quigley MA, Brocklehurst P. Donor breast milk versus infant formula for preterm infants: systematic review and meta-analysis. Arch Dis Child Fetal Neonatal Ed. 2007 May;92:F169-75.

16. Healthychildren.org [Internet]. Elk Grove Village: American Academy of Pediatrics; c2011 [updated 2015 Nov 21; cited 2016 Dec 6]. Available from: https://www.healthychildren.org/English/ages-stages/baby/breastfeeding/Pages/Providing-Breatmilk-for-Premature-and-Ill-Newborns.aspx .

17. Okamoto T, Shirai M, Kokubo M, Takahashi S, Kajino M, Takase M, Sakata H, Oki J. Human milk reduces the risk of retinal detachment in extremely low-birthweight infants. Pediatr Int. 2007 Oct;49(6):894-897.

18. National Institute for Health Care Management. Born too early-improving maternal and child health by reducing early elective deliveries. NIHCM Issue Brief, March 2014. [cited 2016 Dec 6]. Available from: http://www.nihcm.org/pdf/Early_Elective_Delivery_Prevention_Brief_2014.pdf .

19. Elective delivery before 39 weeks [Internet]. Washington, DC: American College of Obstetricians and Gynecologists; c2013 [updated 2013 June; cited 2016 Dec 6]. Available from: http://www.acog.org/Patients/FAQs/Elective-Delivery-Before-39-Weeks .

Nutrition Risk Criteria **312 HISTORY OF LOW BIRTH WEIGHT**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Birth of an infant weighing ≤ 5 lbs 8 oz (≤ 2500 g)

Pregnant Women: any history of low birth weight
Breastfeeding/Non-breastfeeding: most recent pregnancy

Required Documentation

Justification

A woman's history of a delivery of a low birth weight (LBW) baby is the most reliable predictor for LBW in her subsequent pregnancy (1). The risk for LBW is 2-5 times higher than average among women who have had previous LBW deliveries and increases with the number of previous LBW deliveries (1). This is true for histories in which the LBW was due to premature birth, fetal growth restriction (FGR) or a combination of these factors. The extent to which nutritional interventions (dietary supplementation and counsel) can decrease risk for repeat LBW, depends upon the relative degree to which poor nutrition was implicated in each woman's previous poor pregnancy outcome. Nutritional deficiencies and excesses have been shown to result in LBW and pregnancy loss. The pregnant woman's weight gain is one of the most important correlates of birth weight and of FGR (2, 3).

References

1. Institute of Medicine, Committee to Study the Prevention of Low Birth Weight. Preventing low birth weight. National Academy Press, Washington, D.C.; 1985.
2. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
3. Kramer MS. Intrauterine growth and gestational duration determinants. Pediatrics 1987; 80:502-11.

Nutrition Risk Criteria **321 HISTORY OF SPONTANEOUS ABORTION, FETAL OR NEONATAL LOSS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

A spontaneous abortion (SAB) is the spontaneous termination of a gestation at < 20 weeks gestation or < 500 g.
Fetal death is the spontaneous termination of a gestation at ≥ 20 weeks
Neonatal death is the death of an infant within 0-28 days of life.

Pregnant women: any history of fetal or neonatal death or 2 or more spontaneous abortions
Breastfeeding women: most recent pregnancy in which there was a multifetal gestation with one or more fetal or neonatal deaths but with one or more infants still living
Non-breastfeeding: most recent pregnancy

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Document source of information (i.e. as self-reported diagnosis by physician or written documentation by physician).

Justification

Pregnancy:

Previous fetal and neonatal deaths are strongly associated with preterm low birth weight (LBW) and small for gestational age (SGA) and the risk increases as the number of previous poor fetal outcomes goes up.

Spinnillo et al found that the risk for future small for gestational age outcomes increased two fold if a woman had 2 or more SAB. Adverse outcomes related to history of SAB include recurrent SAB, low birth weight (including preterm and small for gestational age infants), premature rupture of membranes, neural tube defects and major congenital malformations. Nutrients implicated in human and animal studies include energy, protein, folate, zinc, and vitamin A.

Postpartum women:

A SAB has been implicated as an indicator of a possible neural tube defect in a subsequent pregnancy. Women who have just had a SAB or a fetal or neonatal death should be counseled to increase their folic acid intake and delay a subsequent pregnancy until nutrient stores can be replenished.

The extent to which nutritional interventions (dietary supplementation and counseling) can decrease the risk for repeat poor pregnancy outcomes depends upon the relative degree to which poor nutrition was implicated in each woman's

previous poor pregnancy outcome. WIC Program clients receive foods and services that are relevant and related to ameliorating adverse pregnancy outcomes. Specifically, WIC food packages include good sources of implicated nutrients. Research confirms that dietary intake of nutrients provided by WIC foods improve indicators of nutrient status and/or fetal survival in humans and/or animals.

NOTE: A woman who becomes pregnant within 16 months after a SAB (her first) would qualify for risk #332, Closely Spaced Pregnancies.

Clarification

Self-reporting for "History of…" conditions should be treated in the same manner as self-reporting for current conditions requiring a physicians diagnosis, i.e., the applicant may report to the CPA that she was diagnosed by a physician with a given condition at some point in the past. As with current conditions, self-diagnosis of a past condition should never be confused with self-reporting.

References

1. American College of Obstetricians and Gynecologists. Preterm Labor. Technical Bulletin 206. Washington, DC: ACOG, 1995.
2. Carmi R, Gohar J, Meizner I, Katz M. Spontaneous abortion--high risk factor for neural tube defects in subsequent pregnancy [see comments]. Am.J.Med.Genet. 1994; 51:93-7.
3. Institute of Medicine, Committee to Study the Prevention of Low Birth Weight. Preventing low birth weight. National Academy Press, Washington, D.C.; 1985.
4. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
5. Kramer MS. Intrauterine growth and gestational duration determinants. Pediatrics 1987; 80:502-11.
6. Paz JE, Otano L, Gadow EC, Castilla EE. Previous miscarriage and stillbirth as risk factors for other unfavorable outcomes in the next pregnancy. Br.J.Obstet.Gynecol. 1992; 99:808-12.
7. Shapiro S, Ross LF, Levine HS. Relationship of selected prenatal factors to pregnancy outcome and congenital anomalies. Am.J.Public Health 1965; 55; 2:268-282.
8. Spinillo A, Capuzzo E, Piazzi G, Nicola S, Colonna L, Iasci A. Maternal high-risk factors and severity of growth deficit in small for gestational age infants. Early Hum.Dev. 1994; 38:35-43.
9. Thorn DH. Spontaneous abortion and subsequent adverse birth outcomes. Am.J.Obstet.Gyn. 1992; 111-6.

Nutrition Risk Criteria **331★ PREGNANCY AT YOUNG AGE**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Conception ≤ 17 years of age.

Pregnant Women: current pregnancy
Breastfeeding/Non-breastfeeding: most recent pregnancy

Required Documentation

DOB
EDC

Justification

Pregnancy before growth is complete, is a nutritional risk because of the potential for competition for nutrients for the pregnancy needs and the woman's growth.

The pregnant teenager is confronted with many special stresses that are superimposed on the nutritional needs associated with continued growth and maturation.

Younger pregnant women of low socioeconomic status tend to consume less than recommended amounts of protein, iron, and calcium, and are more likely to come into pregnancy already underweight. Pregnant teens who participate in WIC have been shown to have an associated increase in mean birth weight and a decrease in LBW outcomes.

Adolescent mothers frequently come into pregnancy underweight, have extra growth related nutritional needs, and because they often have concerns about weight and body image, are in need of realistic, health promoting nutrition advice and support during lactation. Diets of adolescents with low family incomes typically contain less iron, and less vitamin A than are recommended during lactation.

The adolescent mother is also confronted with many special stresses superimposed on the normal nutritional needs associated with continued growth. Nutrition status and risk during the postpartum period follow from the nutritional stresses of the past pregnancy, and in turn have an impact on nutrition related risks in subsequent pregnancies. Poor weight gain and low intakes of a variety of nutrients are more common in pregnant adolescents. Therefore, participation in the WIC Program should be of substantial benefit.

References

1. Endres J, Dunning S, Poon SW, Welch P, Duncan H. Older pregnant women and adolescents: nutrition data after enrollment in WIC. J.Am.Diet.Assoc. 1987; 87:1011-6, 1019.
2. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
3. Kennedy ET, Kotelchuck M. The effect of WIC supplemental feeding on birth weight: a case-control analysis. Am.J.Clin.Nutr. 1984; 40:579-85.
4. Story M, editor. Nutrition management of the pregnant adolescent a practical reference guide. Washington, D.C.: National Clearinghouse; 1990. Sponsored by the March of Dimes Birth Defects Foundation, U.S. Department of Health and Human Services, U.S. Department of Agriculture.

Nutrition Risk Criteria **332★ SHORT PREGNANCY INTERVAL**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Short Interpregnancy Interval, formerly known as *Closely Spaced Pregnancies*, is defined as an interpregnancy interval (IPI) of less than 18 months from the date of a live birth to the conception of the subsequent pregnancy for:

Pregnant women - current pregnancy
Breastfeeding/non-breastfeeding women - most recent pregnancy

Note: The evidence-based information supporting this criterion is specific to live births and did not include women who had miscarriages or stillbirths. Thus, the definition for this criterion is specific only to women who experienced live births. Women whose pregnancies did not result in a live birth may be assigned, as appropriate, risk *321 History of Spontaneous Abortion, Fetal or Neonatal Loss*.

Required Documentation

EDC
Previous pregnancy end date

Justification

Adverse maternal and infant health outcomes have been associated with short Interpregnancy Intervals (IPIs). While there is no standard definition for short IPI, and IPI less than 18 months has been associated with increased risk for adverse outcomes (1, 2). An interval of 18 to 24 months has been associated with the lowest relative risk (2). Evidence associated with the lowest relative risk for an IPI following a miscarriage or abortion is still unclear (see Clarification Section for more information) therefore only health effects associated with a short IPI following a live birth were reviewed for this criterion.

Historically, the World Health Organization (WHO) and other international authorities had recommended at least 2-3 years between pregnancies and the United States Agency for International Development (USAID) had suggested an interval of 3-5 years. Given the inconsistency, various counties and regional programs requested the WHO to further review the research and provide recommendations. As a result, the report from the 2005 WHO Technical Consultation and Scientific Review of Birth Spacing recommended an interval of at least 24 months after a live birth to reduce the risk of adverse maternal, perinatal, and infant outcomes (3). A more recent review of data suggests that there are increased risks for adverse perinatal and maternal outcomes with an IPI less than 18 months (1, 2, 4) and increased risks for perinatal (1, 4) and maternal (4, 5, 6) outcomes longer than 59 months while 18-24 months was associated with the lowest relative risk (2). Parallel to recent findings, Healthy People 202 has proposed a 10% improvement in reducing the proportion of pregnancies conceived within 18 months of a previous birth (7).

Outcomes associated with short IPI have included maternal complications such as uterine rupture in women attempting a vaginal birth after a previous cesarean

delivery (also referred to as VBAC) (8, 9); and perinatal and neonatal complications such as preterm birth (1, 2, 10), low birth weight (1, 2), small for gestational age (1, 2), birth defects (11), and autism (12, 13).

Short IPI has been identified as a risk for increasing uterine rupture in women attempting a VBAC delivery (8, 9, 14). Yet when comparing short interpregnancy interval to labor type – induced labor and spontaneous, there was a decrease rate in VABC success in women who were induced, and no difference with spontaneous labor (15). Given the lack of a specific IPI recommendation for women with a previous cesarean delivery and the inconsistencies in study designs there appears to be no specific guidelines for interval length after a cesarean delivery (16). The short interpregnancy interval definition cut-off of 18 months, however, appears to be inclusive of women who delivered by cesarean with their previous pregnancy.

Factors contributing to adverse outcomes and short IPI remain controversial. It was thought that socioeconomic factors contributed to adverse outcomes. However, when controlled for possible cofounders, short IPI remained an independent risk factor (1, 2). Nutrition-related hypothetical causal mechanisms have been proposed to explain the effects short IPIs have on health, yet research remains inconclusive (4). The Maternal Depletion Syndrome hypothesized that mothers who have a short IPI often do not have adequate time to replenish macro- and micro-nutrients which may lead to the mother and fetus competing for nutrients (17). However, a recent systematic review of the literature found no evidence to support this hypothesis (4). Studies to support the folate depletion theory have had differing results (11, 18). When folate intake is inadequate, concentrations begin to decrease in the fifth month of pregnancy and for several weeks after birth (19). Women who did not take folic acid supplementation during pregnancy, compared to women who did, were at greater risk of fetal growth restriction with a short (less than six months) IPI and, this risk was found to decrease as IPI increased (18). Or interest, a retrospective Canadian study of 46,243 women found an association between IPI (less than six months) and folate-independent anomalies, however not for folate-dependent anomalies such as neural tube defects, cleft lip and palate, and cardiovascular defects (11). In addition, the association between short IPI and anemia was found inconclusive (2).

Implications for WIC Nutrition Service

Findings from a small pilot study found coordination of primary health care and social support services reduced adverse pregnancy outcomes and the average number of pregnancies conceived within 18 months among low-income African American who previously delivered a very low birth weight baby (20). Results from a 2007 U.S. survey found that among women of childbearing age, those aged 18-24 years were the least aware of the need for folic acid prior to pregnancy and least likely to report daily use of supplements containing folic acid. Of equal concern, only 17% of women aged 18-24 years were likely to hear about folic acid from their healthcare provider (21).

Initiations of healthcare referrals for family planning, early prenatal care, and folic acid supplementation have the potential to improve health outcomes for women, infants, and children. Given that half of all pregnancies nationwide are unintended (22), WIC can help to reduce the risk of adverse pregnancy outcomes by:

- Encouraging postpartum women and their partner to meet with their healthcare provider to discuss developing a reproductive plan and birth spacing, as appropriate. http://www.cdc.gov/preconception/documents/rlphealthproviders.pdf

- Encouraging folic acid supplementation. http://www.cdc.gov/features/folicacidbenefits/

- Encouraging healthful eating patterns consistent with the Dietary Guidelines for Americans. http://www.cnpp.usda.gov/DietaryGuidelines

Clarification

Study results for an optimal IPI following a termination or miscarriage have been inconsistent (3, 10, 23, 24). The WHO Technical Consultation on Birth Spacing Report recommended a minimum interval of at least six months between a miscarriage or induced abortion and the next pregnancy. This recommendation was based on a large retrospective cross-sectional study, a review of 258,108 hospital records from several Latin American countries between 1985-2002, that found women whose previous pregnancy resulted in a spontaneous or induced abortion and had an IPI shorter than 6 months had an increased risk for adverse maternal and perinatal outcomes (21). Given several limitations in the study the WHO cautioned against generalizing the results to other regions or even within the Latin American region since service operations and conditions may differ from the study sample (3). However, more recently a review of approximately a million California births found a decreased risk for preterm birth for women with an IPI of less than six months after a terminated pregnancy (10). An overview of the research found that there may be little benefit from delaying pregnancy after an uncomplicated miscarriage, and to that end pregnancy spacing recommendations following a miscarriage should be individually tailored to the person (25).

References

1. Conde-Agudeloa A, Rosas-Bermudez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. JAMA. 2006;295(15):1809-1823.
2. Shachar BZ and Deirdre JL. Interpregnancy Interval and Obstetrical Complications. Obstet Gynecol Surv. 2012;67(9):584-596.
3. World Health Organization. Report of a WHO technical consultation on birth spacing. Geneva, Switzerland, 13-15 June 2005. 2006;1-44.
4. Conde-Agudelo A, Rosas-Bermudez A, Castaño F, Norton MH. Effects of birth spacing on maternal, perinatal, infant, and child health: a systematic review of causal mechanisms. Stud Fam Plann. 2012;43(2):93-114.
5. Conde-Agudelo A, Belizán JM. Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study. BMJ. 2000;321(7271):1255-9.
6. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: a systematic review. Am J Obstet Gynecol. 2007;196(4):297-308.
7. U.S. Department of Health and Human Services. Office of Disease Prevention and Health Promotion. Healthy People 2020. [cited 2015 Feb 1]. Available from: http://www.healthypeople.gov/2020/topicsobjectives2020/objectiveslist.aspx?topicId=13.

8. Shipp TD, Zelop CM, Repke JT, Cohen A, Lieberman E. Interdelivery Interval and Risk of Symptomatic Uterine Rupture. Obstet Gynecol. 2001;97:175-7.
9. Stamillo DM, DeFranco E, Paré E, Odibo AO, Peipert JF, Allsworth JE, et al. Short Interpregnancy Interval: Risk of Uterine Rupture and Complications of Vaginal Birth After Cesarean Delivery. Obstet Bynecol. 2007;110:1075-82.
10. Schachar BZ, Mayo J, Lyell D, Stevenson D, Shaw G. Interpregnancy interval length and risk of preterm birth, a large US study. American Journal of Obstetrics & Gynecology. Wo 14;210(1)Suppl:S373. Abstract. Poster Session V, Number 760.
11. Chen I, Jhangri GS, Chandra S. Relationship between interpregnancy interval and congenital anomalies. Am J Obstet Gynecol. 2014;210(6):564.e1-564.e8.
12. Cheslack-Postava K, Liu K, Bearman PS. Closely Spaced Pregnancies Are Associated With Increased Odds of Autism in California Sibling Births. Pediatrics 2001;127(2):246-53.
13. Gunes N, Surén P, Breshahan M, Hornig M, Lie KK, Lipkin WI, et al. Interpregnancy Interval and Risk of Autism Disorder. Epidemiology. 2013;24:906-912.
14. Bujold E, and Gauthier RJ. Risk of Uterine Rupture Associated with an Interdelivery Interval Between 18 and 24 Months. Obstet Gynecol. 2010;115(5):1003-6/
15. Huang WH, Nakashima DK, Rumney PJ, Keegan KA Jr, Chan K. Interdelivery interval and the success of vaginal birth after cesarean delivery. Obstet Gynecol. 2002;99(1):41-4.
16. The American College of Obstetricians and Gynecologists, Women's Health Care Physicians. ACOG Practice bulletin no. 115: Vaginal birth after previous cesarean delivery. Obstet Gynecol. 2010;116(NO.2 Pt 1):450-63.
17. King JC. The Risk of Maternal Nutritional Depletion and Poor Outcomes Increases in Early or Closely Spaced Pregnancies. J. Nutri. 2003;133:1732S-1736S.
18. Van Eijsden M, Smiths LJ, van der Wal MF, Bonsel GJ. Association between short interpregnancy intervals and term birth weight: the role of folate depletion. Am. J. Clin. Nutr. 2008;88(1):147-53.
19. Smits LJ, Exxed GG. Short interpregnancy intervals and unfavorable pregnancy outcome: role of folate depletion. Lancet. 2001;353:2074-7.
20. Dunlop AL, Dubin C, Raynor BD, Bugg GW Jr, Schmotzer B, Brann AW Jr. Interpregnancy primary care and social support for African-American women at risk for recurrent very-low-birthweight delivery: a pilot evaluation. Matern Child Health J. 2008;12(4):461-8.
21. Centers for Disease Control and Prevention. Use of Supplements Containing Folic Acid Among Women of Childbearing Age – United States, 2007. MMWR. 2008;57(01);5-8.
22. Finer LB, Zolna MR. Shifts in intended and unintended pregnancies in the United States, 2001-2008. Am J Public Health. 2014;104: S43-S48.
23. Conde-Agudeloa A, Belizána JM, Bremanb R, Brockmanb SC, Rosas-Bermudez A. Effect of the interpregnancy interval after an abortion on maternal and perinatal health in Latin America. Int J Gynaecol Obstet. 2005;89(1):S34-S40.
24. Love ER, Bhattacharya, Siladitya; Smith NC, Bhattacharya Sohinee. Effect of interpregnancy interval on outcomes of pregnancy after miscarriage: retrospective analysis of hospital episode statistics in Scotland. BMJ. 2010;341:c3967.
25. Bhattacharya, Sohinee; Smith N. Pregnancy following miscarriage: what is the optimum interpregnancy interval? Women's Health. 2011;7(2):139-141.

Nutrition Risk Criteria **334 LACK OF OR INADEQUATE PRENATAL CARE**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-

Definition

Prenatal care beginning after the 1st trimester (after 13th week), or based on an Inadequate Prenatal Care Index published in a peer review articlesuch as the one by Kessner et al. (4).

First prenatal visit in the third trimester (7-9 months) or:

Weeks Gestation	Number of Prenatal Visits (2)
14-21	0 or unknown
22-29	1 or less
30-31	2 or less
32-33	3 or less
34 or more	4 or less

Required Documentation

Number of prenatal care visits (or no visits).

Justification

Women who do not receive early and adequate prenatal care are more likely to deliver premature, growth retarded, or low birth weight infants (3). The Kessner Index can be used to assess the adequacy of prenatal care for a woman with an uncomplicated pregnancy. Women with medical or obstetric problems, as well as younger adolescents, may need closer management; the frequency of prenatal visits should be determined by the severity of identified problems (1). Several studies have reported significant health and nutrition benefits for pregnant women enrolled in the WIC Program (3).

Clarification

The Centers for Disease Control and Prevention (CDC) defines a trimester as a term of three months in the prenatal gestation period with the specific trimesters defined as follows in weeks:

- First Trimester: 0-13 weeks

- Second Trimester: 14-26 weeks
- Third Trimester: 27-40 weeks

Further, CDC begins the calculation of weeks starting with the first day of the last menstrual period. If that date is not available, CDC estimates that date from the established date of confinement (EDC). This definition is used in interpreting CDC's Prenatal Nutrition Surveillance System data, comprised primarily of data on pregnant women participating in the WIC Program.

References

1. American Academy of Pediatrics, American College of Obstetricians and Gynecologists. Guidelines for Perinatal Care. Washington, D.C.: AAP, ACOG; 1997.
2. Centers for Disease Control and Prevention. Prenatal Nutrition Surveillance System User's Manual. Atlanta: CDC, 1994.
3. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
4. Kessner DM, Singer J, Kalk CE, Schlesinger ER. Infant Death: An analysis by maternal risk and health care. Contrasts in Health Status; Vol. I. Washington, D.C.: Institute of Medicine, National Academy of Sciences; 1973.

Nutrition Risk Criteria **335★ MULTIFETAL GESTATION**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	-

Definition

Pregnant Women: more than one (> 1) fetus in current pregnancy.

Breastfeeding and Non-breastfeeding Women: most recent pregnancy.

Required Documentation

Document source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Multifetal gestations are associated with low birth weight, fetal growth restriction, placental and cord abnormalities, preeclampsia, anemia, shorter gestation and an increased risk of infant mortality. Twin births account for 16% of all low birth weight infants. The risk of pregnancy complications is greater in women carrying twins and increases markedly as the number of fetuses increases.

For twin gestations, the 2009 IOM recommendations provide provisional guidelines: normal weight women should gain 37-54 lbs; overweight women, 31-50 lbs; and obese women, 25-42 lbs (3). There was insufficient information for the IOM committee to develop even provisional guidelines for underweight women with multiple fetuses. A consistent rate of weight gain is advisable. A gain of 1.5 lbs per week during the second and third trimesters has been associated with a reduced risk of preterm and low-birth weight delivery in twin pregnancy (2). In triplet pregnancies the overall gain should be around 50 lbs with a steady rate of gain of approximately 1.5 lbs per week throughout the pregnancy (2). Education by the WIC nutritionist should address a steady rate of weight gain that is higher than for singleton pregnancies.

Pregnant or breastfeeding women with twins have greater requirements for all nutrients than women with only one infant. Postpartum, non-breastfeeding women delivering twins are at greater nutritional risk than similar women delivering only one infant. All three groups of women would benefit greatly from the nutritional supplementation provided by the WIC Program.

References

1. Brown JE and Carlson M. Nutrition and multifetal pregnancy. J Am Diet Assoc. 2000; 100:343-348.
2. Institute of Medicine. WIC nutrition risk criteria: a scientific assessment. National Academy Press, Washington, D,C.; 1996.
3. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines (Prepublication Copy). National Academy Press, Washington, D.C.; 2009. www.nap.edu. Accessed June 2009.

Additional Related References

1. Brown JE, Schloesser PT. Pregnancy weight status, prenatal weight gain,

and the outcome of term twin gestation. Am.J.Obstet.Gynecol. 1990;162:182-6.
2. Suitor CW, editor. Maternal weight gain a report of an expert work group. Arlington, Virginia: National Center for Education in Maternal and Child Health; 1997. Sponsored by Maternal and Child Health Bureau, Health Resources and Services Administration, Public Health Service, U.S. Department of Health and Human Services.
3. Williams RL, Creasy RK, Cunningham GC, Hawes WE, Norris FD, Tashiro M. Fetal growth and perinatal viability in California. Obstet.Gynecol. 1982;59:624-32.
4. Worthington-Roberts, BS. Weight gain patterns in twin pregnancies with desirable outcomes. Clin.Nutr. 1988;7:191-6.

Nutrition Risk Criteria **336** **FETAL GROWTH RESTRICTION**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	PREGNANT WOMEN	1	RD

Definition

Fetal Growth Restriction (FGR) (replaces the term Intrauterine Growth Retardation (IUGR)), may be diagnosed by a physician with serial measurements of fundal height, abdominal girth and can be confirmed with ultrasonography. FGR is usually defined as a fetal weight $< 10^{th}$ percentile for gestational age.

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Fetal Growth Restriction (FGR) usually leads to low birth weight (LBW), which is the strongest possible indicator of perinatal mortality risk. Severely growth restricted infants are at increased risk of fetal and neonatal death, hypoglycemia, polycythemia, cerebral palsy, anemia, bone disease, birth asphyxia, and long term neurocognitive complications. FGR may also lead to increased risk of ischemic heart disease, hypertension, obstructive lung disease, diabetes mellitus, and death from cardiovascular disease in adulthood. FGR may be caused by conditions affecting the fetus such as infections and chromosomal and congenital anomalies. Restricted growth is also associated with maternal height, prepregnancy weight, birth interval, and maternal smoking. WIC's emphasis on preventive strategies to combat smoking, improve nutrition, and increase birth interval, may provide the guidance needed to improve fetal growth.

References

1. Altman DG, Hytten FE. Intrauterine growth retardation: let's be clear about it. Br.J.Obstet.Gynaecol. 1989;96:1127-32.
2. Barros FC, Huttly SR, Victora CG, Kirkwood BR, Vaughan JP. Comparison of the causes and consequences of prematurity and intrauterine growth retardation: a longitudinal study in southern Brazil. Pediatrics 1992;90:238-44.
3. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
4. Institute of Medicine. Nutrition during pregnancy; part I, weight gain and part II, nutrient supplements. National Academy Press, Washington, D.C. 1990.
5. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
6. Kramer MS, Olivier M, McLean FH, Dougherty GE, Willis DM, Usher RH. Determinants of fetal growth and body proportionality. Pediatrics 1990;86:18-26.

7. Kramer MS, Olivier M, McLean FH, Willis DM, Usher RH. Impact of intrauterine growth retardation and body proportionality on fetal and neonatal outcome. Pediatrics 1990;86:707-13.
8. Stein ZA, Susser M. Intrauterine growth retardation: epidemiological issues and public health significance. Semin.Perinatol. 1984;8:5-14.
9. Williams SR. Nutrition and diet therapy. St. Louis, Missouri: Times Mirror/Mosby College Pub, 1989.
10. Worthington-Roberts BS, Williams SR. Nutrition During Pregnancy and Lactation. St. Louis: Mosby, 1989.

Nutrition Risk Criteria **337 HX LARGE FOR GESTATIONAL AGE (LGA) INFANT**

Category, Priority, and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	**RD**
BREASTFEEDING WOMEN	1	**-**
NON-BREASTFEEDING WOMEN	6	**-**

Definition

Pregnant Women: **Any** history of giving birth to an infant weighing greater than or equal to (≥) 9 lbs (4000 g).

Breastfeeding/Non-breastfeeding Women: Most recent pregnancy; or history of giving birth to an infant weighing greater than or equal to (≥) 9 lbs (4000 g).

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. as self-report of diagnosis by physician or written documentation by physician).

Justification

Women with a previous delivery of an infant weighing more than 9 lbs (4000 g) are at an increased risk of giving birth to a large for gestational age infant. Macrosomia may be an indicator of maternal diabetes (current or gestational) or a predictor of future diabetes.

The incidence of maternal, fetal, and neonatal complications is high with neonates weighing 9 lbs (4000 g). Risks for the infant include dystocia, meconium aspiration, clavicular fracture, brachia plexus injury, and asphyxia.

References

1. Boyd, M.E, et al. (1983) Fetal Macrosomia: Prediction, risks, proposed management. *Obstet. Gynecol*. 1983; 61:715-722.
2. Institute of Medicine. (1990) *Nutrition During Pregnancy*. National Academy Press. p. 117, 187.
3. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. Washington (DC): National Academy Press; 1996. p. 117.

Nutrition Risk Criteria **338** **PREGNANT WOMAN CURRENTLY BREASTFEEDING**

	Category	Priority	Referral
Category, Priority and Referral	PREGNANT WOMEN	1	RD

Definition

Breastfeeding woman now pregnant.

Required Documentation

Justification

Breastfeeding during pregnancy can influence the mother's ability to meet the nutrient needs of her growing fetus and nursing baby. Generally, pregnancy hormones cause the expectant mother's milk supply to drastically decline (until after delivery). If the mother conceived while her nursing baby was still solely or predominantly breastfeeding, the baby could fail to receive adequate nutrition. In addition to changes in milk volume and composition, mothers who breastfeed throughout a pregnancy usually report that their nipples, previously accustomed to nursing, become extremely sensitive (presumably due to pregnancy hormones). When women nurse through a pregnancy it is possible that oxytocin released during breastfeeding could trigger uterine contractions and premature labor. When a mother chooses to nurse through a pregnancy, she needs breastfeeding counseling.

Reference

Mohrbacher N, Stock J, La LL, I. The breastfeeding answer book. Schaumburg, Ill: La Leche League International, 1997.

Nutrition Risk Criteria **339** **HISTORY OF BIRTH WITH NUTRITION RELATED CONGENITAL OR BIRTH DEFECT**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

A woman who has given birth to an infant who has a congenital or birth defect linked to inappropriate nutritional intake, e.g., inadequate zinc, folic acid, excess vitamin A.

Pregnant Women: any history of birth with nutrition-related congenital or birth defect
Breastfeeding/Non-breastfeeding: most recent pregnancy

Presence of condition diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The single greatest risk factor for a pregnancy with a neural tube defect is a personal or family history of such a defect. More than 50% of recurrences can be prevented by taking folic acid before conception. Recent studies suggest that intake of folic acid may also be inversely related to the occurrence of cleft lip and palate. The WIC Program provides nutrition education and folic acid-rich foods to women to help prevent future birth defects.

Recurrent birth defects can also be linked to other inappropriate nutritional intake prior to conception or during pregnancy, such as inadequate zinc (LBW) or excess vitamin A (cleft palate or lip). The food package and nutrition education provided to WIC participants help women at risk make food choices that provide appropriate nutrient levels.

Clarification

Self-reporting for “History of…”conditions should be treated in the same manner as self-reporting for current conditions requiring a physicians diagnosis, i.e., the applicant may report to the CPA that she was diagnosed by a physician with a given condition at some point in the past. As with current conditions, self-diagnosis of a past condition should never be confused with self-reporting.

References

1. Federal Register, Part III, DHHS, FDA, 21 CFR Part 101, Food Labeling: Health Claims and Label Statements, Folate and Neural Tube Defects. Proposed and Final Rule. March 5, 1996;61;44:8752-8781.
2. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Diagnosis of nutritional deficiencies or a disease caused by insufficient dietary intake of macro and micronutrients. Diseases include, but are not limited to, Protein Energy Malnutrition, Scurvy, Rickets, Beri Beri, Hypocalcemia, Osteomalacia, Vitamin K Deficiency, Pellagra, Cheilosis, Menkes Disease, Xerophthalmia.

Presence of nutrient deficiency diseases diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The presence of macro- and micro-nutrient deficiencies indicates current nutrition health risks.

Persistent malnutrition may lead to elevated morbidity and mortality rates. Important functional disturbances may occur as a result of single or multiple nutrient deficiencies. Examples include impaired cognitive function, impaired function of the immune system, and impaired function of skeletal muscle. Participation in the WIC Program provides key nutrients and education to help restore nutrition status and promote full rehabilitation of those with an overt nutrient deficiency.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
2. Worthington-Roberts BS, Williams SR. Nutrition throughout the life cycle, 4th edition. Boston: McGraw-Hill, 2001.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Disease(s) or condition(s) that interferes with the intake, digestion, and/or absorption of nutrients. The diseases and/or conditions include, but are not limited to:

- peptic ulcer
- post-bariatric surgery
- short bowel syndrome
- inflammatory bowel disease, including ulcerative colitis or Crohn's disease
- liver disease
- pancreatitis
- gastroesophageal reflux disease (GERD)
- biliary tract diseases

Presence of gastro-intestinal disorders diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician’s orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician.

Justification

Gastrointestinal disorders increase nutrition risk through a number of ways, including impaired food intake, abnormal deglutition, impaired digestion of food in the intestinal lumen, generalized or specific nutrient malabsorption, or excessive gastrointestinal losses of endogenous fluids and nutrients. Frequent loss of nutrients through vomiting, diarrhea, malabsorption, or infections can result in malnourishment and lowered resistance to disease (1, 2).

Gastroesophageal Reflux Disease (GERD)
GERD is irritation and inflammation of the esophagus due to reflux of gastric acid into the esophagus (3). Nutritional care of GERD includes avoiding eating within 3 hours before going to bed; avoiding fatty foods, chocolate, peppermint, and spearmint, which may relax the lower esophageal sphincter; and coffee and alcoholic beverages, which may increase gastric secretion (4). Consumption of these items may need to be limited depending on individual tolerance.

Peptic Ulcer
Peptic ulcer normally involves the gastric and duodenal regions of the gastrointestinal tract (4). Because the primary cause of peptic ulcers is Helicobacter pylori infection, the focus of treatment is the elimination of the bacteria with antibiotic and proton pump inhibitor therapy. Dietary advice for

persons with peptic ulcers is to avoid alcohol, coffee (with and without caffeine), chocolate, and specific spices, such as black pepper (4, 5).

Post-bariatric Surgery

Many types of surgical procedures are used for the intervention of morbid obesity. These procedures promote weight loss by restricting dietary intakes, e.g., adjustable gastric banding (AGB), and/or bypassing some portion of intestine to cause incomplete digestion and/or malabsorption of nutrients, e.g., Roux-y gastric bypass (RYGB). Therefore, the risks for developing nutritional deficiencies after bariatric surgery are greatly increased. Since gastric bypass individuals have both a decreased availability of gastric acid and intrinsic factor, vitamin B_{12} deficiency can develop without supplementation. Taking daily nutritional supplements and eating foods high in vitamins and minerals are important aspects of the nutritional management for the individuals who have had bariatric surgery (6).

Short Bowel Syndrome (SBS)

SBS is the result of extensive small bowel restriction. SBS in infants is mostly the result of small bowel resection for the treatment of congenital anomalies, necrotizing enterocolitis, and congenital vascular. In adults, Crohn's disease, radiation enteritis, mesenteric vascular accidents, trauma, and recurrent intestinal obstruction are the most common conditions treated by small bowel resection and resulting in SBS (4). The loss of a large segment of the small bowel causes malabsorption syndrome. Total parental nutrition usually is started within the first few days after intestinal resection. Gradual supplementation with enteral feeding promotes intestinal adaptation in order to wean from parenteral nutrition therapy. Supplementation with fat soluble vitamins and vitamin B_{12} may be needed (7). The pediatric client's nutritional status must be assessed and growth closely monitored (8).

Inflammatory Bowel Disease (IBD)

Inflammatory bowel disease includes Crohn's disease and ulcerative colitis. Weight loss, growth impairment, and malnutrition are the most prevalent nutritional problems observed in IBD. Nutritional support is essential. Exclusive elemental nutrition has been used in attaining the remission of Crohn's disease. However, symptoms tend to recur promptly after resuming the conventional diet (9).

Liver Disease

Since the liver plays an essential role in the metabolic processes of nutrients, liver disorders have far-reaching effects on nutritional status. Acute liver injury is often associated with anorexia, nausea and vomiting. Therefore, inadequate nutritional intakes are common. Decreased bile salt secretion is associated with the maldigestion and impaired absorption of fat and fat-soluble vitamins. Defects in protein metabolism associated with chronic liver failure include decreased hepatic synthesis of albumin, coagulation factors, urea synthesis and metabolism of aromatic amino acids. For nutritional therapy, an important consideration should be the balance between preventing muscle wasting and promoting liver regeneration without causing hepatic encephalopathy. It is recommended that persons with chronic liver disease consume the same amount of dietary protein as that required by normal individuals (0.74 g/kg) (10).

Pancreatic Disease

In chronic pancreatitis, there is reduced secretion of pancreatic enzymes leading to malabsorption. In severe cases, tissue necrosis can occur. It is suggested that

for patients with pancreatitis, a high carbohydrate, low-fat, low protein diet may be helpful (11).

Biliary Tract Diseases

Common diseases of the biliary tract are:

- Cholelithiasis (gallstones, without infection)
- Choledocholithiasis (gallstone in the bile duct causing obstruction, pain and cramps)
- Cholecystitis (inflammation of gallbladder caused by bile duct obstruction).

Obesity or sever fasting may increase risk for these disorders. Since lipids stimulate gallbladder contraction, a low fat diet with 25% to 30% of total calories as fat is recommended. Greater fat limitation is undesirable as some fat is required for stimulation and drainage of the biliary tract. Supplementation with fat-soluble vitamins may be needed for persons with fat malabsorption or a chronic gall bladder condition (12).

WIC nutritionists can provide counseling to support the medical nutrition therapy given by clinical dietitians, and monitor compliance with therapeutic dietary regimens. They can also review and provide WIC-approved medical foods or formulas prescribed by the health care providers. In certain circumstances, WIC staff may recommend an appropriate medical food or formula to the health care provider. They should also make referrals to an appropriate health care provider for medical nutrition therapy by a clinical dietitian when indicated.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
2. American Dietetic Association, Pediatric Nutrition Practice Group. Pediatric manual of clinical dietetics. Chicago: Pediatric Nutrition Dietetic Practice Group, American Dietetic Association, 1998.
3. Stenson W. The esophagus and stomach. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins 1999. p. 1125-1133.
4. Beyer PL. Medical nutrition therapy for upper gastrointestinal tract disorders. In: Mahan LK, Escott-Slump S, editors. Krause's food nutrition and diet therapy. 11th ed. Philadelphia: Saunders; 2004. p. 688-690.
5. American Dietetic Association. Nutrition Care Manual. Gastrointestinal disease; Peptic ulcers; 2006. http://www.nutritioncaremanual.org. Accessed 1/08.
6. Allied Health Sciences Section Ad Hoc Nutrition committee: Aills L, Blankenship J, Buffington C, Furtado M and Parrot J. Bariatric nutrition: suggestions for the surgical weight loss patient. Review. Surgery for Obesity and Related Diseases 2008 May 17.

7. Scolapio JS, Fleming R. Short Bowel Syndrome. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins; 1999. p. 1135-1140.
8. Farrell M. Gastrointestinal disorders of infancy and childhood (with nutrition support and probiotics) In: Ekvall SW, Ekvall VK. Editors. Pediatric nutrition in chronic diseases and developmental disorders. 2nd ed. Oxford University Press; 2005. p. 248-249.
9. Griffiths A. Inflammatory bowel disease. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkins; 1999. p. 1141-1149.
10. Lieber CS. Nutrition in liver disorders. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams 7 Wilkins; 1999. p. 1177-1189.
11. Raimondo m, Dimagno EP. Nutrition in pancreatic disorders. In: Maurice ES, Olson JA, Shike M, Ross AC, editors. Modern nutrition in health and disease. 9th ed. Lippincott Williams & Wilkinds; 1999. p. 1169-1176.
12. Hasse JM, Matarese JE. Medical nutrition therapy for liver, biliary system and exocrine pancreas disorders. In: Mahan LK, Escott-Stump S, editors. Krause's food nutrition and diet therapy. 11th ed. Philadelphia: Saunders; 2004. p. 758-760.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Diabetes mellitus consists of a group of metabolic diseases characterized by inappropriate hyperglycemia resulting from defects in insulin secretion, insulin action or both (1).

Presence of diabetes mellitus diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Diabetes mellitus may be broadly described as a chronic, systemic disease characterized by:

- Abnormalities in the metabolism of carbohydrates, proteins, fats and insulin; and
- Abnormalities in the structure and function of blood vessels and nerves (2).

The chronic hyperglycemia of diabetes is associated with long-term damage, dysfunction, and failure of various organs, especially the eyes, kidneys, nerves, heart, and blood vessels (1,2) and includes type 1 diabetes mellitus, type 2 diabetes mellitus, and Maturity Onset Diabetes of the Young (MODY). MODY is a series of familial disorders characterized by early onset and mild hyperglycemia. Specific genetic defects have been identified on chromosomes 7, 12, and 20 (2). MODY is often diagnosed before the age of 25 years. It is caused by dominantly inherited defect of insulin secretion. Persons with MODY are often non-obese and without metabolic syndrome (3).

The two major classifications of diabetes are Type 1 Diabetes (beta-cell destruction, usually leading to absolute insulin deficiency) and Type 2 Diabetes (ranging from predominantly insulin resistance with relative insulin deficiency to predominantly an insulin secretory defect with insulin resistance). The Expert Committee (see reference below), working under the sponsorship of the American Diabetes Association, has identified the following as criteria for the diagnosis of diabetes mellitus (1, 2) (see clarification).

Long-term complications of diabetes include retinopathy with potential loss of vision, nephropathy leading to renal failure; peripheral neuropathy with risk of foot ulcers, amputations, and Charcot joints; and, autonomic neuropathy causing gastrointestinal, genitourinary, cardiovascular symptoms and sexual dysfunction. Patients with diabetes have an increased incidence of atherosclerotic cardiovascular, peripheral arterial and cerebrovascular diseases. Hypertension

and abnormalities of lipoprotein metabolism are often found in people with diabetes (1).

WIC nutrition services can reinforce and support the medical and dietary therapies (such as Medical Nutrition Therapy) that participants with diabetes receive from their health care providers (4).

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Clarification

Diabetes mellitus is sometimes described by both patients and health professionals as "a little bit of sugar" or "high sugar." In reality, "sugar" is only one component of the pathology and clinical manifestations of the multifaceted syndrome of diabetes mellitus (2).

Diabetes mellitus is diagnosed by a licensed medical provider using any one of the following three methods:

1. Fasting plasma glucose ≥ 126 mg/dL (7.0 mmol/l). Fasting is defined as no caloric intake for at least 8 hours.
2. Symptoms of hyperglycemia plus casual plasma glucose concentration ≥ 200 mg/dl (11.1 mmol/L).
    - Casual implies any time of day without regard to time since last meal.
    - The classic symptoms of hyperglycemia include polyuria, polydipsia, and unexplained weight loss.
3. Two-hour plasma glucose ≥ 200mg/dL (11.1 mmol/L) during a 75 g oral glucose tolerance test (OGTT) (1).

In the absence of unequivocal hyperglycemia, these criteria should be confirmed by repeat testing on a different day. The third measure (OGTT) is not recommended for routine clinical use.

Reference

1. American Diabetes Association. Diagnosis and classification of diabetes mellitus. Diabetes Care. Jan 2008;31 Suppl 1:S55-60.
2. Franz MT, Ratner RE. Diabetes and Complications. In: Pathophysiology of the diabetes disease state: a Core Curriculum for Diabetes Educators American Association of Diabetes Educators. 5th ed. 2003.
3. Dean L, McEntrye J. The genetic landscape of diabetes. 2004; Bethesda: NCBI, 2004.
4. Http://www.ncbi.nlm.nih.gov/books/bv.fcgi?call=by.view.,showtoc&rid=diabetes.toc&depth=1.
5. American Diabetes Association. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. Diabetes Care. 2006; 29: 2140-2157:S48-S65.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Thyroid Dysfunction	Definition
Hyperthyroidism	Excessive thyroid hormone production (most commonly known as Graves' disease and toxic multinodular goiter).
Hypothyroidism	Low secretion levels of thyroid hormone (can be overt or mild/subclinical). Most commonly seen as chronic autoimmune thyroiditis (Hashimoto's thyroiditis or autoimmune thyroid disease). It can also be caused by severe iodine deficiency.
Congenital Hyperthyroidism	Excessive thyroid hormone levels at birth, either transient (due to maternal Grave's disease) or persistent (due to genetic mutation).
Congenital Hypothyroidism	Infants born with an under active thyroid gland and presumed to have had hypothyroidism in-utero.
Postpartum Thyroiditis	Transient or permanent thyroid dysfunction occurring in the first year after delivery based on an autoimmune inflammation of the thyroid. Frequently, the resolution is spontaneous.

Presence of condition diagnosed, documented, or reported by a physician or someone working under physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The thyroid gland manufactures three thyroid hormones: thyroxine (T_4), triiodothyronine (T_3), and calcitonin. The thyroid hormones regulate how the body gets energy from food (metabolism). Iodine is an essential component of the T_4 and T_3 hormones (1) and must come from the diet. (Note: In nature, iodine does not exist as a free element; rather, it forms compounds such as sodium iodide (2, 3). For more information see Clarification section.) Iodine is available from various foods, and is present naturally in soil and sea water. A dysfunctional thyroid gland can become enlarged (goiter) as a result of an overproduction of thyroid hormones (hyperthyroidism) or conversely, from

insufficient thyroid hormone production (hypothyroidism). Thyroid hormones influence virtually every organ system in the body.

Maternal needs for dietary iodine and thyroid hormone medication (if prescribed) increase during pregnancy as maternal thyroid hormones and iodine are transferred to the fetus along with an increased loss of iodine through the maternal kidneys (3). Concurrently, the fetus is unable to produce thyroid hormones during the first trimester and is entirely dependent on the maternal supply of thyroid hormones. As a result, maternal production of T_4 must increase by at least 50% during pregnancy (4). If the pregnant woman is receiving thyroid hormone therapy, often a 30% - 50% increase in thyroid hormone medication is also needed.

Hyperthyroidism

Hyperthyroidism is a condition in which the thyroid gland is overactive, manufacturing too much thyroid hormone (T_4 and T_3). An excessive consumption or iodine (> 1000 µg/d) may cause fetal and maternal hyperthyroidism (5). In other circumstances, the thyroid might develop nodules which secrete excessive amounts of thyroid hormone regardless of iodine status (5). Enlargement of the thyroid gland (goiter) is a common symptom, as well as weight loss, fatigue, muscle weakness and an irregular heartbeat.

Hyperthyroidism is relatively uncommon in pregnancy (4). However, when it occurs, uncontrolled hyperthyroidism (especially in the second half of pregnancy) may result in infection, miscarriage, preterm delivery, preeclampsia, or congestive heart failure. Fetal complications may include prematurity, small for gestational age, fetal or neonatal thyrotoxicosis, or death (6). Postpartum maternal hyperthyroidism is likely in women with prenatal hyperthyroidism (7).

The primary medical therapy for hyperthyroidism is radioactive iodine therapy which is contraindicated during pregnancy and lactation 97). If hyperthyroidism occurs during this period, low doses of thiomide (antithyroid drug) are given instead.

Hypothyroidism

Hypothyroidism is a condition in which the thyroid gland does not make enough thyroid hormone. Maternal and fetal hypothyroidism may occur when preconception maternal iodine stores are insufficient and there is inadequate maternal iodine intake in early pregnancy. In this instance, the maternal iodine balance may become negative and may never be restored, even with eventual iodine supplementation (4).

Mothers with iodine deficiency during the first half of pregnancy may produce offspring with severe, irreversible brain damage (8). Maternal thyroid deficiency has been associated with neonatal developmental problems which may cause lasting changes in the brain structure and cognitive function.

Uncontrolled hypothyroidism in the second half of pregnancy can cause maternal complications such as anemia, preeclampsia, miscarriage, premature delivery, and postpartum thyroid disease. Fetal or neonatal complications include prematurity, low birth weight, congenital anomalies, poor neuropsychological development, and stillbirth (6).

When iodine nutrition status is adequate, autoimmune thyroid disease (AITD) – also called Hashimoto's thyroiditis – is the most common type of hypothyroidism during pregnancy (4). Pregnant women with AITD are at increased risk of miscarriage and postpartum thyroid disease (including thyroiditis, hyperthyroidism and hypothyroidism). There is an increased risk of permanent and significant impairment in cognitive function for their infants (9).

Congenital Hyperthyroidism and Hypothyroidism

Congenital hyperthyroidism is rare in neonates. Transient congenital hyperthyroidism is caused by maternal Graves' disease. Thyroid stimulating immunoglobulin passes from the mother to the fetus via the placenta and causes thyrotoxicosis in the fetus and subsequently, the neonate. After the baby is born, improvement is rapid if the condition is treated using antithyroid drugs and the hyperthyroidism will subside within several weeks (10). Persistent congenital hyperthyroidism is a familial non-autoimmune disease. It is caused by a genetic mutation resulting in an increase in the constitutive activity of the TSH receptor (11).

Congenital hypothyroidism due to maternal iodine deficiency is a leading cause of preventable mental retardation (10). Over-treatment of thyroid hormone, during pregnancy, as well as prolonged maternal iodine therapy (more than two weeks of therapy or more than 1000 µg/iodine) can also cause congenital hypothyroidism (6). The condition is exacerbated by coexisting selenium and vitamin A deficiencies or iron deficiency (5). Treatment for neonatal hypothyroidism should be started as soon as possible, as every day of delay may result in loss of IQ. Unless treated shortly after birth (within the first 18 days of life), the resulting mental retardation will be irreversible (10).

Postpartum Thyroiditis

Postpartum thyroiditis, an autoimmune inflammation of the thyroid, occurs within the first year after delivery or sometimes after termination of pregnancy. It can be a transient thyroid dysfunction with a brief thyrotoxic phase followed by hypothyroidism, usually with a spontaneous resolution (10). Smoking is a significant precipitating factor in the onset of postpartum thyroiditis (9). Women with a past history of postpartum thyroiditis have a risk of long-term permanent hypothyroidism and recurrence of postpartum thyroiditis in subsequent pregnancies (12). Tests for this condition consist of radioactive products necessitating a temporary cessation of breastfeeding (usually up to 3 days).

Implications for WIC Nutrition Services

Individuals with thyroid disorders can benefit from WIC foods and WIC nutrition services can reinforce and support the medical and dietary therapy prescribed by the participants' health care provider. The following nutrition education messages may be appropriate depending on the type of thyroid disorder:

Encourage iodine sufficiency, unless contraindicated, with an adequate intake of foods high in iodine such as iodized table salt, bread, saltwater fish, kelp, egg yolks (because of iodine supplementation in chicken feed), milk and milk products (because of the treatment of cows with supplemental dietary iodine) (5). It is important to note that the salt used in manufactured foods is not iodized.

Advise women to review the iodine content of their prenatal supplement. It is recommended that all prenatal vitamin-mineral supplements for use

during pregnancy and lactation contain at least150 micrograms of iodine a day (13). Currently, less than 50 percent of prenatal vitamins on the market contain iodine (5, 7).

Promote breastfeeding, as there are no contraindications to breastfeeding and thyroid hormone replacement therapy as long as normal thyroxine levels in the maternal plasma are maintained. Breast milk provides iodine to the infant and is influenced by the dietary intake of the pregnant and lactating mother (14). Hyperthyroidism can develop for the first time during the postpartum period, but the mother's ability to lactate is not affected. However, if a woman with untreated hypothyroidism breastfeeds, her milk supply may be insufficient. In such instances, replacement thyroid hormone therapy is necessary to help increase milk production.

Weight management – hyperthyroidism: The elevated plasma levels of thyroid hormones may cause increased energy expenditure and weight loss along with increased appetite. Following medical treatment, individuals with hyperthyroidism usually regain their typical body weight with a concurrent decrease in appetite (4). Therefore, the monitoring of weight status and dietary adequacy are recommended.

Weight management – hypothyroidism: Many individuals with hypothyroidism experience an increase in weight due to both a decrease in basal metabolic rate and an excessive accumulation of water and salt. Most of the weight gained is due to the excess water and salt retention. After medical treatment, a small amount of weight may be lost, usually less than 10% of body weight (15). Once hypothyroidism has been treated and thyroid hormones are within normal levels, it is less likely that the weight gain is solely due to the thyroid. If an overweight condition persists, weight control therapy may be necessary.

Recommend the cautionary use of soy formula and the avoidance of foods or supplements rich in soy, fiber, or iron when therapeutic thyroid medications are prescribed, since soy, iron, calcium, fiber and phytates may interfere with the absorption of oral thyroid hormone therapy (16, 17).

Discourage smoking as the compound thiocynate found in tobacco smoke inhibits iodine transport (9).

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Iodine (I_2) is an element. In the ambient temperature, it is volatile and forms blue-violet gas. In nature, it does not exist as free element. Instead, it forms compounds, such as sodium **iodide** (NaI), and potassium **iodide** (KI). To prevent iodine deficiency, potassium iodide is added to the salt (most commonly to table salt) to form iodized salt (2, 3).

References

1. National Academy of Sciences. Institute of Medicine. Food and Nutrition Board. Dietary reference intakes: The essential guide to nutrient requirements. 2001.
2. Los Alamos National Labs Chemistry Division Periodic Table. http://periodic.lanl.gov/elements/53.html. Accessed December 2009.
3. WebElements: The periodic table on the web. http://www.webelements.com/iodine/. Accessed December 2009.
4. Smallridge RD, Glinore D, Hollowell JG, Brent G. Thyroid function inside and outside of pregnancy: What do we know and what don't we know? Thyroid. 2005; 15 (1): 54-59.
5. Lee SL, Ananthakrishnan S, Pearce EN. Iodine deficiency. [updated July27, 2006]. Available from: http://www.emedicine.com/med/topic1187.htm.
6. Nguyen PH. Autoimmune thyroid disease and pregnancy. [updated December 21, 2004; cited 2007 Sept 7]. Available from: http://www.emedicine.com.
7. American Association of Clinical Endocrinologists (AACE) Thyroid Task Force. AACE Medical guidelines for clinical practice for the evaluation and treatment of hyperthyroidism and hypothyroidism. 2006 Amended Version. Endocrine Practice 2002; 8 (6): 457-469.
8. LaFranchi SH, Haddow JE, Hollowell JG. Is thyroid inadequacy during gestation a risk factor for adverse pregnancy and developmental outcomes? Thyroid 2005; 15 (1): 60-71.
9. Muller AF, Drexhage HA, Berghout A. Postpartum thyroiditis and autoimmune thyroiditis in women of childbearing age: Recent insights and consequences for antenatal and postnatal care. Endocrine Reviews. 2001 Oct; 22 (5): 605-630.
10. Association for Clinical Biochemistry, British Thyroid Association, British Thyroid Foundation. UK guidelines for the use of thyroid function tests. 2006 July; 1-86.
11. Polak M, Legac I, Vuillard E, Guibourdenche J, Castanet M, Luton D. Congenital hyperthyroidism: The fetus as a patient. Horm Res. 2006; 65: 235-242 (DOI: 10.1159/000092454)
12. O'Malley B, Hickey H, Nevens E. Thyroid dysfunction – weight problems and the psyche: The patients' perspective. J Hum Nutr Dietet. 2000 May; 13: 243-248.
13. American Thyroid Association. American Thyroid Association statement on early maternal thyroidal insufficiency: Recognition, clinical management and research directions. Consensus statement #2. Thyroid. 2005; 15 (1): 77-79.
14. Riordan J. Breastfeeding and human lactation, 3rd ed. Jones & Bartlett Publishers, Inc., 2005; p. 157, 461.
15. American Thyroid Association. Thyroid and weight, 2005. Available from: http://www.thyroid.org
16. American Academy of Pediatrics. American Thyroid Association, Lawson Wilkins Endocrine Society. Update of newborn screening and therapy for congenital hypothyroidism. Pediatrics. 2006; Jun; 117 (6): 2290-2303.
17. American Academy of Pediatrics Committee on Nutrition. Use of Soy protein-based formulas in infant feeding. Pediatrics. 2008; 121 (5): 1062-1068 (doi: 10.1542/peds.2008-0564).

Additional Reference

Hashimoto's Thyroiditis online reference:
http://www.medicinenet.com/hashimotos_thyroiditis/article.htm

Nutrition Risk Criteria **345 HYPERTENSION AND PREHYPERTENSION (INCLUDES CHRONIC AND PREGNANCY INDUCED)**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Presence of hypertension or prehypertension diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Hypertension, commonly referred to as high blood pressure, is defined as persistently high arterial blood pressure with systolic blood pressure above 140 mm Hg or diastolic blood pressure above 90 mm Hg (1). People with high blood pressure can be asymptomatic for years (2). Untreated hypertension leads to many degenerative diseases, including congestive heart failure, end-stage renal disease, and peripheral vascular disease.

There is a large segment of the population that falls under the classification of prehypertension, with blood pressure readings between 130/80 to 139/89 mm Hg (3). People with prehypertension are twice as likely to develop hypertension (3).

There is no cure for hypertension (2); however, lifestyle modifications can prevent high blood pressure and are critical in the management of hypertension and prehypertension (3).

Risk factors for hypertension include (4):

- Age (increases with age)
- Race/ethnicity (occurs more often and earlier in African Americans)
- Overweight and obesity
- Male gender
- Unhealthy nutrient consumption and lifestyle habits (e.g. high sodium intake, excessive alcohol consumption, low potassium intake, physical inactivity, and smoking)
- Family history
- Chronic stress

Management of hypertension includes lifestyle modifications and medication. In prehypertensive individuals, implementing lifestyle changes can prevent or delay the onset of hypertension (3, 5). In hypertensive individuals, dietary intervention is not only effective in reducing blood pressure but also in delaying drug treatment (6).

Lifestyle changes to manage hypertension and prehypertension include:

- Consuming a diet consistent with the Dietary Guidelines for Americans or following the DASH (Dietary Approaches to Stop Hypertension) eating plan, if recommended by a physician
- Limiting dietary sodium
- Engaging in regular physical activity
- Achieving and maintaining a healthy weight
- Smoking cessation

The WIC Program provides fruits, vegetables, low fat milk and cheese, which are important components of the DASH eating plan. WIC nutritionists provide nutrition education counseling to reduce sodium intakes, achieve/maintain proper weight status, promote physical activity, and make referrals to smoking cessation programs, which are the lifestyle interventions critical to the management of hypertension/prehypertension.

Pregnant Women: Hypertension is the most common medical complication of pregnancy, occurring in 7% of all pregnancies. Hypertension during pregnancy may lead to low birth weight, fetal growth restriction, and premature delivery, as well as maternal, fetal and neonatal morbidity (7). Hypertensive disorders of pregnancy are categorized as (8, 9):

- Chronic Hypertension: Hypertension that was present before pregnancy. It increases perinatal mortality and morbidity through an increased risk of SGA (small for gestational age) infants. Women with chronic hypertension are at risk for complications of pregnancy such as preeclampsia. There is a 25% risk of superimposed preeclampsia and an increased risk for preterm delivery, fetal growth restriction, congestive heart failure and renal failure.
- Preeclampsia: A pregnancy-specific syndrome observed after the 20th week of pregnancy with elevated blood pressure accompanied by significant proteinuria.
- Eclampsia: The occurrence of seizures, in a woman with preeclampsia, that cannot be attributed to other causes.
- Preeclampsia superimposed upon chronic hypertension: Preeclampsia occurring in a woman with chronic hypertension. It is the major leading factor of maternal and infant mortality and morbidity.
- Gestational Hypertension: Blood pressure elevation detected for the first time after midpregnancy without proteinuria. It presents minimal risks to mother and baby, when it does not progress to preeclampsia.

The term “pregnancy-induced hypertension” includes gestational hypertension, preeclampsia and eclampsia. For more information about preeclampsia, please see Nutrition Risk Criteria 304, History of Preeclampsia.

The following conditions are associated with an increased incidence of pregnancy-induced hypertension (4):

- Inadequate diet
- Nutritional deficiencies, including low protein, essential fatty acid, or magnesium intake
- Inadequate calcium intake in early pregnancy (7)
- Obesity
- Primigravidity
- Age (pregnancy before age 20 or after age 40)
- Multi-fetal gestation

- Genetic disease factors
- Familial predisposition

The impact of hypertension continues after delivery. Special consideration must be given to lactating women with high blood pressure, especially if their care plan includes medication. It is important that the hypertensive lactating woman inform her physician of her breastfeeding status if she is also taking medication to determine whether they pose any risks to the infant. However, hypertension is not a contraindication for lactation. Lactation, as suggested in research, is thought to present some therapeutic advantages in the management of the disease in women (10, 11, 12).

Children: Hypertension during childhood is age-specific, and is defined as blood pressure readings greater than the 95th percentile for age, gender, and height on at least three separate occasions. Blood pressure reading between the 90th and 95th percentile is considered prehypertension (13). Children with high blood pressure are more likely to become hypertensive adults (15). Therefore, they should have their blood pressure checked regularly beginning at the age of three (14, 15).

Epidemiologic data suggests an association between childhood obesity and high blood pressure (16). Blood pressure and overweight status have been suggested as criteria to identify hypertensive children. Weight control decreases blood pressure, sensitivity to salt and other cardiovascular risk factors (13).

Nutrition-related prevention efforts in overweight hypertensive children should aim at achieving a moderate weight loss or preventing further weight gain. Additionally, a decrease in time spent in sedentary activities with subsequent increase in physical activity should be emphasized.

Dietary changes conductive to weight management in children include:

- Portion control
- Decreased consumption of sugar-containing beverages and energy-dense snacks
- Increased consumption of fresh fruits and vegetables
- Regular meals, especially breakfast

The WIC Program provides nutritious supplemental foods and nutrition education compatible with changes needed to promote a healthy weight and decrease the impact of hypertension in children.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis (“My doctor says that I have/my son or daughter has…”) should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. American Dietetic Association. Nutrition Care Manual. Hypertension: 2006. http://www.nutritioncaremanual.org. Accessed May 2009.
2. Krummel DA. Medical nutrition therapy in hypertension. In: Manhan LK, Escott-Stum[S, editors. Krause's food nutrition and diet therapy. 11th ed. Philadelphia Saunders; 2004, p. 901-902.
3. Chobian, AV, Blakris, GL, Black, HR, et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure, the JNC 7 report, JAMA, 2003. Downloaded from www.jama.com at the University of Florida on May/2009.
4. National Heart Lung and Blood Institute. www.nhlbi.nih.gov. Accessed May 2009.
5. Chobian, AV. Prehypertension Revisited. Hypertension 2006. Downloaded from hyper.ahajournals.org at Health Science Library on May/2009.
6. Appel LJ, Brands MW, Daniels SR, Karanja N, Elmer PJ, Sacks FM; American Heart Association. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. Hypertension 2006; 47:296-308.
7. Roberts JM, Bodnar LM. Report on the WIC nutrition risk criterion for hypertension in pregnancy. July 2007. Unpublished.
8. National Heart Lung and Blood Institute. Report of the working group on research on hypertension during pregnancy, 2001. www.nhlbi.nih.gov. Accessed May 2009.
9. National Heart, Lung and Blood Institute. Working group report on high blood pressure in pregnancy; 2000 Jul. NIH Publication No. 00-3029.
10. Lawrence RA, Lawrence RM. Breastfeeding a guide for the medical profession. 6th ed. Philadelphia: Elsevier Mosby; 2005. p. 590.
11. American Academy of Pediatrics and the American College of Obstetricians and Gynecologists. Breastfeeding handbook for physicians. 2006. p. 179.
12. Lee, SY, Kim, MT, Jee, SH, Yang, HP. Does long-term lactation protect premenopausal women against hypertension risk? A Korean women's cohort study. Preventive Medicine, 2005. Available online at www.sciencedirect.com. Accessed May 2009.
13. US Department of Health and Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. Revised May 2005.
14. Hassink SG. Pediatric obesity: Prevention, intervention, and treatment strategies for primary care. American Academy of Pediatrics; 2007. p. 179.
15. Luma GB, Spiotta RT. Hypertension in children and adolescents. Am Fam Physician 2006 May 6;73: 1158-68.
16. Committee on Nutrition, American Academy of Pediatrics. Pediatric nutrition handbook. 6th ed. Elk Grove, Ill: American Academy of Pediatrics; 2009.

Nutrition Risk Criteria **346** **RENAL DISEASE**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Any renal disease including pyelonephritis and persistent proteinuria, but excluding urinary tract infections (UTI) involving the bladder. Presence of renal disease diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Renal disease can result in growth failure in children and infants. In pregnant women, fetal growth is often limited and there is a high risk of developing a preeclampsia-like syndrome. Women with chronic renal disease often have proteinuria, with risk of azotemia if protein intake becomes too high.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Reference

Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Nutrition Risk Criteria **347** **CANCER**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

A chronic disease whereby populations of cells have acquired the ability to multiply and spread without the usual biologic restraints. The current condition, or the treatment for the condition, must be severe enough to affect nutritional status.

Presence of cancer diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

An individual's nutritional status at the time of diagnosis of cancer is associated with the outcome of treatment. The type of cancer and stage of disease progression determines the type of medical treatment, and if indicated, nutrition management. Individuals with a diagnosis of cancer are at significant health risk and under specific circumstances may be at increased nutrition risk, depending upon the stage of disease progression or type of ongoing cancer treatment.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Reference

Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Conditions which affect energy requirements and may affect the individual's ability to feed self that alter nutritional status metabolically, mechanically, or both. Includes, but is not limited to:

- epilepsy
- cerebral palsy (CP) and
- neural tube defects (NTD), such as spina bifida or myelomeningocele
- Parkinson's disease
- multiple sclerosis (MS)

Presence of central nervous system disorders diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Epileptics are at nutrition risk due to alterations in nutritional status from prolonged anti-convulsant therapy, inadequate growth, and physical injuries from seizures (1). The ketogenic diet has been used for the treatment of refractory epilepsy in children (2). However, children on a ketogenic diet for six months or more have been observed to have slower gain in weight and height (3,4). Growth monitoring and nutrition counseling to increase energy and protein intakes while maintaining the ketogenic status are recommended (4). In some cases, formula specifically prepared for children on a ketogenic diet is necessary. Women on antiepileptic drugs (AEDs) present a special challenge. Most AEDs have been associated with the risk of neural tube defects on the developing fetus. Although it is unclear whether folic acid supplementation protects against the embryotoxic and teratogenic effects of AEDs, folic acid is recommended for women with epilepsy as it is for other women of childbearing age (5-7).

Oral motor dysfunction is associated with infants and children with cerebral palsy (CP). These infants and children often have poor growth due to eating impairment, such as difficulty in spoon feeding, biting, chewing, sucking, drinking from a cup and swallowing. Rejection of solid foods, choking, coughing, and spillage during eating are common among these children (8,9). Growth monitoring and nutrition counseling to modify food consistency and increase energy and nutrient intakes are recommended. Some children may require tube feeding and referral to feeding clinics, where available.
Limited mobility or paralysis, hydrocephalus, limited feeding skills, and genitourinary problems, put children with neural tube defects (NTDs) at

increased risk of abnormal growth and development. Ambulatory disability, atrophy of the lower extremities, and short stature place NTDs affected children at high risk for increased body mass index (10). Growth monitoring and nutrition counseling for appropriate feeding practices are suggested.

In some cases, participants with Parkinson's disease require protein redistribution diets to increase the efficacy of the medication used to treat the disease (11). Participants treated with levodopa-carbidopa may also need to increase the intake of B vitamins (12). Participants with Parkinson's disease will benefit from nutrition education/counseling on dietary protein modification, which emphasizes adequate nutrition and meeting minimum protein requirements. Additionally, since people with Parkinson's often experience unintended weight loss (13), it is important to monitor for adequate maternal weight gain. Individuals with multiple sclerosis (MS) may experience difficulties with chewing and swallowing that require changes in food texture in order to achieve a nutritionally adequate diet (14). Obesity and malnutrition are frequent nutrition problems observed in individuals with MS. Immobility and the use of steroids and anti-depressants are contributing factors for obesity. Dysphagia, adynamia, and drug therapy potentially contribute to malnutrition. Both obesity and malnutrition have detrimental effects on the course of the disease. Adequate intakes of polyunsaturated fatty acids, vitamin D, vitamin B_{12} and a diet low in animal fat have been suggested to have beneficial effects in relapsing-remitting MS (15-17). Breastfeeding advice to mothers with MS has been controversial. However, there is no evidence to indicate that breastfeeding has any deleterious effect on women with MS. In fact, breastfeeding should be encouraged for the health benefits to the infant (18). In addition, mothers who choose to breastfeed should receive the necessary support to enhance breastfeeding duration. As a public health nutrition program, WIC plays a key role in health promotion and disease prevention. As such, the nutrition intervention for participants with medical conditions should focus on supporting, to the extent possible, the medical treatment and/or medical/nutrition therapy a participant may be receiving. Such support may include: investigating potential drug-nutrient interactions; inquiring about the participant's understanding of a prescribed special diet; encouraging the participant to keep medical appointments; tailoring the food package to accommodate the medical condition; and referring the participant to other health and social services.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Institute of Medicine. Food and Nutrition Board. WIC nutrition risk criteria: A scientific assessment. Washington, DC: National Academy Press; 1996.
2. Nelson JK, Mayo C. Mayo clinic diet manual a handbook of nutrition practices. St. Louis: Mosby; 1994.
3. Peterson SJ, Tangney CC, Pimentel-Zablah EM, Hjelmgren B, Booth F, Berry-Kravis E. Changes in growth and seizure reduction in children on the ketogenic diet as a treatment for intractable epilepsy. JADA. 2005 May;105(5):718-724.
4. Santoro KB, O'Flaherty T. Children and the ketogenic diet. JADA. 2005 May;105(5):725-726.
5. Yerby MS. Management issues for women with epilepsy: neural tube defects and folic acid supplementation. Neurology. 2003 Sep;1:61 (6 Suppl 2): S23-6.
6. Champel V, Radal M, Moulin-Vallez M, Jonville-Bera AP, Autret-Leca E. Should folic acid be given to women treated with valproic acid and/or carbamazepine? Folic acid and pregnancy in epilepsy. (Abstract) Rev Neurol (Paris). 1999 Mar;155(3): 220-4.
7. Yerby MS. Clinical care of pregnant women with epilepsy: neural tube defects and folic acid supplementation. Epilepsia. 2003;44 Suppl 3:33-40.
8. Fung EB, Samson-Fang L, Stallings VA, Conaway M, Liptak G, Henderson RC, Worley G, O'Donnell M, Calvert R, Rosenbaum P, Chumlea W, Stevenson RD. Feeding dysfunction is associated with poor growth and health status in children with cerebral palsy. JADA. 2002; 102(3):361-373.
9. Yilmaz S, Basar P, Gisel EG. Assessment of feeding performance in patients with cerebral palsy. Int J Rehabil Res. 2004 Dec; 27(4):325-329.
10. Ekvall SW. Pediatric nutrition in chronic diseases and developmental disorders: prevention, assessment, and treatment. New York Oxford University Press; 1993.
11. Karstaedt PJ, Pincus JH. Protein redistribution diet remains effective in patients with fluctuating parkinsonism. Arch Neurol. 1992 Feb;49(2):149-151.
12. Valkovic P, Benetin J, Blazicek P, Valkovicova L, Gmitterova K, Kukumberg P. Reduced plasma homocysteine levels in levodopa/entacapone treated Parkinson patients. Parkinsonism Relat Disord. 2005 Jun;11(4):253-6. Epub 2005 Apr 20.
13. Chen H, Zhang SM, Heman MA, Willett WC, Ascherio A. Weight loss in Parkinson's disease. Ann Neurol. 2003 May;53(5):676-9.
14. Schapiro R. Managing the symptoms of multiple sclerosis. 4th ed. New York: Demos Medical Publishing; 2003. Ch.13 Swallowing Difficulties.
15. Payne A. Nutrition and diet in the clinical management of multiple sclerosis. J Hum Nutr Dietet. 2001;14:349-357.
16. Schwarz S, Leweling H. Multiple sclerosis and nutrition. Multiple Sclerosis. 2005;11:24-32.
17. Mark BL, Carson JS. Vitamin D and autoimmune Disease-Implications for practice from the multiple sclerosis literature. JADA. 2006 Mar; 106(3): 418-424.
18. Gulick EE, Johnson S. Infant health of mothers with multiple sclerosis. West J Nurs Res. 2004 Oct; 26(6): 632-49.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Hereditary or congenital condition at birth that causes physical or metabolic abnormality. The current condition must alter nutrition status metabolically, mechanically, or both. May include, but is not limited to, cleft lip or palate, Down's syndrome, thalassemia major, sickle cell anemia (not sickle cell trait), and muscular dystrophy.

Presence of genetic and congenital disorders diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

For women, infants, and children with these disorders, special attention to nutrition may be required to achieve adequate growth and development and/or to maintain health.

Severe cleft lip and palate anomalies commonly cause difficulty with chewing, sucking and swallowing, even after extensive repair efforts (5). Surgery is required for many gastrointestinal congenital anomalies. (Examples are trachea-esophageal fistula, esophageal atresia, gastroschisis, omphalocele, diaphragmatic hernia, intestinal atresia, and Hirschsprung's Disease.)

Impaired esophageal atresia and trachea-esophageal fistula can lead to feeding problems during infancy. The metabolic consequences of impaired absorption in short bowel-syndrome, depend on the extent and site of the resection or the loss of competence. Clinical manifestations of short bowel syndrome, include diarrhea, dehydration, edema, general malnutrition, anemia, dermatitis, bleeding tendencies, impaired taste, anorexia, and renal calculi. Total parenteral feedings are frequently necessary initially, followed by gradual and individualized transition to oral feedings. After intestinal resection a period of adaptation by the residual intestine begins and may last as long as 12-18 months (3). Even after oral feedings are stabilized, close follow-up and frequent assessment of the nutritional status of infants with repaired congenital gastro-intestinal anomalies is recommended (5).

Sickle-cell anemia is an inherited disorder in which the person inherits a sickle gene from each parent. Persons with sickle-cell trait carry the sickle gene, but under normal circumstances are completely asymptomatic. Good nutritional status is important to individuals with sickle-cell anemia to help assume adequate growth (which can be compromised) and to help minimize

complications of the disease since virtually every organ of the body can be affected by sickle-cell anemia (i.e., liver, kidneys, gall ladder, and immune system). Special attention should be given to assuring adequate caloric, iron, folate, vitamin E and vitamin C intakes as well as adequate hydration.

Muscular dystrophy is a familial disease characterized by progressive atrophy and wasting of muscles. Changes in functionality and mobility can occur rapidly and as a result children may gain weight quickly (up to 20 lbs in a 6 month period). Early nutrition education that focuses on foods to include in a balanced diet, limiting foods high in simple sugars and fat and increasing fiber intake can be effective in minimizing the deleterious effects of the disease.

References

1. American Dietetic Association, Pediatric Nutrition Practice Group. Pediatric manual of clinical dietetics. Chicago: Pediatric Nutrition Dietetic Practice Group, American Dietetic Association, 1998.
2. Ekvall S. Pediatric nutrition in chronic diseases and developmental disorders prevention, assessment, and treatment. New York: Oxford University Press 1993. p. 289-292.
3. Grand RJ, Sutphen JL, Dietz WH. Pediatric nutrition theory and practice. Boston: Butterworths, 1987.
4. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
5. Ohio Neonatal Nutritionists. Nutritional care for high risk newborns. Philadelphia, PA: G.F. Stickley Publishers, 1985.

Nutrition Risk Criteria **351 INBORN ERRORS OF METABOLISM**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD
CHILDREN	3	RD
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD

Definition

Inherited metabolic disorders caused by a defect in the enzymes or their co-factors that metabolize protein, carbohydrate, or fat.

Inborn errors of metabolism (IEM) generally refer to gene mutations or gene deletions that alter metabolism in the body, including but not limited to:

Inborn Errors of Metabolism*	
Amino Acid Disorders	Urea Cycle Disorders
Organic Acid Metabolism Disorders	Carbohydrate Disorders
Fatty Acid Oxidation Disorders	Peroxisomal Disorders
Lysosomal Storage Diseases	Mitochondrial Disorders
*For information about additional IEM, please see Clarification.	

Presence of condition diagnosed, documented, or reported by a physician or someone working under physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The inheritance of most metabolic disorders is rare. IEM disorders may manifest at any stage of life, from infancy to adulthood. Early identification of IEM correlates with significant reduction in morbidity, mortality, and associated disabilities for those affected (1).

All states screen newborns for IEM, although the type and number of IEM screened for may vary from state to state. Typically, infants are screened for amino acid disorders, urea cycle disorders, organic acid disorders, and fatty acid oxidation defects. A few States are working toward including lysosomal storage diseases and peroxisomal disorders among their newborn screening panels (2).

In most states, treatment of an IEM is referred to a specialized metabolic treatment facility. Please see Clarification for contact information for treatment facilities. IEM treatment is based on symptomatic therapy which may include the following strategies: substrate restriction; stimulation or stabilization of residual enzyme activity; replacement of deficient products; removal of toxic metabolites or blocking their production; and enzyme replacement therapy (3). Avoidance of catabolism is essential at all treatment stages.

Nutrition therapy is integral to the treatment of IEM. Nutrition therapy should both correct the metabolic imbalance and ensure adequate energy, protein, and nutrients for normal growth and development among affected individuals. Continual monitoring of nutrient intake, laboratory values, and the individual's growth are needed for evaluation of the adequacy of the prescribed diet (4). It is important that caregivers of infants and children with IEM ensure that the patient follows the prescribed dietary regimen.

Amino Acid Metabolism Disorders (3)

- Phenylketonuria (included clinically significant hyperphenylalaninemia variants)
- Maple syrup urine disease
- Homocystinuria
- Tyrosinemia

Amino Acid Metabolism Disorders are characterized by the inability to metabolize a certain essential amino acid. The build-up of the amino acid that is not metabolized can be toxic. Treatment of amino acid disorders involves restricting one or more essential amino acids to the minimum required for growth and development and supplying the missing product due to the blocked reaction.

Carbohydrate Disorders (5)

- Galactosemia
- Glycogen storage disease type 1
- Glycogen storage disease type II (see also Pompe disease)
- Glycogen storage disease type III
- Glycogen storage disease type IV (Anderson Disease)
- Glycogen storage disease type V
- Glycogen storage disease type VI
- Hereditary Fructose Intolerance (Fructose 1-phosphate aldolase deficiency, Fructose 1, 6, biphosphatase deficiency, fructose kinase deficiency)

This group of disorders includes an enzyme deficiency or its cofactor that affects the catabolism or anabolism of carbohydrate. Carbohydrate disorders are complex and affect neurological, physical, and nutritional status.

Fatty Acid Oxidation Defects (5)

- Medium-chain acyl-CoA dehydrogenase deficiency
- Long-chain 3-hydroxyacl-CoA dehydrogenase deficiency
- Trifunctional protein deficiency type 1 (LCHAD deficiency)
- Trifunctional protein deficiency type 2 (mitochondrial trifunctional protein deficiency)
- Carnitine uptake defect (primary carnitine deficiency)
- Very long-chain acyl-CoA dehydrogenase deficiency

Fatty acid oxidation defects include any enzyme defect in the process of mitochondrial fatty acid oxidation (FAO) system. The biochemical characteristic of all FAO defects is abnormal low ketone production as a result of the increased energy demands. This results in fasting hypoglycemia with severe acidosis secondary to the abnormal accumulation of intermediate metabolites of FAO, which can result in death.

Organic Acid Disorders (AKA organic aciduria or organic academia) (6)

- Isovaleric acidemia
- 3-Methylcrotonyl-CoA carboxylase deficiency
- Glutaric acidemia type I
- Glutaric acidemia type II
- 3-hydroxy-3-methylglutaryl-coenzyme A lyase deficiency
- Multiple carboxylase deficiency (Biotinidase deficiency, Holocarboxylase synthetase deficiency)
- Methylmalonic acidemia
- Propionic acidemia
- Beta-ketothiolase deficiency

Organic Acid Disorders are characterized by the excretion of non-amino organic acids in the urine. Most of the disorders are caused by a deficient enzyme involving the catabolism of specific amino acid(s). As a result, the non-metabolized substance accumulates due to the blockage of the specific metabolic pathway, which is toxic to certain organs and may also cause damage to the brain (7).

Lysomomal Storage Diseases (6, 8)

- Fabry disease (α-galactosidase A deficiency)
- Gauchers disease (glucocerebrosidase deficiency)
- Pompe disease (glycogen storage disease Type II, or acid α-glucosidase deficiency)

Lysosomal storage diseases are a group of related conditions characterized by increased storage of undigested large molecule in lysosomes. Lysosome is a cellular organelle responsible for intracellular degradation and recycling of macromolecules. Due to a defect in a specific lysosomal enzyme, the macromolecule that normally would be metabolized is not broken down; instead, it accumulates in the lysosomes. This leads to tissue damage, organ failures and premature death. Common clinical features include bone abnormalities, organomegaly, developmental impairment and central, peripheral nervous system disorders.

Mitochondrial Disorders (6, 8)

- Leber hereditary optic neuropathy
- Mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes (MELAS)
- Mitochondrial neurogastrointestinal encephalopathy disease (MNGIE)
- Myoclonic epilepsy with ragged-red fibers (MERRF)
- Neuropathy, ataxia, and retinitis pigmentosa (NARP)
- Pyruvate carboxylase deficiency

Mitochondrial Disorders are caused by the dysfunction of the mitochondrial respiratory chain, or electron transport chain (ETC). Mitochondria play an essential role in energy production. The ETC dysfunction increases free radical production, which causes mitochondrial cellular damage, cell death and tissue necrosis and further worsens ETC dysfunction and thus forms a vicious cycle. The disorders can affect almost all organ systems. However, the organs and

cells that have the highest energy demand, such as the brain and muscles (skeletal and cardiac) are most affected. The clinical features vary greatly among this group of disorders, but most have multiple organ dysfunctions with severe neuropathy and myopathy.

Peroxisomal Disorders (6, 8, 9)

- Zellweger Syndrome Spectrum
- Adrenoleukodystrophy (x-ALD)

There are two types of peroxisomal disorders: single peroxisomal enzyme deficiencies and peroxisomal biogenesis disorders. These disorders cause severe seizures and psychomotor retardation (9). Peroxisomes are small organelles found in cytoplasm of all cells. They carry out oxidative reactions which generate hydrogen peroxides. They also contain catalase (peroxidase), which is important in detoxifying ethanol, formic acid and other toxins. Single peroxisomal enzyme deficiencies are diseases with dysfunction of a specific enzyme, such as acyl coenzyme A oxidase deficiency. Peroxisomal biogenesis disorders are caused by multiple peroxisome enzymes such as Zellweger syndrome and neonatal adrenoleukodystrophy.

Urea Cycle Disorders (6, 5)

- Citrullinemia
- Argininosuccinic aciduria
- Carbamoyl phosphate synthetase deficiency

Urea Cycle Disorders occur when any defect or total absence of any of the enzymes or the cofactors used in the urea cycle results in the accumulation of ammonia in the blood. The urea cycle converts waste nitrogen into urea and excretes it from the kidneys. Since there are no alternate pathways to clear the ammonia, dysfunction of the urea cycle results in neurologic damages.

Implications for WIC Nutrition Services

WIC can provide exempt infant formulas and WIC-eligible medical foods, including those specifically formulated for IEM. Most of the dietary regimens for IEM require a combination of medical food (special formula in most cases) and standard infant formula or prescribed conventional foods. For example, participants with IEM related to essential amino acid metabolism (such as PKU, MSUD), who are not developmentally ready for conventional foods; require both medical food without the offending amino acid(s), and human milk or standard infant formula.

It is recommended that WIC nutritionists collaborate with the clinical dietitians at the metabolic treatment facility, where available, to prescribe WIC food packages (Food Package III) according to the therapeutic diet ordered by the metabolic team, monitor the compliance of the restricted diet, and follow up on the growth and developmental status of the participants with IEM.

Note: Infants with classic galactosemia cannot be breastfed due to lactose in human milk.

Clarification

IEM not listed within this write-up may be found under: http://rarediseases.info.nih.gov/GARD. Please keep in mind these additional resources are not meant for medical advice nor to suggest treatment.

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

The link below lists newborn screening coordinators. The coordinator can direct families to appropriate metabolic treatment facilities based on the newborn screening result: http://genes-r-us.uthscsa.edu/State_contacts.pdf

References

1. Metabolic backgrounder: The Ross metabolic formula system for meeting special nutrition needs. Columbus, OH: Ross Products Division; 2007.
2. Levy PA. Inborn Errors of Metabolism: part 1: Overview. Pediatr Rev. 2009 Apr; 30 (4): 131-7.
3. Wilcken B. An introduction to nutritional treatment in inborn errors of metabolism – different disorders, different approaches. Southeast Asian J Trop med Public Health. 2003; 34 Suppl 3: 198-201.
4. Hendricks KM, Duggan C. Manual of pediatric nutrition, 4th ed. 2005; 626-657.
5. Ekval S, Ekvall, VK, editors. Pediatric nutrition in chronic diseases and developmental disorders: prevention, assessment and treatment. 2nd ed. Oxford University Press; 2005. Part III. Chapters 37-59.
6. GeneReviews are expert-authored, peer-reviewed, current disease descriptions that apply genetic testing to the diagnosis, management, and genetic counseling of patients and families with specific inherited conditions. Available at: http://www.ncbi.nlm.nih.gov/sites/GeneTests/review?db=GeneTests (last accessed 8/5/2010).
7. Stanley CA. Disorders of fatty acid oxidation. In: Fernandes J, et al editors. Inborn metabolic diseases. Berlin Springer; 2000. P. 141-150.
8. Agamanolis D. Inherited metabolic disorders in neuropathology: an illustrated interactive course for medical students and residents. Akron OH: Akron Children's Hospital, Northeastern Ohio University College of Medicine. Available at: http://www.neuropathologyweb.org/chapter10/chapter10aLSDgeneral.html.
9. Van Veldhoven, PP, Leuven KU. Biochemistry and genetic disorders of inherited disorders of peroxisomal fatty acid metabolism. J Lipid Res. 2010 June. Available at: http://www.jlr.org/cgi/rapidpdf/jlr.R005959v1.

Nutrition Risk Criteria **352.1 INFECTIOUS DISEASES – ACUTE (also known as 352a)**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

A disease which is characterized by a single or repeated episode of relatively rapid onset and short duration. Infectious diseases come from bacteria, viruses, parasites, or fungi and spread directly or indirectly from person to person (1). Infectious diseases may also be zoonotic, which are transmitted from animals to humans, or vector-borne, which are transmitted from mosquitoes, ticks, and fleas to humans (1, 2). These diseases and/or conditions include, but are not limited to (an extensive listing of infectious diseases can be found at: http://www.nlm.nih.gov/medlineplus/infections.html):

Most common Acute Infectious Diseases	
Hepatitis A	Listeriosis
Hepatitis E	Pneumonia
Meningitis (Bacterial/Viral)	Bronchitis (3 episodes in last 6 months)
Parasitic Infections	

The infectious disease must be present within the past 6 months, and diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders. See Clarification for more information about self-reporting diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Both chronic and acute infectious diseases can lead to: 1) poor appetite, 2) low nutrient absorption, 3) accelerated nutrient utilization, and/or 4) rapid nutrient loss, depending on the individual's nutritional state before becoming infected and the individual's diet during the improvement period (3). The following information pertains to some of the more prevalent and/or serious acute infectious diseases.

VIRAL HEPATITIS

Hepatitis is inflammation of the liver. It is most often caused by viruses, but can also be caused by excessive alcohol consumption, toxins, and medicines such as acetaminophen, as well as other medical conditions linked to liver inflammation (4). Viral hepatitis is caused by a series of viruses labeled A, B, C, D and E – with A, B, and C being the most common forms in the United States. Viral hepatitis A and E are the only forms that are acute and do not become chronic, whereas B, C, and D can both be acute and chronic in nature (5). (For

more information on chronic infectious diseases see Risk # 352b Infectious Diseases – Chronic). Regardless of the type of hepatitis, infected individuals with signs of the infection will typically experience anorexia, nausea, vomiting, diarrhea, jaundice, epigastria pain, tiredness, and weakness, all of which affect one's diet and health (5). In addition, darker urine and pale stools may be present in infected individuals. It is important to note that viral hepatitis is the leading cause of liver cancer and the most frequent need for liver transplants in the United States (6).

Hepatitis A: Hepatitis A is an acute infection caused by exposure to the Hepatitis A virus. It is transmitted through the fecal-oral route, with transmission most commonly spread through close contact with an infected household member or sexual partner. Outbreaks can also be caused by fecal-contaminated food or water. Because the symptoms of all types of acute hepatitis infections are the same, suspected diagnosis must be confirmed through either positive laboratory testing, or epidemiologic link to a confirmed case. (7)

A large majority of those infected with Hepatitis A are asymptomatic, with 70% showing no clinical signs of infection. Hepatitis A does not progress to a chronic disease, and symptoms typically resolve without treatment in two months, however in 10-15% of cases periodic relapses can occur for up to six months. (8)

The Hepatitis A virus can survive for months outside of the body, therefore proper hygiene and food safety are important preventative measures. However, the most effective method of preventing infection is through vaccination, which has reduced the incidence of Hepatitis A by 95% since its introduction. Emphasis should be placed on preventing an unvaccinated child from close personal contact with someone who is at high risk, or suspected of Hepatitis A infection. (7)

Hepatitis E: Hepatitis E is an acute infection caused by exposure to the Hepatitis E virus. It is transmitted through the fecal-oral route, most commonly through ingestion of contaminated drinking water. However recent cases have been linked to uncooked/undercooked meat and shellfish, indicating the potential for foodborne exposure. While Hepatitis E is believed to be uncommon in the United States, those who frequently travel to developing counties with poor water and environmental sanitation are at risk of becoming infected. Diagnosis for Hepatitis E can be confirmed only by testing for the presence of antibodies to the virus or viral RNA. There are currently no serological tests approved for use in the United States. (9)

Hepatitis E symptoms typically resolve on their own, and there is currently no therapeutic treatment or approved vaccine for the disease. Supportive therapy should be offered and hospitalization recommended for severe cases. The predominant forms of prevention are good sanitation and only relying on clean drinking water when in areas at high risk for infection. (10)

Pregnant women are especially at risk when infected with Hepatitis E. While in general most people will recover completely and the death rate among confirmed cases is about 1%, the mortality rate can reach 10-30% for women in their third trimester. (9)

MENENGITIS

Characterized by an inflammation of the protective membranes known as the meninges, meningitis is typically caused by an infection of the fluid surrounding the brain and the spinal cord. Most commonly meningitis is caused by a bacterial or viral infection, but it can also result as a response to physical injury, cancer, or certain drugs. Due to the severity of meningitis and resulting treatment differing depending on the cause, it is important to correctly diagnose the agent responsible for the disease. (11)

Bacterial Meningitis: While most people with meningitis typically recover, bacterial meningitis is typically severe and can result in serious complications, including brain damage, hearing loss, or learning disabilities. The leading causes of bacterial meningitis in the United States include Haemophilus influenzae, Streptococcus pneumoniae, Listeria monocytogenes, and Neisseria meningitidis. The causes of meningitis vary by age group. In adults, including pregnant women, it is most commonly caused by Streptococcus pneumoniae, Nisseria meningitidis, and Listeria monocytogenes. The cause in newborns is most typically Group B Streptococcus, E. coli, and Listeria. Infants and children most commonly develop meningitis in response to Streptococcus pneumoniae, Neisseria meningitidis, and Haemophilus influenza type b. (12)

In addition, *Cronobactor* may cause severe meningitis in infants. Although *Cronobactor* infection is rare (the Centers for Disease Control and Prevention reports 4-6 infections in infants per year), meningitis due to *Cronobactor* occurs almost exclusively among infants in the first 2 months of life. *Cronobactor* infections have been associated with consumption of reconstituted powdered infant formula. In several outbreak investigations, *Cronobactor* has been found in powdered infant formula that had been contaminated in the factory. In other cases, the powdered infant formula might have been contaminated with *Cronobactor* after it was opened at home or elsewhere. It is recommended that manufacturesr's preparation instructions be adhered to in order to prevent *Cronobactor* infection in infants consuming reconstituted powdered infant formula. (13)

Risk factors for bacterial meningitis include, but are not limited to, age, with infants at higher risk than other age groups; congregate living settings, with groups such as military personnel and college students at increased risk; medical conditions that weaken the immune system; and travel to the meningitis belt in sub-Saharan Africa. Transmission from an infected person usually requires prolonged, close, contact. Additionally, healthy people may carry the bacteria in their nose and throat without developing an illness and most healthy people who carry the disease never become sick. Pregnant women infected with any of the bacteria responsible for causing meningitis are capable of passing the bacteria to their baby, putting them at increased risk of developing meningitis. (12)

Meningitis symptoms are characterized by as sudden onset of fever, headache, and stiff neck. Other symptoms are also often present, including nausea, vomiting, sensitivity to light, and confusion. Diagnosis must be confirmed through laboratory testing of the blood or cerebrospinal fluid. Bacterial meningitis is effectively treated with antibiotics, though it is important to begin treatment as early as possible. (12)

The most effective method of preventing meningitis is vaccination. There are currently vaccines available for three types of meningitis causing bacteria – *Neisseria meningitidis* (meningococcus), *Streptococcus pneumonia* (pneumococcus), and *Haemophilus influenza* type b (Hib). Additionally for

individuals in close contact with those with the disease, antibiotics may be recommended as a preventive measure. The risk of meningitis resulting from Listeria can be prevented by properly preparing and refrigerating food as well as avoiding certain foods. Women diagnosed with group B strep are also given antibiotics during labor to prevent transmission to their newborn. (12)

Viral Meningitis: Viral meningitis is the most common type of meningitis and is often less severe than bacterial caused cases. In the United States it is most commonly caused by non-polio enteroviruses, as well as others including the mumps, herpes, measles, influenza, and arboviruses. While few people infected with these viruses develop meningitis, the risk is especially high from summer to fall. Children younger than five and people with weakened immune systems are at higher risk of developing the disease, with infants younger than one month old and people with weakened immune systems more likely to develop severe illness. (14)

Transmission of a virus that can lead to meningitis may occur due to close contact with a person who has viral meningitis, however it is unlikely meningitis will develop. Symptoms in infants include fever, irritability, poor eating, sleepiness or trouble walking, and lethargy. Adults most commonly experience fever, headache, stiff neck, light sensitivity, sleepiness or trouble walking, nausea, vomiting, lack of appetite, and lethargy. As with bacterial meningitis, diagnosis requires lab tests to confirm the illness. (14)

Typically viral meningitis resolves without treatment in 7-10 days. However those with meningitis caused by the herpes virus or influenza may benefit from antiviral medication. While there are no vaccines available for the non-polio enteroviruses that can cause meningitis, the following steps can be taken to reduce the risk of infection:

- Washing hands often with soap and water, especially after changing diapers, using the toilet, or coughing or blowing your nose.
- Avoiding face touching with unwashed hands.
- Avoiding close contact with infected persons.
- Cleaning and disinfecting frequently touched household surfaces.
- Staying home when sick.

Additionally children should be vaccinated against the other viruses that can cause meningitis, including measles, mumps, chickenpox, and influenza. (14)

LISTERIOSIS

Listeriosis is a serious infection caused by the bacteria *Listeia monocytogenes*. It is most commonly transmitted through contaminated food; however it is also naturally present in the soil, water, and animals, including poultry and cattle (15). Listeria is especially dangerous due to its ability to grow in cold temperatures, unlike many other pathogens (16). Common food sources include ready-to-eat deli meats and hot dogs, unpasteurized milk and dairy products, raw sprouts and others. Symptoms include fever, stiff neck, confusion, weakness, vomiting, and diarrhea (17).

Pregnant women and newborns are at exceptionally high risk for listeriosis, with pregnant women 10-20 times as likely as the general population to become infected (18). It can lead to miscarriage, stillbirth, or lifelong health issues for the child (19). Additionally, those with weakened immune systems are also at

heightened risk. Listeriosis is treated with antibiotics and for severe cases referral to a medical facility may be necessary. The best methods of prevention are associated with proper food safety, handling, and storage. Additionally, raw milk and raw dairy products should be avoided. There is currently no vaccine available. (17)

PNEUMONIA

Pneumonia is an infection of the lungs that can cause mild to severe illness. It can be caused by viruses, bacteria, and fungi. In the United States the most common causes of viral and bacterial pneumonia are respiratory synctal virus (RSV) and Streptococcus pneumonia (pneumococcus), respectively, however Human Parainfluenza Viruses are the leading cause of pneumonia in infants and children. Symptoms include fever, muscle aches, fatigue, enlarged lymph nodes in the neck, chest pain, sore throat, coughing, shortness of breath, and rapid breathing. (20)

Children younger than five years of age are considered at especially high risk of pneumonia. Additionally, pneumonia contracted during pregnancy has been associated with increased morbidity and mortality when compared with non-pregnant women. It can lead to negative outcomes including low birth weight, increased risk of pre-term birth, and serious complications for the mother including respiratory failure. Treatment includes administering antimicrobial and antiviral drugs depending on the pathogen responsible for the infection. (21)

Vaccination is an effective way to prevent pneumonia, with several vaccinations available for both bacteria and viruses including pneumococcal, Haemophilus influenza type b (Hib), pertussis (whopping cough), varicella (chickenpox), measles, and influenza vaccines. Good hygiene is also another effective method of prevention, including regular hand-washing and disinfecting frequently touched surfaces. (20)

BRONCHITIS

Acute bronchitis is diagnosed by a healthcare provider based on the signs and symptoms present in the patient. It is a condition that occurs when the airways in the lungs swell and produce mucus, resulting in a cough. Bronchitis typically occurs after a chest cold and is usually caused by a virus, with the most common being: Respiratory syncytial virus (RSV), Adenovirus, Influenza viruses, and parainfluenza. Symptoms include, but are not limited to coughing that produces mucus; soreness in the chest; fatigue; headache; body aches; fever; and sore throat. Most symptoms of acute bronchitis resolve on their own after two weeks, but the cough may last up to eight weeks in some cases. In severe cases, such as a fever above 100.4 degrees Fahrenheit, patients should seek assistance from a health care provider. (22)

Since bronchitis is almost never caused by bacteria, antibiotics are not needed or recommended. Furthermore, antibiotic treatment may cause harm in both children and adults (20). The best course of action is to provide symptom relief through rest, over-the-counter medicines, and other self-care methods. It is important to use pain relievers appropriate for the age of the child, and only acetaminophen for babies six months of age and younger (23). Bronchitis may be prevented by avoiding smoking, practicing good hygiene, and remaining current on all immunizations (22).

PARASITIC INFECTIONS

Parasites are organisms that live on or in a host organism and survive by getting their food at the detriment of the host. Pregnant women and children are most at risk from certain types of parasites including Toxoplasma gondii – found in uncooked meat; Giardia intestinalis; Cryptosporidium; lice; and pinworms (24). Toxoplasmosis, caused by Toxoplasma gondii, is considered to be the leading cause of death attributed to foodborne illness in the United States (25). To reduce the risk of parasitic infection, prevention includes good food safety and general good hygiene. Additionally environmental risk can be reduced by wearing gloves when coming into contact with soil, covering sandboxes, and teaching children the importance of hand washing 926).

Most healthy people will recover from parasites without treatment. However for pregnant women, newborns, and infants with toxoplasmosis, treatment can be administered as a combination of drugs such as pyrimethamine and sulfadiazine, plus folinic acid (27). This treatment will reduce the parasitic burden, but will not eliminate it completely as parasites can remain in tissues, which makes it hard for the medication to reach them. Lice and other dermal parasites can be treated with topical drugs, such as medicated shampoo (24).

Implications for WIC Nutrition Services

WIC can improve the management of acute infectious diseases through WIC foods, nutrition education, counseling, and referrals to community resources. The table below provides additional WIC nutrition services recommendations specific to the disease state that can help improve the health outcomes of participants with acute infectious diseases:

WIC Nutrition Services Recommendations for Acute Infectious Diseases (9, 10)	
All Types of Infections	• Encourage sufficient calorie intake to ameliorate accelerated nutrient utilization. • Recommend the *Dietary Guidelines* to ensure healthy eating patterns. • Provide suggestions to address poor appetite. • Provide education on safe food handling and storage practices.
All Types of Hepatitis	• Recommend testing to pregnant women and high risk individuals. • Encourage abstinence from alcohol. • Provide information on high calorie, high protein and moderate fat diets. • Recommend high calorie consumption at breakfast as nausea is less common in the morning. • Recommend, in consultation with health care provider, consumption of high calorie and protein liquid formula between meals to boost calorie intake. • Encourage a bland diet with extra fluids depending on the severity of nausea and vomiting.
Hepatitis A	• Encourage the Hepatitis A vaccine for all children, previously unvaccinated adolescents through the age of 18, and high-risk adults. • Promote breastfeeding as being safe, but to avoid breastfeeding when nipples are cracked and bleeding – at

	which time, mothers should pump and discard milk to maintain supply. • Discourage the practice of pre-chewing food for infants, as blood may be present.
Hepatitis E	• Avoid contaminated water.
Meningitis	• Encourage vaccinations for both bacteria and viruses known to cause meningitis. • Provide education on proper food handling and storage practices. • Recommend use of manufacturer's instruction for the preparation of infant formula. • Provide education on good hygiene practices.
Listeriosis	• Recommend alternatives to raw milk and dairy products. • Emphasize importance of safe food handling, preparation and storage practices.
Pneumonia	• Recommend referral to a healthcare provider to administer appropriate antimicrobial or antiviral treatment.
Bronchitis	• Provide education on symptom relief and proper pain-medication practices for children. • Recommend smoking cessation. • Provide education on good hygiene practices. • Encourage appropriate vaccinations.
Parasitic Infections	• Recommend appropriate measures be taken when coming into contact with potential environmental contaminants, e.g., use of gloves when working with soil and covering sandboxes when not in use. • Provide education on proper food handling and storage practices. • Provide education on good hygiene practices.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. World Health Organization. Health topics: Infectious disease. [cited 2015 May 15]. Available from: http://www.who.int/topics/infectious_diseases/en/ .
2. Centers for Disease Control and Prevention (CDC). Division of Vector-Borne Diseases. [cited 2015 May 15]. Available from: http://www.cdc.gov/ncezid/dvbd/about.html .
3. Friis, H. Micronutrients and infection: an introduction. In: Micronutrients and HIV infection. Boca Raton:CRC Press; 2010. P. 3.
4. Centers for Disease Control and Prevention (CDC). Hepatitis information for the public. [cited 2014 Jan 27]. Available from: http://www.cdc.gov/hepatitis/PublicInfo.htm .
5. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases. Viral hepatitis: A through E and beyond. [cited 2014 Jul 26]. Available from: http://www.nlm.nih.gov/medlineplus/hepatitis.html .
6. Centers for Disease Control and Prevention. Division of Viral Hepatitis and National Center for HIV/AIDS. Viral hepatitis, STD, and TB prevention. [cited 2015 May 1]. Available from: http://www.cdc.gov/hepatitis/ .

7. Centers for Disease Control and Prevention (CDC). Viral Hepatitis – Hepatitis A Information. [cited 2012 Aug 18]. Available from: http://www.cdc.gov/hepatitis/hav/havfaq.htm#general .
8. National Institute of Allergy and Infectious Disease (NIAID). Hepatitis A. [cited 2012 Aug 18]. Available from: http://www.niaid.nih.gov/topics/hepatitis/types/Pages/hepatitis A.aspx .
9. Centers for Disease Control and Prevention (CDC). Hepatitis E FAQs for Health Professionals. [cited 2012 Aug 18]. Available from: http://www.cdc.gov/hepatitis/hev/hevfaq.htm#section1 .
10. National Institute of Allergy and Infectious Disease (NIAID). Hepatitis E. [cited 2012 Aug 18] Available from: http://www.niaid.nih.gov/topics/hepatitis/types/Pages/hepatitisE.aspx .
11. Centers for Disease Control and Prevention (CDC). Meningitis. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/meningitis/index.html .
12. Centers for Disease Control and Prevention (CDC). Bacterial Meningitis. [cited 2014 Feb 12]. Available from: http://www.cdc.gov/meningitis/bacterial.html .
13. Centers for Disease Control and Prevention (CDC). Cronobactor. [cited 2014 Feb 12]. Available from: http://www.cdc.gov/meningitis/viral.html .
14. Centers for Disease Control and Prevention (CDC). Viral Meningitis. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/meningitis/viral.html .
15. Food Safety.gov. Listeria. [cited 2012 Aug 26] Available from: http://www.foodsafety.gov/poisoning/causes/bacteriaviruses/listeria/# .
16. Centers for Disease Control and Prevention (CDC). Listeria: People at Risk. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/listeria/risk.html .
17. Schlech WF, 3rd, Lavigne PM, Bortolussi RA, Allen AC, Haldane EV, Wort AJ, Hightower AW, Johnson SE, King SH, Nicholls ES, Broome CV. Epidemic listeriosis—evidence for transmission by food. The New England journal of medicine. 1983;308(4):203-6.
18. Centers for Disease Control and Prevention. Vital signs: listeria illnesses, deaths, and outbreaks – United States, 2009-2011. Morbidity and Mortality Weekly Report (MMWR). 2013;62(22):448-52.
19. Jackson KA, Iwamoto M, Swerdlow D. Pregnancy-associated listeriosis. Epidemiology and infection. 2010:138(10):1503-9.
20. Centers for Disease Control and Prevention (CDC). Pneumonia. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/pneumonia/ .
21. Goodnight W H, & Soper D E (2005). Pneumonia in pregnancy. *Critical care medicine, 33*(10), S390-S397.
22. Centers for Disease Control and Prevention (CDC). Bronchitis. [cited 2012 Sep 16]. Available from: http://www.cdc.gov/getsmart/community/for-patients/common-illnesses/bronchitis.html .
23. Centers for disease Control and Prevention (CDC). Symptom Relief. [cited 2012 Sep 16]. Available from: http://www.cdc.gov/getsmart.community/for-patients/symptom-relief.html .
24. Centers for Disease Control and Prevention (CDC). Parasites. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/parasites/about.html .
25. Centers for Disease Control and Prevention (CDC). Toxoplasmosis. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/parasites/toxoplasmosis/indeshtml .

26. Centers for Disease Control and Prevention (CDC). Toxoplasmosis – Prevention. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/paratistes/toxoplasmosis/prevent.html .
27. Centers for Disease Control and Prevention (CDC). Toxoplasmosis – Treatment. [cited 2012 Aug 26]. Available from: http://www.cdc.gov/parasties/toxoplasmosis/treatment.html .

352.2 INFECTIOUS DISEASES – CHRONIC (also known as 352b)

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Conditions likely lasting a lifetime and require long-term management of symptoms. Infectious diseases come from bacteria, viruses, parasites, or fungi and spread directly or indirectly, from person to person (1). Infectious diseases may also be zoonotic, which are transmitted from animals to humans, or vector-borne, which are transmitted from mosquitoes, ticks, and fleas to humans (1, 2). These diseases and/or conditions include, but are not limited to (an extensive listing of infectious diseases can be found at : http://www.nlm.nih.gov/medlineplus/infections.html):

Chronic Infectious Diseases	
HIV Human Immunodeficiency Virus	Hepatitis B
AIDS Acquired Immunodeficiency Syndrome	Hepatitis C
Hepatitis D	

Presence of condition diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Both chronic and acute infectious diseases can lead to: 1) poor appetite, 2) low nutrient absorption, 3) accelerated nutrient utilization, and/or 4) rapid nutrient loss, depending on the individual's nutritional state before becoming infected and the individual's diet during the improvement period (3). The following information pertains to some of the more prevalent and/or serious chronic infectious diseases.

Human Immunodeficiency Virus (HIV)/ Acquired Immunodeficiency Syndrome (AIDS)

The Human Immunodeficiency Virus (HIV) is a chronic virus that reduces an individual's ability to fight off infections and diseases (4). HIV destroys white blood cells found in the immune system, also known as CD4 (cluster differentiation) or T cells (T lymphocytes) (5). HIV is transmitted only through blood, semen, pre-seminal fluid, rectal fluids, vaginal fluids, and breast milk from an HIV-infected person (6). HIV can lead to Acquired Immunodeficiency Syndrome (AIDS) if left untreated (4). Individuals who are aware of their HIV status and are undergoing antiretroviral therapy (ART) to stop the replication of the virus, can typically live decades – while those unaware of their status or are not on ART, can usually remain in this stage about ten years before progressing to the AIDS stage. Some individuals may progress to the AIDS stage sooner than 10 years. During the time period a person progresses from HIV to AIDS, the immune system becomes extremely weakened and can no longer protect against other infections or opportunistic illnesses** - which are infections generally not detrimental to healthy individuals, but can be life-threatening in people infected with HIV. A person with AIDS and an opportunistic illness that goes untreated has a life expectancy of approximately one year (4).

Getting tested is the only way individuals know they are infected with HIV. Many people infected with the virus display no symptoms for as long as ten years or more. The Centers for Disease Control and Prevention (CDC) currently estimates that 1 in 6 people in the United States infected with HIV do not know they have the virus and therefore recommends that everyone between the ages of 13-64 get tested at least once as part of a regular heath screening. The CDC further recommends that all pregnant women be tested early in their pregnancy, via an "opt-out" testing measure – which is when pregnant women are told that an HIV test will be included in the standard group of prenatal tests and that they may decline the test. Unless the HIV test is specifically declined, they will be tested for the virus. (7)

An early diagnosis in pregnant women can reduce the transmission of HIV in babies to 2%, if the expectant mother (8):

- Receives Active Antiretroviral Therapy (ART) during pregnancy, labor, and delivery.
- Delivers the baby by cesarean, or C-section.
- Avoids breastfeeding.

There is a 20% chance of transmission if the HIV positive, expectant mother does none of the prevention measures listed above (8). In addition, women living in certain geographic areas or women considered high risk, such as those with sexually transmitted infections, multiple partners, or have substance abuse issues, are encouraged to be retested in the third trimester, preferably when less than 36 weeks pregnant (9).

PrEP (Pre-Exposure Prophylaxis) is a daily pill containing two medicines (tenofovir and emtricitabine), recommended for HIV negative people who are at substantial risk of becoming infected with HIV. PrEP, when taken consistently, reduces HIV transmission by up to 92%, and is recommended for (10):

- Individuals in an HIV discordant relationship in which one partner is HIV positive and the other partner is HIV negative.
- Heterosexual women who do not regularly use condoms with sex partners of unknown HIV status.

- Women who share injectable drug paraphernalia or were in treatment for injectable drug use in the past six months.

**Extensive listing of opportunistic illness can be found at: http://womenshealth.gov/hiv-aids/opportunitstic-infections-and-other-conditions/ .

HIV/AIDS and Nutrition: Dietary needs for an HIV positive individual are determined by the presence of symptoms (11, 12). **Symptomatic** individuals experiencing unintended weight loss, or wasting, and are dealing with: 1) poor food intake due to medication side effects, sore mouth, or mental health issues; 2) altered metabolism due to disease progression; or 3) nutrient malabsorption caused by gastrointestinal problems resulting from medications or just the presence of the virus. In symptomatic participants, the main goals are to: 1) increase or maintain a normal body weight; 2) retain or increase lean body mass; and 3) ensure adequate intake of macro- and micronutrients. In most cases, these individuals usually require diets higher in protein and potentially a multivitamin, as vitamins A, B_6, C, and E are lower in symptomatic people. In instances when wasting cannot be alleviated through regular dietary means, enteral and parenteral nutrition therapy may be necessary. For **asymptomatic** individuals or those with a stable weight, the goals should focus on adequate intake of nutrients to prevent wasting – and if food intake is low, these individuals could potentially include a multivitamin or mineral supplement to avoid deficiencies (11, 12).

It is important to note that taking large amounts of iron supplements, leading to iron-overload, encourages disease progression from HIV to AIDS,and should be avoided. In addition, Vitamin A and Zinc, in the form of supplements, can have a negative impact on adults living with HIV/AIDS (12). Participants should always consult with their health care providers before taking dietary supplements over the Recommended Dietary Allowance to prevent adverse reactions and interactions with medications used to treat HIV/AIDS. (13)

HIV/AIDS Medication Nutritional Problems: Even though people with HIV are able to manage the disease and live longer with Highly Active Antiretroviral Therapy (HAART), the side effects can have a negative impact on a person's nutritional status. Common side effects include: gastrointestinal problems, lipid disorders, and insulin resistance/glucose intolerance. Participants experiencing these problems should: reduce total fat intake and cholesterol; increase dietary fiber; increase physical activity; reduce alcohol consumption; and reduce the consumption of simple sugars. (11, 12)

HIV/AIDS and Food Safety: Participants living with HIV are more susceptible to contracting a food-borne illness due to weakened immune systems and therefore should be encouraged to: store and prepare foods safely; check expiration dates; and avoid raw or semi raw foods, such as meat, non-pasteurized dairy, and soft cheeses (11, 12). Infants born to HIV positive mothers, regardless of their HIV status, should drink ready-to-feed or liquid concentrate infant formula as powdered infant formula is not sterile and may not be microbiologically safe (14).

HIV/AIDS Care and Support: HIV-affected families often experience a lack of financial and psychological support needed to deal with an HIV/AIDS diagnosis, including the effects of social stigma which negatively impacts their ability to comply with the medical treatment needed to control the disease (15). Further, to fully benefit from current treatment protocols required to manage HIV and reduce the progression to AIDS, infected individuals who know their status, must get

care, stay in care, and adhere to an effective antiretroviral treatment plan known as an HIV/AIDS Care Continuum (16). WIC agencies should proactively refer participants to health care services and various community resources, including other FNS nutrition assistance programs, to improve health outcomes among HIV-infected WIC participants.

Implications for WIC Nutrition Services

WIC can improve the management of chronic infectious diseases through WIC foods, nutrition education, counseling, and referrals to community resources that provide support in the long-term management of chronic infectious diseases.

HIV/AIDS

The table below summarizes the WIC Nutrition Services that can help improve the health and birth outcomes of participants with HIV/AIDS.

Participant Category	WIC Nutrition Services Recommendations for HIV/AIDs
All Categories	**Nutrition and Health Tips to Manage HIV/AIDS Symptoms (12, 17, 18, 19)** • Use MyPlate as the guide for dietary needs. • Consult health care providers when using supplements and herbs to avoid adverse reactions or medication interactions that could reduce effectiveness. • Eat small, frequent meals when gastrointestinal problems are present or persistent. • Eat soft foods with manageable textures at tolerable temperatures when oral lesions and dental problems are present (i.e. mashed potatoes, scrambled/boiled eggs, bananas, non-citrus juices, puddings, custards, milk, cooked vegetables, rice, oatmeal, non-fizzy drinks, cottage cheese, non-spicy foods). • Add canned tuna, beans, cheese, peanut butter, dried milk for inexpensive extra protein. • Add moderate amounts of concentrated sources of calories to diet when heeded (e.g. butter, cream cheese, gravies, whole milk, ice cream). • Consume nutritious, high caloric foods when appetite is normal or has returned. • Drink adequate water to stay hydrated, replace fluid loss from diarrhea and vomiting, and help medications move through the body. • Consume foods high in fiber or fiber supplements to slow digestion if foods are moving too quickly through the body. • Eat yogurt or foods with *Lactobacillus acidophilus* culture to help with bacterial over-growth resulting from prolonged use of antibiotics. • Avoid caffeinated beverages to prevent dehydration. • Avoid or reduce sugar-free foods with sorbitol as diarrhea may be exacerbated. • Consult with health care provider about use of complete oral nutritional supplements to help nutritional status. • Avoid alcohol and illegal drugs for overall good health and to help protect the liver.

	• Use pancreatic enzymes when medically prescribed to help with digestion. • Prepare and store food safely. • Avoid expired and moldy foods or foods with rotten spots. • Participate in weight-bearing exercises to strengthen and maintain bones. • Refer HIV-affected families to other community resources for food, housing, and medical resources to improve compliance with HIV treatment.

Women	• Encourage all women to be tested to prevent mother-to-child HIV transmission through delivery and breastfeeding (7). Women who are considered high risk, such as those with sexually transmitted infections, multiple partners, or have substance abuse issues, are encouraged to be retested during late gestation, preferably before 36 weeks (9). Note: HIV testing is not a standard medical test administered to pregnant women in many states, in addition, pregnant women can opt-out in those states in which HIV testing is part of the standard test. Therefore, WIC can impact the spread of HIV/AIDS by making referrals to participants for early and late gestation testing, given that some populations served by WIC are most at risk for contracting HIV (7).
	• Advise infected pregnant women to consume a diet adequate in nutrients, achieve appropriate weight gain, and discuss taking a multivitamin with their health care provider (11).
	• Educate mothers with HIV/AIDS to avoid breastfeeding. This is especially important for recent immigrants and refugees from developing nations, as the recommendations are different in developing countries (15). In some developing countries, breastfeeding is encouraged due to the lack of available clean water to prepare infant formula and other sanitation problems).
	• More information about **women and HIV** can be found at:
	  ○ http://www.womenshealth.gov/hiv-aids/
	  ○ http://www,cdc.gov/hiv/risk/gender/pregnantwomen/facts/

Infants	• Inform mothers/caregivers that formula feeding is the standard for infants born to HIV positive mothers in the United States as breastfeeding is not recommended – especially to the immigrant and refugee population (13). • Ensure that liquid concentrate, or ready-to-feed infant formula, prescribed with medical documentation, is provided to HIV-exposed infants or babies born to HIV positive mothers, even if the infant has tested negative for HIV. Powdered infant formula is not sterile and therefore may not be microbiologically safe for immune-compromised participants (14). • Discourage giving pre-chewed food, regardless of HIV status, as the individual's HIV status, who is pre-chewing the food is unknown (6). • More information about **infants and HIV** can be found at: ○ http://www.cdc.gov/hiv/risk/gender/pregnantwomen/facts/
Children	• Discourage giving pre-chewed food, regardless of HIV status, as the individual's HIV status, who is pre-chewing the food is unknown (6). • More information about **children and HIV** can be found at:

	○ http://www.cdc.gov/hiv/risk/gender/pregnantwomen/facts/ ○ http://aids.gov/hiv-aids-basics/just-diagnosed-with-hiv-aids/overview/children/

Viral Hepatitis

Hepatitis is inflammation of the liver. It is most often caused by viruses, but can also be caused by excessive alcohol consumption, toxins, and medicines such as acetaminophen, as well as other medical conditions linked to liver inflammation (20). Viral hepatitis is caused by a series of viruses labeled A, B, C, D, and E with A, B, and C being the most common forms in the United States. Viral hepatitis A and E are the only forms that are acute and do not become chronic, whereas B, C, and D can both be acute and chronic in nature (20). Regardless of the type of hepatitis, infected individuals with signs of the infection will typically experience: anorexia, nausea, vomiting, diarrhea, jaundice, epigastria pain, tiredness, and weakness, all of which affect one's diet and health (21). In addition, darker urine and pale stools may be present in infected individuals. It is important to note that viral hepatitis is the leading cause of liver cancer and the most frequent needs for liver transplants in the United States (22).

Hepatitis B: Hepatitis B is both acute and chronic, and is transmitted through contact with hepatitis B virus (HBV) infected blood, sexual intercourse with an infected person, and from mother to child by both vaginal or cesarean section births (20). Those at higher risk of becoming infected with hepatitis B are those: living with a hepatitis B infected person; coming into contact with blood, needles, or body fluids through work; working or living in a prison system; from Asian and Pacific Islands nations; undergoing kidney dialysis; infected with HIV or hepatitis; and who have an immigrant or refugee status (21).

Treatment for Hepatitis B involves the use of interferon and antiviral drugs to interfere with the course of the virus. Early diagnosis and treatment of hepatitis B can help prevent damage to the liver. In addition, the Hepatitis B vaccination can prevent Hepatitis B. (22)

Hepatitis B is not spread through breast milk. Given that Hepatitis B is spread through blood, mothers who breastfeed should care for their nipples to avoid cracking and bleeding. If a mother with Hepatitis B has cracked and bleeding nipples, she should temporarily stop breastfeeding until her nipples heal – but continue to pump and discard pumped milk to maintain her milk supply (23). If a mother with HBV has concerns with providing her milk to her infant or concerns with drug treatment for the HBV, she should consult her physician.

Hepatitis C: Hepatitis C is both acute and chronic; however, most cases are chronic and commonly spread through sharing needles during intravenous drug use (20). It can also spread through sexual intercourse; having a blood transfusion or organ transplant before July 1992; or using the razor, toothbrush, or nail clippers of an infected person. Being infected with a sexually transmitted disease or HIV can increase the chances of becoming infected with Hepatitis C. Getting tattoos and body piercings from unlicensed facilities, in casual settings, or with the use of non-sterile instruments can also transmit Hepatitis C (20).

By the time symptoms appear with hepatitis C, the liver has been damaged, which in most cases can be as long as ten years after being infected. There is no

vaccine for Hepatitis C. Medicines are used to slow or stop the virus from damaging the liver in chronic hepatitis. Severe damage to the liver leading to failure may require a liver transplant. (20)

Infants born to mothers with hepatitis C can become infected; however, breastfeeding is not contraindicated, as Hepatitis C is not transmitted through breast milk, unless the mother's nipples are cracked and bleeding. (See information above in **Hepatitis B** about breastfeeding with cracked and/or bleeding nipples.)

Hepatitis D: Hepatitis D is both acute and chronic. Though not common in the United States, viral hepatitis D can only be contracted when an individual also has hepatitis B (20, 22). The virus is present in blood and other body fluids of infected persons and is most commonly transmitted through: engaging in sexual activity; mother to child during delivery; sharing injection drug paraphernalia, razors, or toothbrushes; or coming in direct contact with the blood of an infected person. Chronic hepatitis D resulting from a super-infection, in which an individual has chronic hepatitis B, can progress to end-stage liver diseases (cirrhosis) or liver cancer. In some patients, interferon may be useful for treating hepatitis D. Although no vaccine exists for Hepatitis D, it can be prevented in persons who do not have Hepatitis B, by getting the Hepatitis B vaccination (20, 22).

Implications for WIC Nutrition Services

WIC can improve the management of chronic infectious diseases through WIC foods, nutrition education, counseling, and referrals to community resources that provide support in the long-term management of chronic infectious diseases.

Hepatitis
The table below summarizes the WIC Nutrition Services recommendations that can help improve the health outcomes of participants with Hepatitis.

Types of Hepatitis	WIC Nutrition Services Recommendations for Chronic Hepatitis (24, 25)
All Types	• Recommend testing to pregnant women and high risk individuals. • Encourage abstinence from alcohol. • Provide information on high calorie, high protein and moderate fat diets. • Recommend high calorie consumption at breakfast to mitigate nausea. (Typically nausea is less common in the morning.) • Recommend, in consultation with health care provider, consumption of high calorie and protein liquid formula between meals to boost calorie intake. • Encourage a bland diet with extra fluids depending on the severity of nausea and vomiting.
Hepatitis B	• Encourage the Hepatitis B vaccine for all newborns, previously unvaccinated adolescents through the age of 18, and high-risk adults. • Promote breastfeeding as being safe, but to avoid breastfeeding when nipples are cracked and bleeding – at which time, mothers should pump and discard milk to maintain supply.

	• Discourage the practice of pre-chewing food for infants, as blood may be present.
Hepatitis C	• Promote breastfeeding as being safe, but to avoid breastfeeding when nipples are cracked and bleeding – at which time, mothers should pump and discard milk to maintain supply.
Hepatitis D	• Recommend Hepatitis B vaccine.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. World Health Organization. Health topics: Infectious disease. [cited 2015 May 15]. Available from: http://www.who.int/topics/infectious_diseases/en/ .
2. Centers for Disease Control and Prevention (CDC). Division of Vector-Borne Diseases. [cited 2015 May 15]. Available from: http://www.cdc.gov/ncezid/dvbd/about.html .
3. Friis, H. Micronutrients and infection: an introduction. In: Micronutrients and HIV infection. Boca Raton:CRC Press; 2010. P. 3.
4. Centers for Disease Control and Prevention (CDC). HIV/AIDS.: about HIV/AIDS. [cited 2015 May 15]. Available from: http://www.cdc.gov/hiv/basics/whatishiv.html .
5. AIDS.gov. CD4 count. [cited 2015 May 15]. Available from: https://www.aids.gov/hiv-aids-basics/just-diagnosed-with-hiv-aids/understand-your-test-results/cd4-count/ .
6. Centers for Disease Control and Prevention (CDC). HIV/AIDS transmission. [cited 2015 May 15]. Available from: http://www.cdc.gov/hiv/basics/transmission.html .
7. Centers for Disease Control and Prevention (CDC). HIV/AIDS: testing. [cited 2015 May 15]. Available from: http://www.cdc.gov/hiv/basics/testing.html .
8. Centers for Disease Control and Prevention (CDC). HIV/AIDS: prevention. [cited 2015 May 15]. Available from: http://www.cdc.gov/hiv/basics/prevention.html .
9. Branson BM, Handsfield HH, Lampe MA, Janssen RS, Taylor AW, Lyss SB, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. MMWR Recommendations and Reports 2006;55(RR-14):1-17. [cited 2015 May 15]. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5514a1.htm .
10. Centers for Disease Control and Prevention (CDC). HIV/AIDS: PrEP. [cited 2015 May 15]. Available from: http://www.cdc.gov/hiv/bascis/prep.html .
11. Poulia KA. HIV/AIDS. In: Katsilambros N, Dimosthenopoulos C, Kontogianni MD, Manglara E, editors. Clinical nutrition practice. Wiley.com; 2010. P. 147-153.

12. Lutz CA, MA, Mazurr EE, Litch NA. Nutrition and diet therapy: evidence-based applications. 6th edition. F.A. Davis Company, 2015. Chapter 23.
13. World Health Organization. Nutrient requirements for people living with HIV/AIDS: report of a technical consultation. World Health Organization, Geneva, 13-15 May 2003. [cited 2015 May 15]. Avialable from: http://www.who.int/nutrition/publications/hivaids/9241591196/en/ .
14. Food and Agriculture Organization of the United Nations. Enterobacter sakazakii and other microorganisms in powdered infant formula. WHO, Geneva, 2-5 February 2004.
15. McFarland EJ. Human immunodeficiency virus infection. In: Hay WW, Levin MJ, Deterding RR, Abzug MJ, In: Current diagnosis & treatment: Pediatrics. 22nd edition. McGraw-Hill; 2014. Chapter 41.
16. AIDS.gov. HIV/AIDS care continuum. [cited 2015 May 15]. Available from: https://www.aids.gov/federal-resources/policies/care-continuum/ .
17. Shiau S, Arpadi SM, Yin MT. HIV/AIDS and bone health: the role of nutrition. In: Holick MF, Nieves JW, editors. Nutrition and Bone Health. 2nd edition. New York: Springer; 2015. Chapter 38.
18. Knox TA, Jerger L, Tang, AM. Alcohol, HIV/AIDS, and liver disease. In: Watson RR, Preedy VR, Zibadi S, editors. Alcohol, Nutrition, and Health Consequences. New York: Humana Press; 2013. P. 287-303.
19. Staying Healthy with Diet and Exercise When You have HIV/AIDS. In: Judd S, Judd S, Judd S, editors. Health Reference Series: AIDS Sourcebook. Detroit: Omnigraphics; 2011. Section 41.1
20. Centers for Disease Control and Prevention (CDC). Hepatitis information for the public. [cited 2014 Jan 27]. Available from: http://www.cdc.gov/hepatitis.PublicInfo.htm .
21. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases. Viral hepatitis: A through E and beyond. [cited 2014, July 26]. Available from: http://www.nlm.nih.gov/medlineplus/hepatitis.html .
22. Centers for Disease Control and Prevention. Division of Viral Hepatitis and National Center for HIV/AIDS. Viral hepatitis, STD, and TB prevention. [cited 2015 May 1]. Available from: http://www.cdc.gov/hepatitis/ .
23. Centers for Disease Control and Prevention. Breastfeeding: hepatitis B and C infections. [cited 2015 May 1]. Available from: http://www.cdc.gov/breastfeeding/disease/hepatitis.htm .
24. Carroll A & Lutz, MA. Diet in gastrointestinal disease. In: nutrition and diet therapy. Philadelphia: F.A. Davis Company; 2011. Chapter 20.
25. Juve C, Schadewald D, Youngkin EQ, Davis MS, eds. 2013. Women's health: a primary care clinical guide – 4th Edition. New Jersey. Pearson Education; 2013.

Nutrition Risk Criteria **353 FOOD ALLERGY**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	1	RD
CHILDREN	3	RD

Definition

Food allergies are adverse health effects arising from a specific immune response that occurs reproducibly on exposure to a given food. (1)

Presence of condition (food allergy) diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The actual prevalence of food allergies is difficult to establish due to variability in study designs and definitions of food allergies; however recent studies suggest a true increase in prevalence over the past 10 to 20 years (1). A meta-analysis conducted by the National Institute of Allergy and Infectious Disease (NIAID) found the prevalence of food allergy among all age groups between 1-10% (2). Further research has found that food allergy affects more children than recently reported with the prevalence estimated to be 8% (2). Food allergies are a significant health concern as they can cause serious illness and life-threatening reactions. Prompt identification and proper treatment of food allergies improves quality of life, nutritional well-being and social interaction.

Food allergy reactions occur when the body's immune system responds to a harmless food as if it were a threat (3). The most common types of food allergies involve immunoglobulin E (IgE)-mediated responses. The immune system forms IgE against offending food(s) and causes abnormal reactions. IgE is a distinct class of antibodies that mediates an immediate allergic reaction. When food allergens enter the body, IgE antibodies bind to them and release chemicals that cause various symptoms. (1)

According to an expert panel sponsored by the National Institute of Allergy and Infectious Disease, individuals with a family history of any allergic disease are susceptible to developing food allergies and are classified as "at risk" or "high risk." Individuals who are "at risk" are those with a biological parent of sibling with existing, or history of, allergic rhinitis, asthma or atopic dermatitis. Individuals who are "high risk" are those with preexisting severe allergic disease and/or family history of food allergies. (1)

Food Allergies vs. Intolerances

Food intolerances are classified differently from food allergies based on the pathophysiological mechanism of the reactions. Unlike food allergies, food

intolerances do not involve the immune system. Food intolerances are adverse reactions to food caused either by the properties of the food itself, such as a toxin, or the characteristics of the individual, such as a metabolic disorder (4). Food intolerances are often misdiagnosed as food allergies because the symptoms are often similar. Causes of food intolerances may include food poisoning, histamine toxicity, food additives such as monosodium glutamate (MSG), or sulfites (5). The most common food intolerance is lactose intolerance (see Nutrition Risk Criterion 355 *Lactose Intolerance*).

Common Food Allergens

Although reactions can occur from the ingestion of any food, a small number of foods are responsible for the majority of food-induced allergic reactions (6). The foods that most often cause allergic reactions include:

-Cow's milk (and foods made from cow's milk)
-Eggs
-Peanuts
-Tree nuts (walnuts, almonds, cashews, hazelnuts, pecans, brazil nuts)
-Fish
-Crustacean shellfish (e.g. shrimp, crayfish, lobster, and crab)
-Wheat
-Soy

For many individuals, food allergies appear within the first two years of life. Allergies to cow's milk, eggs, wheat and soy generally resolve in early childhood. In contrast, allergy to peanuts and tree nuts typically persist to adulthood. Adults may have food allergies continuing from childhood or may develop sensitivity to food allergens encountered after childhood, which usually continue through life. (1)

Symptoms

There are several types of immune responses to food including IgE-mediated, non-IgE-mediated or mixed. In an IgE-mediated response, the immune system produces allergen-specific IgE antibodies (sigE) when a food allergen first enters the body. Upon re-exposure to the food allergen, the sigE identifies it and quickly initiates the release of chemicals, such as histamine (3). These chemicals cause various symptoms based on the area of the body in which they were released. These reactions occur within minutes or up to 4 hours after ingestion and include symptoms such as urticaria (hives), angioedema, wheezing, cough, nausea, vomiting, hypotension and anaphylaxis (7).

Food-induced anaphylaxis is the most severe form of IgE-mediated food allergies. It often occurs rapidly, within seconds to a few hours after exposure, and is potentially fatal without proper treatment. Food-induced anaphylaxis often affects multiple organ systems and produces many symptoms, including respiratory compromise (e.g., dyspnea, wheeze and bronchospasm), swelling and reduced blood pressure (7). Prompt diagnosis and treatment is essential to prevent life-threatening reactions. Tree nuts, peanuts, milk, egg, fish and crustacean fish are the leading causes of food-induced anaphylaxis (1).

Food allergens may also induce allergic reactions which are non-IgE-mediated. Non-IgE-mediated reactions generally occur more than 4 hours after ingestion, primarily result in gastrointestinal symptoms and are more chronic in nature (7). Examples of non-IgE-mediated reactions to specific foods include celiac disease

(see nutrition risk criterion 354 Celiac Disease), food protein-induced enterocolitis syndrome (FPIES), food protein-induced proctocolitis (FPIP), food protein-induced gastroenteropathy, food-induced contact dermatitis and food-induced pulmonary hemosiderosis (Heiner's syndrome) (accessed May 2012) (8).

The diagnosis of food allergies by a health care provider (HCP) is often difficult and can be multifaceted (see Clarification for more information). Food allergies often coexist with severe asthma, atopic dermatitis (AD), eosinophilic esophagitis (EoE) and exercise-induced anaphylaxis. Individuals with a diagnosis of any of these conditions should be considered for food allergy evaluation. (1)

Prevention

Currently, there is insufficient evidence to conclude that restricting highly allergenic foods in the maternal diet during pregnancy or lactation prevents the development of food allergies in the offspring (9). Adequate nutrition intake during pregnancy and lactation is essential to achieve positive health outcomes. Unnecessary food avoidance can result in inadequate nutrition. There is also a lack of evidence that delaying the introduction of solids beyond 6 months of age, including highly allergenic foods, prevents the development of food allergies. If the introduction of developmentally appropriate solid food is delayed beyond 6 months of age, inadequate nutrient intake, growth deficits and feeding problems can occur. (1)

The protective role that breastfeeding has in the prevention of food allergies remains unclear. There is some evidence for infants at high risk of developing food allergies that exclusive breastfeeding for at least 4 months may decrease the likelihood of cow's milk allergy in the first 2 years of life (9). The American Academy of Pediatrics (AAP) continues to recommend that all infants, including those with a family history of food allergies, be exclusively breastfed until 6 months of age, unless contraindicated for medical reasons (1, 10). For infants who are partially breastfed or formula fed, partially hydrolyzed formulas may be considered as a strategy for preventing the development of food allergies in at-risk infants. According to the AAP, there is no convincing evidence for the use of soy formula as a strategy for preventing the development of food allergies in at-risk infants and therefore it is not recommended. (9)

Management

Food allergies have been shown to produce anxiety and alter the quality of life of those with the condition. It is recommended that individuals with food allergies and their caregivers be educated on food allergen avoidance and emergency management that is age and culturally appropriate. Individuals with a history of severe food allergic reactions, such as anaphylaxis, should work with their HCP to establish an emergency management plan. (1)

Food allergen avoidance is the safest method for managing food allergies. Individuals with food allergies must work closely with their HCP to determine the food(s) to be avoided. This includes the avoidance of any cross-reactive foods, i.e., similar foods within a food group (see Clarification for more information). Nutrition counseling and growth monitoring is recommended for all individuals with food allergies to ensure a nutritionally adequate diet. Individuals with food allergies should also be educated on reading food labels and ingredient lists. (1)

Infants who are partially breastfed or formula fed, with certain non-IgE mediated allergies, such as, FPIES and FPIP may require extensively hydrolyzed casein or amino acid-based formula. According to food allergy experts, children with FPIES can be re-challenged every 18-24 months and, infants/children with FPIP be re-challenged at 9-12 months of age. The re-challenging of foods should be done with HCP oversight. (8)

Implications for WIC Nutrition Services

Through client-centered counseling, WIC staff can assist families with food allergies in making changes that improve quality of life and promote nutritional well-being while avoiding offending foods. Based on the needs and interests of the participant, WIC staff can (as appropriate):

-Facilitate and encourage the participant's ongoing follow-up with the HCP for optimal management of the condition.
-Promote exclusive breastfeeding until six months of age and continue through the first year (10).
-Provide hypoallergenic formula for participants with appropriate medical documentation, as needed.
-Tailor food packages to substitute or remove offending foods.
-Educate participants about reading food labels and identifying offending foods and ingredients.
See resources below:

http://www.fda.gov/downlaods/ForConsumers/ConsumerUpdates/UCM254727.pdf
Accessed May 2012.
http://www.webmd.com/allergies/foodtriggers. Accessed May 2012.
http://www.foodallergy.org/section/how-to-read-a-label. Accessed May 2012.

-Educate participants on planning meals and snacks for outside the home.
-Refer participants to their HCP for a re-challenge of offending foods, as appropriate.
-Establish/maintain communication with participant's HCP.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Food allergies are diagnosed by a HCP by evaluating a thorough medical history and conducting a physical exam to consider possible trigger foods to determine the underlying mechanism of the reaction, which guides testing. Along with a detailed history of the disorder, such as symptoms, timing, common triggers and associations, there are several types of tests that the HCP may use in diagnosing food allergies. These include the following:

- Food Elimination Diet
- Oral Food Challenges
- Skin Prick Test (SPT)
- Allergen-specific serum IgE (sigE)

- Atopy Patch Test

Diagnosing food allergies is difficult because the detection of sigE does not necessarily indicate a clinical allergy. Often, more than one type of test is required to confirm a diagnosis. The double-blind, placebo-controlled food challenge is considered the gold standard in testing for food allergies. (11)

Children often outgrow allergies to cow's milk, soy, egg, and wheat quickly; but are less likely to outgrow allergies to peanut, tree nuts, fish, and crustacean shellfish. If the child has had a recent allergic reaction, there is no reason to retest. Otherwise, annual testing may be considered to see if the allergy to cow's milk, soy, egg, or wheat has been outgrown so the diet can be normalized. (1)

Cross-reactive food: When a person has allergies to one food, he/she tends to be allergic to similar foods within a food group. For example, all shellfish are closely related; if a person is allergic to one shellfish, there is a strong chance that person is also allergic to other shellfish. The same holds true for tree-nuts, such as almonds, cashews and walnuts. (1)

References

1. Boyce, J. et al. Guidelines for the diagnosis and management of food allergy in the United States: Report of the NIAID-Sponsored Expert Panel. Journal of Allergy and Clinical Immunology. 2012; 126(6):S1-S58. http://www.jacionline.org/artical/S0091-6749(10)01566-6/fulltext. Accessed May 2012.
2. Gupta, R. et al. The prevalence, severity and distribution of childhood food allergy in the United States. Pediatrics. 2011;128(1):e9-e17. Available at: http://pediatrics.aappublications.org/content/128/1/e9.full.html. Accessed May 2012.
3. National Institute of Allergy and Infectious Disease website: How do allergic reactions work? Available at: http://www.niaid.nih.gov/topics/foodallergy/understanding/pages/whatisit.aspx. Accessed May 2012.
4. Cianferoni, A, Spergel, JM. Food Allergy: review, classification and diagnosis. Allergology International. 2009; 58:457-466.
5. National Institute of Allergy and Infectious Disease. Food allergy: An overview. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, 2010 (NIH Publication No. 11-5518). Available at: http://www.niaid.nih.gov/topics/foodallergy/understanding/pages/whatisit.aspx. Accessed May 2012.
6. Sampson, HA. Food Allergy. Part 1: Immunopathogenesis and clinical disorders. Journal of Allergy and Clinical Immunology. 1999; 103(5):717-728.
7. Davis, C. Food allergies: clinical manifestations, diagnosis, and management. Current Problems in Pediatric Adolescent Health Care. 2009: 39:236-254.
8. Metcalfe DD, Sampson HA, Simon RA, editors. Food allergy: adverse reactions to food and food additives. 4th ed. Malden (MA): Blackwell Publishing; 2008.
9. Greer, F. et al. American Academy of Pediatrics Committee on Nutrition. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. Pediatrics. 2008; 121(1), pages 183-191.

10. Gartner LM, Morton J, Lawrence RA, et al. American Academy of Pediatrics. Policy Statement: Breastfeeding and the use of human milk. Pediatrics. 2005; 115 (2): 496-506.
11. Sicherer S.H., Sampson, HA. Food allergy. Journal of Allergy and Clinical Immunology. 2010; (125):S116-S125.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Celiac Disease (CD) is an autoimmune disease precipitated by the ingestion of gluten (a protein in wheat, rye, and barley) that results in damage to the small intestine and malabsorption of the nutrients from food. (1). (For more information about the definition of CD, please see the Clarification section)

CD is also known as:

- Celiac Sprue
- Gluten-sensitive Enteropathy
- Non-tropical Sprue

Presence of Celiac Disease diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

CD affects approximately 1% of the U.S. population (2,3). CD can occur at any age and the treatment requires strict adherence to a gluten-free diet for life. CD is both a disease of malabsorption and an abnormal immune reaction to gluten. When individuals with CD eat foods or ingest products containing gluten, their immune system responds by damaging or destroying villi---the tiny, fingerlike protrusions lining the small intestine. Villi normally allow nutrients from food to be absorbed through the walls of the small intestine into the bloodstream (4). The destruction of villi can result in malabsorption of nutrients needed for good health. Key nutrients often affected are iron, calcium and folate as they are absorbed in the first part of the small intestine. If damage occurs further down the small intestinal tract, malabsorption of carbohydrates (especially lactose), fat and fat-soluble vitamins, protein and other nutrients may also occur (2,5).

In addition to the gastrointestinal system, CD affects many other systems in the body, resulting in a wide range and severity of symptoms. Symptoms of CD may include chronic diarrhea, vomiting, constipation, pale foul-smelling fatty stools and weight loss. Failure to thrive may occur in infants and children. The vitamin and mineral deficiencies that can occur from continued exposure to gluten may result in conditions such as anemia, osteoporosis and neurological disorders such as ataxia, seizures and neuropathy.

Individuals with CD who continue to ingest gluten are also at increased risk for developing other autoimmune disorders (e.g. thyroid disease, type 1 diabetes, Addison's disease) and certain types of cancer, especially gastrointestinal malignancies (2).

Continued exposure to gluten increases the risk of miscarriage or having a low birth weight baby, and may result in infertility in both women and men. A delay in diagnosis for children may cause serious nutritional complications including growth failure, delayed puberty, iron-deficiency anemia, and impaired bone health. Mood swings or depression may also occur (2, 6). See Table 1 for Nutritional Implications and Symptoms.

Table 1. Nutritional Implications and Symptoms of CD
Common in Children:
Digestive Symptoms- more common in infants and children, may include: • Vomiting • Chronic diarrhea • Constipation • Abdominal bloating and pain • Pale, foul-smelling, or fatty stool
Other Symptoms: • Delayed puberty • Dental enamel abnormalities of the permanent teeth • Failure to thrive (delayed growth and short stature) • Weight loss • Irritability
Common in Adults:
Digestive Symptoms- same as above, less common in adults
Other Symptoms- adults may instead have one or more of the following: • Unexplained iron-deficiency anemia • Other vitamin and mineral deficiencies (A,D,E, K, calcium) • Lactose intolerance • Fatigue • Bone or joint pain • Arthritis • Depression or anxiety • Tingling numbness in the hands and feet • Seizures • Missed menstrual periods • Infertility (men and women) or recurrent miscarriage • Canker sores inside the mouth • Itchy skin rash- dermatitis herpetiformis • Elevated liver enzymes
Sources:

Case, Shelley, Gluten-Free Diet, A Comprehensive Resource Guide, Case Nutrition Consulting Inc., 2008.

National Institute of Diabetes and Digestive and Kidney Diseases, Celiac Disease, NIH Publication No. 08-4269 September 2008. http://digestive.niddk.nih.gov/ddiseases/pubs/celiac/#what. Accessed May 2012.

The risk for development of CD depends on genetic, immunological, and environmental factors. Recent studies suggest that the introduction of small amounts of gluten while the infant is still breast-fed may reduce the risk of CD. Both breastfeeding during the introduction of dietary gluten, and increasing the duration of breastfeeding were associated with reduced risk in the infant for the development of CD. It is not clear from studies whether breastfeeding delays the onset of symptoms or provides a permanent protection against the disease. Therefore, it is prudent to avoid both early (<4 months) and late (≥7 months) introduction of gluten and to introduce gluten gradually while the infant is still breast-fed, as this may reduce the risk of CD. (7)

The only treatment for CD is a gluten-free diet. Individuals with CD should discuss gluten-free food choices with a dietitian or physician that specializes in CD. Individuals with CD should always read food ingredient lists carefully to make sure that the food does not contain gluten. Making informed decisions in the grocery stores and when eating out is essential for the successful treatment of the disease (5, 8).

Implications for WIC Nutrition Services

Through client-centered counseling, WIC staff can assist participants with CD in making gluten-free food choices that improve quality of life and promote nutritional well-being. WIC can provide nutrition education/counseling on alternatives to gluten-containing food products as well as provide gluten-free grain selections available in the WIC food packages. Based on the needs and interests of the participant, WIC staff may (as appropriate):

- Promote breastfeeding throughout the first year of life, with exclusive breastfeeding until 4-6 months of age.
- In consultation with the guidance of a medical provider, introduce gluten-containing foods between 4 and 6 months to infants at risk of CD, including infants with a parent or sibling with CD.
- Tailor food packages to substitute or remove gluten-containing foods.
- Educate participants on meeting nutritional needs in the absence of gluten-containing foods.
- Encourage high fiber, gluten-free grain selections.
- Monitor participant's growth pattern and weight status.
- Educate participants on planning gluten-free meals and snacks for outside the home.
- Provide educational materials outlining allowed foods and foods to avoid, for example:
  http://www.celiac.nih.gov/Default.aspx Accessed May 2012.
  http://www.naspghan.org/user-assets/Documents/pdf/diseaseInfo/GlutenFreeDietGuide-E.pdf Accessed May 2012

Provide referrals as appropriate.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

The 2006 American Gastroenterological Association (AGA) Institute Technical Review on the Diagnosis and Management of Celiac Disease refers to CD as "a unique disorder that is both a food intolerance and autoimmune disorder" (9). According to the 2010 NIAID-Sponsored Expert Panel definition, CD is a non-IgE mediated food allergy (10). (See nutrition risk criterion 353 Food Allergy.) However, the Expert Panel did not include information about CD in its report but rather refers readers to existing clinical guidelines on CD including the AGA Institute's Technical Review. (5,9,10)

References

1. National Institute of Allergy and Infectious Disease. Food allergy: an overview. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, 2010 (NIH Publication No. 11-5518). Available at: http://www.niaid.nih.gov/topics/foodallergy/understanding/pages/whatisit.aspx. Accessed May 2012.
2. Case, S. Gluten-free diet: A comprehensive resource guide. Case Nutrition Consulting Inc., 2008.
3. Green, PHR, Cellier, C. Medical progress-celiac disease. The New England Journal of Medicine. 2007 Oct 25:1731-1743.
4. National Institute of Diabetes and Digestive and Kidney Diseases, Celiac Disease, National Institute of Health. Celiac disease. Available at: http://digestive.niddk.nih.gov/ddiseases/pubs/celiac/#what. Accessed May 2012.
5. National Institute of Diabetes and Digestive and Kidney Diseases, Celiac Disease, NIH Publication No. 08-4269 September 2008.
6. Guideline for the Diagnosis and Treatment of Celiac Disease in Children: Recommendation of the North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J Pediatr Gastroenterol Nutr. 2005 Jan;40(1):1-19.
7. ESPGHAN Committee on Nutrition: Agostoni, C. et al. Complementary feeding: A commentary by the ESPGHAN Committee on Nutrition, Medical Position Paper. Journal of Pediatric Gastroenterology. 2006 September: 67-92.
8. Raymond, N, Heap, J, Case, S. The gluten-free diet: An update for health professionals. Practical Gastroenterology. 2006 September: 67-92.
9. American Gastroenterological Association (AGA) Institute Technical Review on the Diagnosis and Management of Celiac Disease. Gastroenterology. 2006 Dec;131(6):1981-2002.
10. Boyce, J. et al. Guidelines for the diagnosis and management of food allergy in the United States: Report of the NIAID-Sponsored Expert Panel. Journal of Allergy and Clinical Immunology. 2010;126(6):S1-S58. http://www.jacionline.org/article/S0091-6749(10)01566-6/fulltext. Accessed May 2012.

Nutrition Risk Criteria **355 LACTOSE INTOLERANCE**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	1	RD
CHILDREN	3	RD

Definition

Lactose intolerance is the syndrome of one or more of the following: diarrhea, abdominal pain, flatulence, and/or bloating, that occurs after lactose ingestion.

Presence of condition diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Lactose intolerance occurs because of a deficiency in the levels of the lactase enzyme (1). Many variables determine whether a person with lactase deficiency develops symptoms. They include: the dose of lactose ingested; the residual intestinal lactase activity; the ingestion of food along with lactose; the ability of the colonic flora to ferment lactose; and, the individual sensitivity to the products of lactose fermentation (1). Some forms of lactase deficiencies may be temporary, resulting from premature birth or small bowel injuries, and will correct themselves, leaving individuals with the ability to digest lactose sufficiently (2).

Primary lactase deficiency is attributable to relative or absolute absence of lactase that develops in childhood, and is the most common cause of lactose malabsorption and lactose intolerance (2).

Secondary lactase deficiency is one that results from small bowel injury, such as acute gastroenteritis, persistent diarrhea, or other causes that injure the small intestine mucosa, and can present at any age, but is more common in infancy. Treatment of secondary lactase deficiency and lactose malabsorption attributable to an underlying condition generally do not require elimination of lactose from the diet. Once the primary problem is resolved, lactose-containing products can be consumed normally. (2)

Congenital lactase deficiency is a rare disorder that has been reported in only a few infants. Affected newborn infants present with intractable diarrhea as soon as human milk or lactose-containing formula is introduced. (2)

Developmental lactase deficiency is the relative lactase deficiency observed among pre-term infants of less than 34 weeks gestation (2). One study in preterm infants reported benefit from the use of lactase-supplemented feedings or lactose-reduced formulas (3). The use of lactose-containing formulas and human milk does not seem to have any short- or long-term deleterious effects in preterm infants (2).

Lactose is found primarily in milk, milk-based formula and other dairy products, which provide a variety of nutrients essential to the WIC population (calcium, vitamin D, protein). Lactose intolerance varies according to individuals. Some individuals may tolerate various quantities of lactose without discomfort, or tolerate it when consumed with other foods. Dairy products that are soured, or otherwise treated with bacteria that secrete lactase (e.g., Lactobacillus acidophilus), such as cheese and yogurt, are easier to digest in lactose-intolerant individuals because they contain relatively low levels of lactose. (4)

Many individuals diagnosed with lactose intolerance avoid dairy all together. Also, lactose intolerance has been shown to be associated with low bone mass and increased risk of fracture (5). Inadequate dairy intake increases the risk of metabolic syndrome, hypertension, preeclampsia, obesity and certain forms of cancer, especially colon cancer (6).

Implications for WIC Nutrition Services

It is important to assess participants individually for lactose tolerances and nutrient needs to determine the best plan of action. WIC can provide client-centered counseling to incorporate tolerated amounts of lactose-containing foods and/or other dietary sources of calcium, vitamin D and protein into participants' diets. WIC foods such as cheese, lactose-free milk, soy beverages, tofu and calcium fortified foods (like juice) can provide these nutrients to participants with lactose intolerance. Based on the needs and interests of the participant, WIC staff can, in addition, also offer the following strategies (as appropriate):

- **Except for infants with congenital lactase deficiency,** promote exclusive breastfeeding until six months of age and continue breastfeeding through the first year. For infants with congenital lactase deficiency, treatment is removal and substitution of lactose from the diet with a commercial lactose-free formula (2).
- Tailor food packages to substitute or remove lactose-containing foods.
- Educate participants on meeting nutritional needs in the absence of lactose-containing foods.
- Educate participants on planning lactose-free/lactose-reduced meals and snacks for outings, social gatherings, school and/or work.

Any WIC participant suspected to have lactose intolerance should be referred to a health care provider for evaluation and appropriate diagnosis (7), if needed (see Clarification for additional information on diagnosing Lactose intolerance).

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Lactose malabsorption can be diagnosed with a hydrogen breath test. The test involves having individuals ingest a standard dose of lactose after fasting. Elevated levels of breath hydrogen, which are produced by bacterial fermentation of undigested lactose in the colon, indicate the presence of lactose

malabsorption (1). The hydrogen breath test is not routinely ordered, and instead, patients are frequently asked to assess symptoms while avoiding dairy products for a period of time followed by a lactose product challenge to determine if they are lactose intolerant (7). The demonstration of lactose malabsorption does not necessarily indicate that an individual will be symptomatic.

References

1. National Institutes of Health Consensus Development Conference Statement: Lactose intolerance and health. February, 2010. Available at: http://consensus.nih.gov/2010/lactosestatement.htm. Accessed May 2012.
2. Heyman MB. Lactose intolerance in infants, children, and adolescents; Pediatrics 2006 September: 118 (#3) 1279-1286. http://aappolicy.aappublications.org/cgi/reprint/pediatrics;118/3/1279.pdf. Accessed May 2012.
3. Shulman RJ, Feste A, Ou C. Absorption of lactose, glucose polymers, or combination in premature infants. J Pediatr. 1995;127:626-631.
4. Ranciaro A, Tishoff SA. Population genetics: evolutionary history of lactose tolerance in Africa [abstract]. NIH Consensus Development Conference Lactose Intolerance and Health; February 2010; 43-47.
5. U.S. Department of Health and Human Services- Office of the Surgeon General. Bone health and osteoporosis: a report of the surgeon general. 2004.
6. Hearney RP. Consequences of excluding dairy or of avoiding milk in adults [abstract]. NIH Consensus Development Conference Lactose Intolerance and Health. February, 2010; 73-77.
7. Chang, Lin MD. Clinical Presentation: But what if it is not lactose intolerance? [abstract]. NIH Consensus Development Conference Lactose Intolerance and Health; February 2010; 39-42.

Additional Reference

1. National Dairy Council [Internet]. Lactose Intolerance Health Education Kit (2011). Available at: http://www.nationaldairycouncil.org/EDUCATIONMATERIALS/HEALTHPROFESSIONALSEDUCATIONKITS/Pages/LactoseIntoleranceHealthEducationKit.aspx. Accessed May 2012.

Nutrition Risk Criteria **356** **HYPOGLYCEMIA**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	1	-
CHILDREN	3	-

Definition

Presence of hypoglycemia diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Hypoglycemia can occur as a complication of diabetes, as a condition in itself, in association with other disorders, or under certain conditions such as early pregnancy, prolonged fasting, or long periods of strenuous exercise (1).

Symptomatic hypoglycemia is a risk observed in a substantial proportion of newborns who are small for gestational age (SGA), but it is uncommon and of shorter duration in newborns who are of the appropriate size for gestational age (2).

WIC can provide nutrition management that concentrates on frequent feedings to support adequate growth for infants and children (2). WIC can also provide nutrition education to help manage hypoglycemia in women that includes consuming a balanced diet, low carbohydrate snacks and exercise (1).

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. National Institute of Diabetes, Digestive and Kidney Diseases. Hypoglycemia. National Diabetes Information Clearinghouse, 199. Available at: http://www.niddk.nih.gov/health/diabetes/pubs/hypo/hypo.htm.
2. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington D.C.; 1996. p.217-218.

Nutrition Risk Criteria **357 DRUG NUTRIENT INTERACTIONS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Use of prescription or over-the-counter drugs or medications that have been shown to interfere with nutrient intake or utilization, to an extent that nutritional status is compromised.

Required Documentation

Medications

Justification

The drug treatment of a disease or medical condition may itself affect nutritional status. Drug induced nutritional deficiencies are usually slow to develop and occur most frequently in long-term drug treatment of chronic disease. Possible nutrition-related side effects of drugs include, but are not limited to, altered taste sensation, gastric irritation, appetite suppression, altered GI motility, and altered nutrient metabolism and function, including enzyme inhibition, vitamin antagonism, and increased urinary loss.

The marketplace of prescribed and over-the-counter drugs is a rapidly changing one. For knowledgeable information on the relationship of an individual's drug use to his/her nutritional status, it is important to refer to a current drug reference such as Physician's Desk Reference (PDR), a text such as Physician's Medication Interactions, drug inserts, or to speak with a pharmacist.

References

1. Allen, M: Food-Medication Interactions; 7th edition; Tempe, Arizona; 1991.
2. Physician’s Desk Reference, 51st edition; Montvale, New Jersey; Medical Economics Company, Inc.; 1997.
3. Diet and Drug Interactions. Daphne A. Roe, M.D., F.R.C.P.
4. Handbook on Drug and Nutrient Interactions: A Reference and Study Guide.
5. Institute of Medicine: WIC Nutrition Risk Criteria: A Scientific Assessment; 1996; pp. 217-218.
6. Pronsky, ZM: Powers and Moore’s Food Medications Interactions; 10th edition; 1997.

Nutrition Risk Criteria **358 EATING DISORDERS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD

Definition

Eating disorders (anorexia nervosa and bulimia), are characterized by a disturbed sense of body image and morbid fear of becoming fat. Symptoms are manifested by abnormal eating patterns including, but not limited to:

- self-induced vomiting
- purgative abuse
- alternating periods of starvation
- use of drugs such as appetite suppressants, thyroid preparations or diuretics
- self-induced marked weight loss

Presence of eating disorder(s) diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders or evidence of such disorders documented by the CPA.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Anorexia nervosa and bulimia are serious eating disorders that affect women in the childbearing years. These disorders result in general malnutrition and may cause life-threatening fluid and electrolyte imbalances. Women with eating disorders may begin pregnancy in a poor nutritional state. They are at risk of developing chemical and nutritional imbalances, deficiencies, or weight gain abnormalities during pregnancy if aberrant eating behaviors are not controlled. These eating disorders can seriously complicate any pregnancy since the nutritional status of the pregnant woman is an important factor in perinatal outcome.

Maternal undernutrition is associated with increased perinatal mortality and an increased risk of congenital malformation. While the majority of pregnant women studied reported a significant reduction in their eating disorder symptoms during pregnancy, a high percentage of these women regressed in the postpartum period. This regression in postpartum women is a serious concern for breastfeeding and non-breastfeeding postpartum women who are extremely preoccupied with rapid weight loss after delivery.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Worthington-Roberts, B., and Williams, SR: Nutrition in Pregnancy and Lactation; 5th ed.; Mosby Pub; St. Louis; pp.270-271.
2. Strober, M: International Journal of Eating Disorders; Vol. 8, No. 3; 1986; pp.285-295.
3. Institute of Medicine: Nutrition Services in Perinatal Care; 1992, p. 20.
4. Clinical Issues Perinatal Womens Health Nursing; 1992;3(4); pp. 695-700.
5. Krummel DA, and Kris-Etherton, PM: Nutrition in Women's Health, Aspen Pub; Gaithersburg, MD; pp. 58-102.

Nutrition Risk Criteria **359 RECENT MAJOR SURGERY, PHYSICAL TRAUMA, BURNS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Major surgery (including cesarean sections - “C-sections”), physical trauma or burns severe enough to compromise nutritional status.

Any occurrence:

- Within the past two (≤ 2) months may be self-reported.
- More than two (> 2) months previous must have the continued need for nutritional support diagnosed by a physician or a health care provider working under the orders of a physician.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

The body's response to injuries such as major surgeries, physical trauma, or burn may affect nutrient requirements needed for recovery and lead to malnutrition. The catabolic response to these injuries causes a hypermetabolic state in the body. The alteration in metabolism not only increases the individual’s calorie and protein needs, but they also increase the needs for certain vitamins, minerals, fatty acids, and amino acids. (1)

Proper wound healing is essential in the recovery of surgeries, physical trauma, and burns. Normal wound healing is a complex process and involves three phases: inflammation, proliferation, and remodeling (1, 2). Each phase of wound healing involves growth factors, other biologically active molecules, and specific vitamins and minerals such as Vitamin A, Vitamin C, and Zinc. The process of wound healing does not always follow the three stages sequentially and can sometimes move forward or regress based on nutrition status and response to treatment (3, 4). Even after a wound is closed, the individual’s metabolic rate and need for additional nutrition can remain high (5).

Factors that can prevent proper wound healing or can increase the time needed for a wound to heal include (2, 6):

- Malnutrition prior to the surgery, injury or burn
- Infections
- Diabetes
- Poor blood flow
- Obesity
- Age
- Heavy alcohol use
- Stress
- Medications
- Smoking

Because healing is a complex process and is impacted by a variety of factors, it is inappropriate to expect a set recovery time for an individual based solely on the type and severity of the injury (7). For some individuals, they may no longer be at increased nutritional risk within a couple weeks of their injury. For others, recovery from the same type and severity of injury may take months.

Major Surgery and Wound Healing

Many types of surgeries are completed as noninvasive procedures and do not result in large incisions that require additional medical and nutritional care to heal. However, many surgical procedures (including cesarean sections) do not involve incisions that, if left unaddressed, could lead to infection. Major surgeries are surgeries that involve a risk to the life of the individual and include operations on organs within the body (8). Removal of a portion of the large or small intestine, heart surgery, and bariatric surgery are examples of major surgeries. Minor surgeries are surgeries that involve little risk to the individual and include operations on the superficial structures of the body (9). Ear tubes, the most common childhood surgery performed with anesthesia, are an example of a minor surgery that does not impact nutrition status (10).

Cesarean sections are considered a major surgery and, therefore, require additional assessment and education in the WIC clinic. In the US, the rate of cesarean delivery rose from 19.7% of singleton births in 1996 to 31.3% of singleton births in 2011 (11). Reasons for a cesarean delivery include: multiple pregnancy, labor fails to progress, medical concerns for the infant, problems with the placenta, a large infant, breech position, maternal infections, and medical conditions in the mother (i.e. diabetes or high blood pressure) (12).

Nutritional Considerations for Major Surgery/Wound Healing

The role of specific nutrients in wound healing continues to be explored and studies are conducted regularly to assess the role vitamins, minerals, fatty acids, amino acids, and carbohydrates play in proper wound healing. Nutrient supplements above the Recommended Dietary Allowances (RDA) may be necessary to aid in wound healing. However, before using any additional supplement to assist in wound healing, energy and protein requirements of the individual must be met (13, 14). Amino acids are essential to the repair of damaged tissue in the body. Amino acids are divided into three categories: essential (must be obtained through foods), nonessential (can be produced in the body), and conditionally essential (produced in the body except in cases of injury or illness). Arginine and Glutamine are examples of conditionally essential amino acids. The following table highlights the roles of these nutrients in the wound healing process:

Nutrient	Role in Wound Healing
Arginine	Involved in secretion of growth hormone (12)
Omega-3fatty acids	Reduces wound infections (12)
Vitamin C	Collagen synthesis (2)
Vitamin A	Immune function and cellular communication (15)

Vitamin E	Antioxidant (16)
Vitamin D	Modulates cell growth Neuromuscular and immune function Reduces inflammation (17)
Magnesium	Co-factor for enzymes involved in protein and collagen synthesis (2)
Copper	Co-factor for cross-linking of collagen (2)
Zinc	Involved in RNA and DNA polymerase (2)
Iron	Aids in the synthesis of some growth hormones and connective tissue (18)

Following a cesarean section, a breastfeeding mother may experience difficulty finding a comfortable nursing position that does not cause pain with the incision. She may also have difficulty breastfeeding if the infant is drowsy due to the pain medication administered during the procedure. A referral to a lactation specialist can help ensure that the mother is successful in reaching her breastfeeding goals.

Physical Trauma

Physical trauma is usually a result of accidents and injuries that often lead to fractures, wounds, and subsequent hospitalization. Physical trauma can be divided into blunt force trauma, penetrating trauma, and trauma from surgery. Blunt force trauma is the result of an object (or force) striking the body, causing concussions, lacerations or fractures. Penetrating trauma is trauma that occurs as a result of an object piercing the skin, causing an open wound (7). Fracture healing is a process that begins with a hemorrhage and progresses through three stages: inflammatory, reparative, and remodeling.

Physical trauma can also be a result of domestic and/or child abuse. In addition to the physical effects of abuse, victims of abuse often experience acute and ongoing psychological and emotional trauma that may also impact an individual's nutrition status. Poor appetite, undesirable food choices, and using food for coping can impact both women and children. Children may also begin hoarding food in cases of abuse or neglect.

Nutritional Considerations for Physical Trauma

In addition to an increase in energy, protein, and micronutrients needed for proper wound healing, physical trauma that includes fractures requires additional nutrients for proper bone healing. In some cases, the physical trauma will lead to temporary or lifelong difficulty with self-feeding. Research on the roles specific nutrients play in fracture healing continues to expand. Key nutrients for bone health include calcium, phosphorus, fluoride, magnesium, sodium, vitamin D, vitamin A, vitamin K, vitamin C, vitamin B6, folate, and vitamin B12. Meeting RDAs set for these nutrients is important for bone health and bone healing (19). For some individuals, intakes above the RDA may be recommended by their medical provider to assist in bone healing; however, some nutrients including fluoride, sodium, and vitamin A may negatively impact bone health when intake is above the recommended level (19).

Burns

Burns can be caused by heat (including hot surfaces, fires, and hot liquids), chemicals, electricity, sunlight or nuclear radiation. There are three stages of burns based on what layers of the skin are burned. A first degree burn only affects the outer layer of the skin (epidermis). A second-degree burn damages the epidermis and the layer directly under the epidermis (dermis). A third-degree burn damages the epidermis, dermis, and damages the tissue underneath the skin. (20)

Burns are also classified based on the surface area pf the body that has been burned (Percent Total Body Surface Area or TBSA). For example, a burn that covers one hand and arm would be 9%TBSA, whereas a burn that covers a person's back would be 18% TBSA (21). Increases in the surface area affected by the burn result in a greater potential for fluid loss and infection (21). Inhalation burns are burns that occur inside an individual's lungs and internal organs. Once discharged from the hospital, enteral feedings may be prescribed to aid in healing.

Nutritional Considerations for Burns

The nutrition status of burn patients is monitored very closely during hospitalization and after discharge. Following a severe burn, the body goes into a catabolic state and the body begins to breakdown skeletal muscle (5). This state increases the requirements for energy, protein, carbohydrates, fats, vitamins, minerals, and antioxidants (22). Damaged blood vessels also increase fluid loss and can lead to dehydration or shock (19). Nutrition care in the hospital setting for individual's recovering from burns may also include parenteral or enteral nutrition support depending on the severity of the burns. Glutamine, a conditionally essential amino acid, can improve the healing of burns (23).

Implications for WIC Nutrition Services

Most surgeries, physical traumas, and burns are unexpected. The education and supplemental food that WIC provides can help ensure that the individual is in good nutritional health prior to the surgery, physical trauma or burn. Following a major surgery, physical trauma, and/or burn, an individual will be at increased nutritional risk until the injury has completely healed. WIC staff can improve outcomes following an injury by:

- Assuring that vitamin and mineral intakes meet the RDAs (unless amounts that exceed the RDAs are recommended by their medical provider).
- Assuring that energy and protein intake preserve lean muscle mass and body weight.
- Recommending a participant speak with their medical provider about a multivitamin supplement when diet alone cannot meet the RDAs for vitamins and minerals.
- Referring to community resources for smoking cessation, support groups, food assistance, and safe living environments (in cases of physical abuse).
- Referring to a lactation educator if women experience difficulty breastfeeding following a cesarean section.

Reference

1. Gurtner GC. Wound Healing: normal and abnormal. In: Thorne CH, editor. Grabb and Smith's Plastic Surgery. Riverwoods, IL;2007. p.15-22.
2. Guo S, DiPietro LA. Factors affecting wound healing. J Dent Res. 2010; 89:219-229.
3. Wild T, Rahbarnia A, Kellner M, Sobotka L, Eberlein T. Basics in nutrition and wound healing. J Nut. 2010;26:862-866.
4. Stechmiller JK. Understanding the role of nutrition and wound healing. Nutr Clin Pract. 201;25:61-68.
5. Prelack K, Dylewski M, Sheridan RL. Practical guidelines for nutritional management of burn injury and recovery. J Burns.2007;33:14-24
6. National Institutes of Health [Internet]. Maryland; c1997-2015 [updated 2015 Jun 15; cited 2015 Jul 6]. How wounds heal; [about 3 screens]. Available from: https://www.nlm.nih.gov/medlineplus/ency/patientinstructions/000741.htm .
7. National Institutes of Health [Internet]. Maryland: [updated 2014 Nov 17; cited 2015 Aug 4]. Trauma Fact Sheet; [about 2 screens]. Available from: http://nigms.nih.gov/Education/Pages?Factsheet_Trauma.aspx .
8. Merriam-Webster Medical Dictionary. [cited 2015 Jul 6]. Available from: http://www.merriam-webster.com/medical/major+surgery .
9. Merriam-Webster Medical Dictionary. [cited 2015 Jul 6]. Available from: http://www.merriam-webster.com/medical/minor+surgery .
10. American Academy of Otolaryngology-Head and Neck Surgery [Internet]. Virginia; c2015 [cited 2015 Aug 12]. Ear Tubes; [about 5 screens]. Available from: http://www.entnet.org/content/ear-tubes .
11. Osterman MJK, Martin JA. Changes in cesarean delivery rates by gestational age: United States, 1996-2011. NCHS Data Brief. 2013;124.
12. National Institutes of Health [Internet]. Maryland: [updated 2012 Nov 30; cited 2015 Jul 6]. When is a cesarean delivery necessary and what are the risks; [about 1 screen]. Available from: http://www.nichd.nih.gov/health/topics/obstetrics/conditionsinfo/pages/risks.aspx .
13. Alexander JW, Supp D. Role of arginine and omega-3 fatty acids in wound healing and infection. Adv Wound Care. 2014;3:682-690.
14. Ellinger S, Stehle P. Efficacy of vitamin supplementation in situations with wound healing disorders: results from clinical intervention studies. Curr Opin Clin Nutr Metab Care. 2009;12:588-95.
15. National Institutes of Health [Internet]. Maryland: [updated 2013 Jun 5; cited 2015 Jul 6]. Vitamin A fact sheet for health professionals; [about 16 screens]. Available from: https://ods.od.nih.gov/factsheets/VitaminA-HealthProfessional/ .
16. National Institutes of Health [Internet]. Maryland: [updated 2013 Jun 5; cited 2015 Jul 6]. Vitamin E fact sheet for health professionals; [about 11 screens]. Available from: https://ods.od.nih.gov/factsheets/VitaminE-HealthProfessional/ .
17. National Institutes of Health [Internet]. Maryland: [updated 2014 Nov 10; cited 2015 Jul 6]. Vitamin D fact sheet for health professionals; [about 9 screens]. Available from: https://ods.od.nih.gov/factsheets/Vitamin D-HealthProfessional/ .
18. National Institutes of Health [Internet]. Maryland: [updated 2015 Feb 19; cited 2015 Jul 6]. Iron fact sheet for health professionals; [about 20 screens]. Available from:

https://ods.od.nih.gov/factsheets/Iron-HealthProfessional/ .
19. Angelo G (Oregon State University, Linus Pauling Institute, Corvallis, OR). Micronutrient Information Center; 2012 Aug.
20. National Institutes of Health [Internet]. Maryland: [updated 2014 Aug 8; cited 2015 May 12]. Burns Fact Sheet; [about 2 screens]. Available from: http://nigms.nih.gov/Education/Pages/Factsheet_Burns.aspx .
21. National Institutes of Health [Internet]. Maryland: [updated 2011 Jun 25; cited 2015 Jul 6]. Burn triage and treatment – thermal injuries [about 11 screens]. Available from: http://chemm.nlm.nih.gov/burns.htm .
22. Chan MM, Chan GM. Nutritional therapy for burns in children and adults. J Nut.2009;25:261-269.
23. Ellinger S. Micronutrients, arginine, and glutamine: does supplementation provide an efficient tool for prevention and treatment of different kinds of wounds. Adv Wound Care. 2014;3:691-707.

Additional Reference:

Mahan LK, Raymond JL. Krause's food & nutrition care process. 14th ed. Philadelphia (PA): Elsevier BV; 2016.

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Diseases or conditions with nutritional implications that are not included in any of the other medical conditions. The current condition, or treatment for the condition, must be severe enough to affect nutritional status. Includes, but is not limited to:

- juvenile rheumatoid arthritis (JRA)
- lupus erythematosus
- cardiorespiratory diseases
- heart disease
- cystic fibrosis
- persistent asthma (moderate or severe) requiring daily medication

Presence of medical condition(s) diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Document source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Juvenile rheumatoid arthritis (JRA) is the most common pediatric rheumatic disease and most common cause of chronic arthritis among children. JRA puts individuals at risk of anorexia, weight loss, failure to grow, and protein energy malnutrition.

Lupus erythematosus is an autoimmune disorder that affects multiple organ systems. Lupus erythematosus increases the risk of infections, malaise, anorexia, and weight loss. In pregnant women, there is increased risk of spontaneous abortion and late pregnancy losses (after 28 weeks gestation).

Cardiorespiratory diseases affect normal physiological processes and can be accompanied by failure to thrive and malnutrition. Cardiorespiratory diseases put individuals at risk for growth failure and malnutrition due to low calorie intake and hypermetabolism.

Cystic fibrosis (CF), a genetic disorder of children, adolescents, and young adults characterized by widespread dysfunction of the exocrine glands, is the most common lethal hereditary disease of the Caucasian race.

Many aspects of the disease of CF stress the nutritional status of the patient directly or indirectly by affecting the patient's appetite and subsequent intake. Gastrointestinal losses occur in spite of pancreatic enzyme replacement therapy.

Also, catch-up growth requires additional calories. All of these factors contribute to a chronic energy deficit, which can lead to a marasmic type of malnutrition. The primary goal of nutritional therapy is to overcome this energy deficit.

Studies have shown variable intakes in the CF population, but the intakes are usually less than adequate and are associated with a less than normal growth pattern.

Asthma is a chronic inflammatory disorder of the airways, which can cause recurrent episodes of wheezing, breathlessness, chest tightness, and coughing of variable severity. Persistent asthma requires daily use of medication, preferably inhaled anti-inflammatory agents. Severe forms of asthma may require long-term use of oral corticosteroids which can result in growth suppression in children, poor bone mineralization, high weight gain, and, in pregnancy, decreased birthweight of the infant. High doses of inhaled corticosteroids can result in growth suppression in children and poor bone mineralization. Untreated asthma is also associated with poor growth and bone mineralization and, in pregnant women, adverse birth outcomes such as low birth weight, prematurity, and cerebral palsy. Repeated asthma exacerbations ("attacks") can, in the short-term, interfere with eating, and in the long-term, cause irreversible lung damage that contributes to chronic pulmonary disease. Compliance with prescribed medications is considered to be poor. Elimination of environmental factors that can trigger asthma exacerbations (such as cockroach allergen or environmental tobacco smoke) is a major component of asthma treatment. WIC can help by providing foods high in calcium and vitamin D, in educating participants to consume appropriate foods and to reduce environmental triggers, and in supporting and encouraging compliance with the therapeutic regimen prescribed by the health care provider.

NOTE: This criterion will usually not be applicable to infants for the medical condition of asthma. In infants, asthma-like symptoms are usually diagnosed as bronchiolitis with wheezing which is covered under Criterion #352, Infectious Diseases.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis (“My doctor says that I have/my son or daughter has…”) Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Institute of Medicine: WIC Nutrition Risk Criteria: A Scientific Assessment; 1996; pp. 185-187, 190-191
2. Queen, Patricia and Lang, Carol: Handbook of Pediatric Nutrition; 1993; pp. 422-425.
3. National Heart, Lung, and Blood Institute: Expert Panel Report 2: Guidelines for the Diagnosis and Management of Asthma; 1997; pp. 3, 20, 67-73.
4. National Heart, Lung, and Blood Institute: Management of Asthma During Pregnancy; 1992; pp. 7, 36-37.
5. JAMA: Asthma Information Center: Asthma Medications Misused, Underused in Inner City Residents; 1998, pp.1-2.

Nutrition Risk Criteria **361 DEPRESSION**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Presence of clinical depression, including postpartum depression.

Presence of condition diagnosed, documented, or reported by a physician, clinical psychologist, or someone working under a physician's orders, or as self-reported by applicant/participant/caregiver. See the **Clarification** section for more information about self-reporting a diagnosis.

Required Documentation

Document source of information (i.e. self-report of diagnosis by medical provider or written documentation by medical provider).

Justification

According to the National Institute of Mental Health (NIMH) nearly 10% of the U.S. population ages 18 and older suffers from depression each year, with 6.7% suffering from major depressive disorders (1). Although depression can occur at any age, the average onset is around age 30 (1, 2). Depression occurs twice as frequently in women as in men. Depression has a variety of symptoms, but the most common are deep feelings of sadness or a marked loss of interest in pleasure or activities. Other symptoms of depression include: appetite changes resulting in unintended weight losses or gains, insomnia or oversleeping, loss of energy or increased fatigue, restlessness or irritability, feelings of worthlessness or inappropriate guilt and difficulty thinking, concentrating or making decisions (1-3). Further, depression can increase the risk for some chronic diseases such as coronary heart disease, myocardial infarction, chronic pain syndromes, premature aging, and impaired wound healing. Therefore, untreated depression has the potential to impact long term health status (4). For information about children and depression, please see the **Clarification** section.

Pregnancy and Depression

Depression is common during pregnancy. Between 14% and 23% of pregnant women will experience depressive symptoms (5, 6). Several studies have found that depression risk is highest during the last trimester of pregnancy (4). Women who experience depression during pregnancy are found to be less likely to seek prenatal care (3). They may also suffer from episodes of nausea/vomiting or initiate/increase the use of drugs, alcohol and nicotine (4). Pregnant women with depression may be at risk for preeclampsia, preterm delivery or delivery of low birth weight infants and have higher perinatal mortality rates (5, 6).

Pregnant Adolescents

In the United States, 10% of women became pregnant during adolescence (7). The prevalence of teen pregnancy is highest among African and Native Americans, lower socioeconomic groups, and those living in stressful family

environments. The prevalence rate of depression among pregnant adolescents is between 16% and 44%, which is almost twice as high as among their adult counterparts and non-pregnant adolescents (7). Adolescence is a stage of rapid metabolic, hormonal, physiological and developmental changes. Depressive symptoms are likely to emerge when the physiologic and psychological changes that occur during pregnancy are superimposed upon normal developmental change. (8)

Teens who are under stress, lack social and/or family support, experience significant loss, or who have attention, learning or conduct disorders are at greater risk for developing clinical depression (9). Depression in young people often occurs with mental disorders, substance abuse disorders, or physical illnesses, such as diabetes (10). Pregnant adolescents with depressive symptoms are more likely to delay or refuse prenatal care and have subsequent, short interval pregnancies (within 24 months), both of which have shown to result in poor pregnancy outcomes (11, 12).

Antidepressant Use in Pregnancy

Negative consequences for the newborn such as fetal growth changes and shorter gestation periods have been associated with both depression symptoms and use of antidepressant medications during pregnancy. Although rare, some studies have linked fetal malformations, cardiac defects, pulmonary hypertension and reduced birth weight to antidepressant use during pregnancy, however, more research in this area is needed. (4, 6, 13) For more information about specific drug therapies used for treating depression, please see the Clarification section (14).

A fetus exposed to antidepressants throughout pregnancy or during the last trimester may, in rare instances, experience temporary withdrawal symptoms – such as jitters or irritability – at birth (15, 16). Some health care providers may suggest tapering dosages until after birth to minimize newborn withdrawal symptoms though it is unclear whether this method can reduce harmful effects. This strategy may also be unsafe for new mothers as they enter the postpartum period – a time of increased risk of mood swings and problems with anxiety. Therefore, it is imperative that prenatal women discuss the risks and benefits of antidepressant therapy with their health care provider.

Postpartum Depression and Related Mood Disorders

Postpartum depression was historically hypothesized to be caused by low estrogen and progesterone levels immediately following birth, however, this hypothesis has been found to have limited scientific support (17). Emerging studies have found that reproductive hormones have an indirect relationship on depression because of the influence on stress hormones, immune markers or sleep quality. The incidence of postpartum depression in new mothers can range from approximately 12% to 25%, to up to 35% or more in some high-risk groups. High risk groups include: women of low income, younger age, low education level and histories of stressful life events or traumatic experiences. Some studies have higher percentage rates for depression because they include both subjects with diagnosed major depression and those with depressive symptoms, thus accounting for the wide range in rates. (4)

Postpartum depression is distinguished from “baby blues” – a common reaction following delivery – both by its duration and the debilitating effects of the indifference the mother has about herself and her children (17). “Baby blues” are

characterized by mild depressive symptoms, tearfulness (often for no discernible reason), anxiety, irritableness, mood fluctuations, increased sensitivity and fatigue. The “blues” typically peak four to five days after delivery, may last hours to days and resolve by the 10th postnatal day (18).

Inflammation and Depression

Inflammation was once recognized as one of several risk factors for depression. New research has found that inflammation is not a risk factor – but rather it is the risk factor that underlies all others. This represents a shift in how inflammation contributes to depression. Emerging research has revealed that depression is associated with inflammation manifested by increased levels of proinflammatory cytokines. Common experiences of new motherhood; sleep disturbance, postpartum pain and past or current psychological trauma, act as stressors that cause proinflammatory cytokine levels to rise. This finding may explain why psychological trauma, act as stressors that cause proinflammatory cytokine levels to rise. This finding may explain why psychosocial, behavioral and physical risk factors increase the risk of depression (19). Additionally, inflammation levels normally rise during the last trimester of pregnancy, which may explain, as stated in the **Pregnancy and Depression** section above, the higher risk for experiencing depression during pregnancy (4).

Breastfeeding and Depression

Successful breastfeeding has a protective effect on maternal mental health because it attenuates stress and modulates the inflammatory response. Conversely, breastfeeding difficulties such as nipple pain can increase the risk of depression and should be addressed promptly. (19)

Implications for WIC Nutrition Services

Individuals diagnosed with depression can benefit from WIC nutrition services and supplemental foods. Through participant-centered counseling, WIC staff can, as necessary:

- Reinforce and support the treatments and therapies prescribed by the participant’s health care provider.
- Make referrals to the primary health care provider and/or to other appropriate mental health and social service programs. A 2010 brief from the Urban Institute, recognized the WIC Program as a viable access point to identify and refer mothers with depressive symptoms (20). To learn more about mental health resources in your area please access the U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration’s website. http://store.samhsa.gov/mhlocator or http://www.samhsa.gov/prevention/.
- Provide follow-up to ensure that the woman is receiving the necessary mental health treatment.
- Encourage food choices that promote nutritional well-being (to include good sources of Omega-3’s for their anti-inflammatory properties).
- Educate about the increased risk of depressive symptoms during the third trimester of pregnancy as well as the prevalence, risks and signs of postpartum depression.

- Provide adequate breastfeeding education, assessment and support (e.g., peer counseling) to women with existing depression; both prenatally and in the postpartum period.

A supplement to this criterion was developed to provide WIC State and local agencies with more information about the treatment of depression and WIC's role in providing nutrition services to women at risk of o diagnosed with depression: ***Guidance for Screening and Referring Women with or At Fisk for Depression.***

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Depression may be present in young children; however, it is generally not diagnosed until later in life. At this time, there is no evidenced-based research to support the diagnosis of depression as a risk criterion for WIC children participants. It is important to note, however, that a child's health may be at risk if the mother has a diagnosis of depression.

Nutrition Risk Criterion ***902 Woman of Infant/Child of Primary Caregiver with Limited ability to Make Feeding Decisions or Prepare Food***, is an appropriate risk criterion assignment for an infant or child of a WIC mother diagnosed with clinical depression.

There are three major classes of antidepressants. Of the three classes listed below, the first two, Tricyclic antidepressants (TCAs) and Selective serotonin reuptake inhibitors (SSRIs) are generally viewed as safe options for pregnant and breastfeeding women. MAOIs such as Nardil (Phenelzine) and Parnate (Tranylcypromine) are always contraindicated during pregnancy and breastfeeding as reproductive safety has not been established. (20)

- **Tricyclic antidepressants (TCAs)** are the oldest, least expensive and most studied of the antidepressants with a proven track record of effectiveness and include medications such as Amitriptyline (Elavil) and Desiprmine (Norpramin). Noted drawbacks are complex dosing, unpleasant side effects and risk of suicide.
- **Selective serotonin reuptake inhibitors (SSRIs)** are used most frequently in pregnant and breastfeeding mothers. Sertraline (Zoloft) are paroxetine (Paxil) are recommended first line treatments for breastfeeding women due to fewer side effects than other antidepressants and a once-a-day dosing schedule. Paroxetine (Paxil) is generally discouraged during pregnancy because it has been associated with fetal heart defects when taken during the first three months of pregnancy. Infants of mothers on these medications should be monitored for the following symptoms: sedation, agitation, irritability, poor feeding and GI distress.
- **Monoamine oxidase inhibitors (MAOIs)** work by inhibiting the enzyme monoamine oxidase to allow for more norepinephrine and serotonin to remain available in the brain. As stated above, these types of medications are always contraindicated during pregnancy and

breastfeeding as reproductive safety has not been established. Furthermore, MAOIs have many drug and diet contraindications.

Nutrition Risk Criterion ***#357 Drug-Nutrient Interactions*** may be assigned, as appropriate, to women taking antidepressants.

Reference

1. National Institute of Mental Health [Internet]. Major depressive disorders among adults. Available at: http://www.nimh.nih.gov/statistics/1MDD_ADULT.shtml. Accessed June 2013.
2. American Psychiatric Association. Consumer Fact Sheet: Let's talk facts about depression. Available at: www.healthyminds.org. Accessed June 2013.
3. Institute of Medicine. WIC nutrition risk criteria: a scientific assessment. National Academy Press, Washington, D.C.; 1996.
4. Kendall-Tackett K. Depression in new mothers: causes, consequences and treatment alternatives, 2nd edition: 2010 pages 17-19, 8, 51-58.
5. Gaynes BN, Gavin N, Meltzer-Brody S, Lohr KN, Swinson T, Gartlehner G, Brody S, Miller WC. Perinatal depression: prevalence, screening accuracy, and screening outcomes. Evidence Report/Technology Assessment No. 119. (Prepared by the RTI-University of North Carolina Evidence-based Practice Center, under Contract No. 290-02-0016.) AHRQ Publication No. 05-E006-2. Rockville, MD: Agency for Healthcare Research and Quality. February 2005.
6. Yonkers KA, Wisner KL, Stewart DE, et al. The management of depression during pregnancy: a report from the American Psychiatric Association and the American College of Obstetricians and Gynecologists September/October 2009.
7. Szigethy EM, Ruiz, P. Depression among pregnant adolescents: an integrated treatment approach. Am J Psychiatry, 2001 Jan; 158(1):22-7.
8. McClanahan KK. Depression in pregnant adolescents: considerations for treatment. J Pediatr Adolesc Gyneco. Feb; 22(1):59-64. 2009.
9. Cash RE, P. N. (2204, February). Depression in children and adolescents: information for parents and educators. Available at: http://www.nasponline.org/resources/handouts/revisedPDFs/depression.pdf. Accessed June 2013.
10. National Institute of Mental Health. Depression in children and adolescents. (Fact Sheet for Physicians). Bethesda, MD: Author (NIH Publication No. 00-4744). August 2001.
11. American Academy of Child and Adolescent Psychiatry. Facts for families: when children have children. July, 2004 (No.31). Available at: www.aacap.org. Accessed June 2013.
12. Barnet B, Liu J, Devoe M. Double jeopardy: depressive symptoms and rapid subsequent pregnancy in adolescent mothers. Arch Pediatric Adolescent Medicine; 162(3), March 2008.
13. Kendall-Tackett K, Hale TW. Review: The use of antidepressants in pregnant and breastfeeding women: a review of recent studies. Hum Lact May 2010 26: 187-195, first published on August 3, 2009.
14. Mayo Clinic. Antidepressants: safe during pregnancy? Last updated 1/10/12. Available at: www.mayoclinic.com/health/antidepressants/DN00007/NSECTIONG. Accessed June 2013.
15. Sanz E, De-las-Cuevas C, Kiuru A, Bate A, Edwards R. Selective serotonin reuptake inhibitors in pregnant women and neonatal withdrawal syndrome: a database analysis. Lancet 2005, 365:451-4.

16. Hale TW, Kendall-Tackett K, Cong Z, Votta R, McCurdy F. Discontinuation syndrome in newborns whose mothers took antidepressants while pregnant or breastfeeding. Breastfeed Med. 2010 Dec;5(6):283-8. Epub 2010 Aug 31.
17. National Alliance on Mental Health Women and Depression Fact Sheet. Reviewed by Frank E, PhD; Novick D, Masalehdan A. November 2003; Updated October 2009, Reviewed by Duckworth K, MD. October 2009. Available at: http://www.nami.org/Template.cfm?Section=Women_and_Depression&Template=/ContentManagement/ContentDisplay.cfm&ContentID=89194. Accessed June 2013.
18. Epperson CN. Postpartum major depression: detection and treatment. American Family Physician. April 15, 1999. Available at: http://www.aafp.org/afp/990415ap/2247.html. Accessed June 2013.
19. Kendall-Tackett K. A new paradigm for depression in new mothers: the central role of inflammation and how breastfeeding and anti-inflammatory treatments protect mental health, International Breastfeeding Journal. March 2007.
20. Vericker T, Macomber J, Golden O. Infants of depressed mothers living in poverty: opportunities to identify and serve. Urban Institute, Brief 1, August 2010. Available at: http://www.urban.org/uploadedpdf/412199-infnats-of-depressed.pdf. Accessed June 2013.

Additional References

1. Healthwise.org [Internet]. Alcohol and drug problems. Available at: http://health.msn.com/health-topics/addiction/alcohol-and-drug-problems-61. Accessed June 2013.
2. U.S. Department of Health and Human Services. Health Resources and Services Administration, Maternal and Child Health Bureau. Depression during & after pregnancy: a resource for women, their families & friends. Avaialbe at: http://www.mchb.hrsa.gov/pregnancyandbeyond/depression/index.html. Accessed June 2013.

Nutrition Risk Criteria **362 DEVELOPMENTAL, SENSORY, OR MOTOR DELAYS INTERFERING WITH THE ABILITY TO EAT**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD
INFANTS	1	RD
CHILDREN	3	RD

Definition

Developmental, sensory or motor disabilities that restrict the ability to intake, chew or swallow food or require tube feeding to meet nutritional needs. Disabilities include but are not limited to:

- minimal brain function
- feeding problems due to a developmental disability such as pervasive development disorder (PDD) which includes autism
- birth injury
- head trauma
- brain damage
- other disabilities

Required Documentation

Developmental, sensory or motor disability

Justification

Infants and children with developmental disabilities are at increased risk for nutritional problems. Education, referrals, and service coordination with WIC will aid in early intervention of these disabilities. Prenatal, lactating and non-lactating women with developmental, sensory or motor disabilities may: 1) have feeding problems associated with muscle coordination involving chewing or swallowing, thus restricting or limiting the ability to consume food and increasing the potential for malnutrition; or 2) require enteral feedings to supply complete nutritional needs which may potentially increase the risk for specific nutrient deficiencies.

Pervasive Developmental Disorder (PDD) is a category of developmental disorders with autism being the most severe. Young children may initially have a diagnosis of PDD with a more specific diagnosis of autism usually occurring at 2½ to 3 years of age or older. Children with PDD have very selective eating habits that go beyond the usual "picky eating" behavior and that may become increasingly selective over time, i.e., foods they used to eat will be refused. This picky behavior can be related to the color, shape, texture or temperature of a food. Common feeding concerns include:

- difficulty with transition to textures, especially during infancy;
- increased sensory sensitivity; restricted intake due to color, texture, and/or temperature of foods;
- decreased selection of foods over time;

- difficulty accepting new foods; difficulty with administration of multivitamin/mineral supplementation and difficulty with changes in mealtime environment.

Nutrition education, referrals, and service coordination with WIC will assist the participant, parent or caregiver in making dietary changes/adaptations and finding assistance to assure she or her infant or child is consuming a nutritionally adequate diet.

References

1. Quinn, Heidi Puelzl; "Nutrition Concerns for Children With Pervasive Developmental Disorder/Autism" published in Nutrition Focus by the Center on Human Development and Disability; University of Washington, Seattle, Washington; September/October 1995.
2. Paper submitted by Betty Lucas, MPH, RD, CD to the Risk Identification and Selection Collaborative (RISC); November, 1999.
3. Zeman. Frances J.; Clinical Nutrition and Dietetics, 2nd Edition; 1991; pp.713-14, 721-22, 729-730.

Nutrition Risk Criteria **363** **PRE-DIABETES**

Category, Priority and Referral

Category	Priority	Referral
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD

Definition

Impaired fasting glucose (IFG) and/or impaired glucose tolerance (IGT) are referred to as pre-diabetes. These conditions are characterized by hyperglycemia that does not meet the diagnostic criteria for diabetes mellitus (1). (See Clarification for more information.)

Presence of pre-diabetes diagnosed by a physician as self-reported by application/participant/caregiver, or as reported or documented by a physician or someone working under a physician's orders.

Required Documentation

Document source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

An individual who is identified as having pre-diabetes is at relatively high risk for the development of Type 2 diabetes and cardiovascular disease (CVD).

The Expert Committee on the Diagnosis and Clarification of Diabetes Mellitus (2, 3) recognized a group of individuals whose glucose levels, although not meeting criteria for diabetes, are nevertheless too high to be considered normal. The blood tests used to measure plasma glucose and to diagnose pre-diabetes include a fasting plasma glucose test and a glucose tolerance test (see Clarification for more information). Individuals with a fasting plasma glucose level between 100-125 mg/dl are referred to as having impaired fasting glucose (IFG). Individuals with plasma glucose levels of 140-199 mg/dl after a 2-hour oral glucose tolerance test are referred to as having impaired glucose tolerance (IGT).

Many individuals with IGT are euglycemic and, along with those with IFG, may have normal or near normal glycosylated hemoglobin (HbA1c) levels. Oftentimes, individuals with IGT manifest hyperglycemia only when challenged with the oral glucose load used in standardized oral glucose tolerance tests.

The prevalence of IFG and IGT increases greatly between the ages of 20 and 49 years. In people who are > 45 years of age and overweight (BMI ≥ 25), the prevalence of IFG is 9.3% and 12.8% for IGT.

Screening for pre-diabetes is critically important in the prevention of Type 2 diabetes. The American Diabetes Association recommends (5) that testing to detect pre-diabetes should be considered in all asymptomatic adults who are overweight (BMI ≥ 25) or obese (BMI ≥ 30) and who have one or more additional risk factors (see Table 1 in Clarification).

IFG and IGT are not clinical entities in their own right, but rather risk factors for future diabetes as well as CVD. (Note: During pregnancy, IFG and IGT are diagnosed as gestational diabetes.) They can be observed as intermediate

stages in many of the disease processes. IFG and IGT are associated with the metabolic syndromes, which include obesity (especially abdominal or visceral obesity), dyslipidemia (the high-triglyceride and/or low HDL type), and hypertension. Dietary recommendations include monitoring of calories, reduced carbohydrate intake and high fiber consumption. Medical nutrition therapy (MNT) aimed and producing 5-10% loss of body weight and increased exercise have been variably demonstrated to prevent or delay the development of diabetes in people with IGT. However, the potential impact of such interventions to reduce cardiovascular risk has not been examined to date (2, 3).

WIC nutrition services can support and reinforce the MNT and physical activity recommendations that participants receive from their health care providers. In addition, WIC nutritionists can play an important role in providing women with counseling to help them achieve or maintain a healthy weight after delivery.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has...") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

Hyperglycemia is identified through a fasting blood glucose or an oral glucose tolerance test (1).

Impaired fasting glucose (IFG) is defined as fasting plasma glucose (FPG) ≥ 100 or ≥ 125 mg/dl (≥ 5.6 or ≥ 6.1 mmol/l), depending on study and guidelines (2).

Impaired glucose tolerance (IGT) is defined as a 75-g oral glucose tolerance test (OGTT) with 2-h plasma glucose values of 140-199 mg/dl (7.8-11.0 mmol/l).

The cumulative incidence of diabetes over 5-6 years was low (4-5%) in those individuals with normal fasting and normal 2-h OGTT values, intermediate (20-34%) in those with IFG and normal 2-h OGTT of IGT and a normal FPG, and highest (38-65%) in those with combined IFG and IGT (4).

Recommendations for testing for pre-diabetes and diabetes in asymptomatic, undiagnosed adults are in Table 1 below.

Table 1. Criteria and Methods for Testing for Pre-Diabetes and Diabetes in Asymptomatic Adults

1. Testing should be considered in all adults who are overweight (BMI ≥ 25*) and have additional risk factors:
    - physical inactivity
    - first-degree relative with diabetes
    - members of a high-risk ethnic population (e.g., African American, Latino, Native American, Asian American, Pacific Islander)
    - women who delivered a baby weighing > 9 lbs or were diagnosed with gestational diabetes mellitus
    - hypertension (blood pressure ≥ 140/90 mmHg or on therapy for hypertension)
    - women with polycystic ovarian syndrome (PCOS)
    - IGT or IFG on previous testing

- other clinical conditions associated with insulin resistance (e.g., severe obesity and acanthosis nigricans)
- history of CVD

2. In the absence of the above criteria, testing for pre-diabetes should begin at age 45 years.

3. If results are normal, testing should be repeated at least at 3-year intervals, with consideration of more frequent testing depending on initial results and risk status.

4. To test for pre-diabetes or diabetes, either an FPG test or 2-hour glucose tolerance (OGTT; 75 g glucose load), or both, is appropriate.

5. An OGTT may be considered in patients with impaired fasting glucose (IFG) to better define the risk of diabetes.

6. In those identified with pre-diabetes, identify and if appropriate, treat other CVD risk factors.

* At-risk BMI may be lower in some ethnic groups.

Reference

1. American Diabetes Association. Clinical practice recommendations: Standards of medical care in diabetes. Diabetes Care. 2008 Jan: 31 Suppl 1:S12-54.
2. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the expert committee on the diagnosis and classification of diabetes mellitus. Diabetes Care. 1997; 20:1183-1197.
3. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Follow-up report on the diagnosis of diabetes mellitus. Diabetes Care. 2003; 26:3160-3167.
4. American Diabetes Association National Institute of Diabetes and Digestive and Kidney Diseases. Position statement on prevention or delay of type 2 diabetes. Diabetes Care. 2004; 27:S47.
5. American Diabetes Association. Executive summary: Standards of medical care in diabetes. Diabetes Care. 2008 Jan; 31 Suppl 1:S5-11.

Nutrition Risk Criteria **371 MATERNAL SMOKING**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	-

Definition

Any daily smoking of tobacco products, i.e., cigarettes, pipes, or cigars.

Required Documentation

Smoking

Justification

Research has shown that smoking during pregnancy causes health problems and other adverse consequences for the mother, the unborn fetus and the newborn infant such as: pregnancy complications, premature birth, low-birth-weight, stillbirth, infant death, and risk for Sudden Infant Death Syndrome (SIDS) (1). Women who smoke are at risk for chronic and degenerative diseases such as: cancer, cardiovascular disease and chronic obstructive pulmonary disease. They are also at risk for other physiological effects such as loss of bone density (2).

Maternal smoking exposes the infant to nicotine and other compounds, including cyanide and carbon monoxide, in-utero and via breastmilk (3). In-utero exposure to maternal smoking is associated with reduced lung function among infants (4). In addition, maternal smoking exposes infants and children to environmental tobacco smoke (ETS). (See #904, Environmental Tobacco Smoke.)

Because smoking increases oxidative stress and metabolic turnover of vitamin C, the requirement for this vitamin is higher for women who smoke (5). The WIC food package provides a good source of vitamin C. Women who participate in WIC may also benefit from counseling and referral to smoking cessation programs.

References

1. Manual of Clinical Dietetics 6th ed., American Dietetic Association. 2000.
2. Women and Smoking: A Report of the Surgeon General – 2001. www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/sgr_women_chapters.htm.
3. Breastfeeding Handbook for Physicians, American Academy of Pediatrics and American College of Obstetrics and Gynecologists. 2006
4. U.S. Department of Health and Human Services. *The Health Consequences of Smoking:* A Report of the Surgeon General—Executive Summary. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2004.
5. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids* (2000). Institute of Medicine, the National Academy of Science.

Nutrition Risk Criteria **372 ALCOHOL AND ILLEGAL DRUG USE**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD
BREASTFEEDING WOMEN	1	RD
NON-BREASTFEEDING WOMEN	6	RD

Definition

For Pregnant Women:

- Any alcohol use
- Any illegal drug use

For Breastfeeding and Non-breastfeeding Postpartum Women:

- Routine current use of ≥ 2 drinks per day (6). A serving or standard sized drink is 1 can of beer (12 fluid oz.); 5 oz wine; and 1½ fluid oz liquor (1 jigger gin, rum, vodka, whiskey (86 proof), vermouth, cordials or liqueurs)
- Binge drinking, i.e. drinks 5 or more (≥ 5) drinks on the same occasion on at least one day in the past 30 days
- Heavy drinking, i.e. drinks 5 or more (≥ 5) drinks on the same occasion on five or more days in the previous 30 days
- Any illegal drug use

Required Documentation

Alcohol/drug use

Justification

Drinking alcoholic beverages during pregnancy can damage the developing fetus. Excessive alcohol consumption may result in low birth weight, reduced growth rate, birth defects, and mental retardation. WIC can provide supplemental foods, nutrition education and referral to medical and social services which can monitor and provide assistance to the family.

“Fetal Alcohol Syndrome” is a name given to a condition sometimes seen in children of mothers who drank heavily during pregnancy. The child has a specific pattern of physical, mental, and behavioral abnormalities. Since there is no cure, prevention is the only answer.

The exact amount of alcoholic beverages pregnant women may drink without risk to the developing fetus is not known as well as the risk from periodic bouts of moderate or heavy drinking. Alcohol has the potential to damage the fetus at every stage of the pregnancy. Therefore, the recommendation is not to drink any alcoholic beverages during pregnancy.

Studies show that the more alcoholic beverages the mother drinks, the greater the risks are for her baby. In addition, studies indicate that factors such as cigarette smoking and poor dietary practices may also be involved. Studies show that the reduction of heavy drinking during pregnancy has benefits for both mother and newborns. Pregnancy is a special time in a woman's life and the majority of heavy drinkers will respond to supportive counseling.

Heavy drinkers themselves may develop nutritional deficiencies and more serious diseases, such as cirrhosis of the liver and certain types of cancer, particularly if they also smoke cigarettes. WIC can provide education and referral

to medical and social services, including addiction treatment, which can help improve pregnancy outcome.

Pregnant women who smoke marijuana are frequently at higher risk of still birth, miscarriage, low birth weight babies and fetal abnormalities, especially of the nervous system. Heavy cocaine use has been associated with higher rates of miscarriage, premature onset of labor, IUGR, congenital anomalies, and developmental/behavioral abnormalities in the preschool years. Infants born to cocaine users often exhibit symptoms of cocaine intoxication at birth. Infants of women addicted to heroin, methadone, or other narcotics are more likely to be stillborn or to have low birth weights. These babies frequently must go through withdrawal soon after birth. Increased rates of congenital defects, growth retardation, and preterm delivery, have been observed in infants of women addicted to amphetamines.

Pregnant addicts often forget their own health care, adding to their unborn babies' risk. One study found that substance abusing women had lower hematocrit levels at the time of prenatal care registration, lower pregravid weights and gained less weight during the pregnancy. Since nutritional deficiencies can be expected among drug abusers, diet counseling and other efforts to improve food intake are recommended.

Heroin and cocaine are known to appear in human milk. Marijuana also appears in a poorly absorbed form but in quantities sufficient to cause lethargy, and decreased feeding after prolonged exposure.

References

1. USDA/DHHS Dietary Guidelines; 1995.
2. Lawrence Ruth: Maternal & Child Health Technical Information Bulletin: A Review of Medical Benefits and Contraindications to Breastfeeding in the United States; October 1997.
3. Weiner, L., Morse, B.A., and Garrido, P.: FAS/FAE Focusing Prevention on Women at Risk; International Journal of the Addictions; 1989; 24:385-395.
4. National Clearinghouse for Alcohol and Drug Information; Office for Substance Abuse Prevention; The fact is …alcohol and other drugs can harm an unborn baby; Rockville; 1989.
5. Institute of Medicine. Nutrition during pregnancy. National Academy Press, Washington, D.C.; 1990.
6. Jones, C. and Lopez, R.: Drug Abuse and Pregnancy; New Perspectives in Prenatal Care; 1990; pp. 273-318.
7. National Household Survey on Drug Abuse, Main Findings 1996; Office of Applied Studies, Substance Abuse and Mental Health services Administration. DHHS.

Nutrition Risk Criteria **381 ORAL HEALTH CONDITIONS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	RD/DENTAL
BREASTFEEDING WOMEN	1	DENTAL
NON-BREASTFEEDING WOMEN	6	DENTAL
INFANTS	1	DENTAL
CHILDREN	3	DENTAL

Definition

Oral health conditions include, but are not limited to:

- Dental caries, often referred to as “cavities” or “tooth decay”, is a common chronic, infectious, transmissible disease resulting from tooth-adherent specific bacteria, that metabolizes sugars to produce acid which, over time, demineralizes tooth structure (1).

- Periodontal diseases are infections that affect the tissues and bone that support the teeth. Periodontal diseases are classified according to the severity of the disease. The two major stages are gingivitis and periodontitis. Gingivitis is a milder and reversible form of periodontal disease that only affects the gums. Gingivitis may lead to more serious, destructive forms of periodontal disease called periodontitis. (2)

  More information on types of periodontal disease is available at: http://www.perio.org/consumer/2a.html.

- Tooth loss, ineffectively replaced teeth or oral infections which impair the ability to ingest food in adequate quantity or quality.

Presence of oral health conditions diagnosed, documented, or reported by a physician, dentist, or someone working under a physician’s orders, or as self-reported by applicant/participant/caregiver. See Clarification for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician/dentist or written documentation by physician/dentist).

Justification

Oral health reflects and influences general health and well-being. Good oral health care and nutrition during pregnancy, infancy and childhood are often overlooked factors in the growth and development of the teeth and oral cavity.

Infants and Children

The Centers for Disease Control and Prevention (CDC) reports that dental caries may be the most prevalent infectious disease in U.S. children. More than 40% of children have tooth decay by the time they reach kindergarten. Infants that consume sugary foods, are of low socioeconomic status, and whose mothers have a low education level, are 32 times more likely to have caries at the age of 3 years than children who do not have those risk factors. Despite its high prevalence, early childhood caries (ECC) is a preventable disease. (3)

ECC may develop as soon as teeth erupt. Bacteria, predominantly mutans streptococci (MS), metabolize simple sugars to produce acid that demineralizes teeth, resulting in cavities. The exact age at which MS colonization occurs in children is controversial, but it does not happen until teeth erupt. The earlier colonization occurs, the greater the risk of carries. MS typically originates in the mother and is transmitted to the child via saliva (often through cup and utensil sharing). Elevated maternal levels of MS, due to active or untreated caries and frequent sugar consumption, increase the risk of transmission. In addition, recent evidence suggests that exposure to environmental tobacco smoke increases the likelihood of MS colonization in children. (4)

Historically, ECC has been attributed to inappropriate and prolonged bottle use; formally called “baby bottle tooth decay.” However, recent studies indicate that the disease is multifactorial, which suggests any feeding practice that allows frequent sugar consumption in the presence of MS may result in caries formation: propped bottles containing sweetened liquids or formula, frequent consumption of juice or sweetened liquids from infant and “sippy” cups, and frequent snacking of high cariogenic foods. (4)

The frequency of sugar consumption is the main dietary variable in caries etiology. After bacteria metabolize sugar into acid, it takes 20-40 minutes for the acid to be neutralized or washed away by saliva. Therefore, if sugars are frequently consumed, the potential for demineralization is greater. Although MS can metabolize many different carbohydrates, they produce acid most efficiently from sugars, especially sucrose. Sugars within the cellular structure of food (such as fructose in whole fruit) are thought to be less cariogenic than sugars intentionally added to foods. (4) See **Table 1** for more information on the cariogenic potential of children’s foods and snacks.

Milk is widely consumed, especially by children, and thus the interaction between different kinds of milk consumed and caries development has been a research topic of interest. Lactose is one of the least cariogenic sugars because it is poorly metabolized by MS. Researchers have reported cows’ milk to be a protective, anticariogenic agent due to its high concentration of calcium and phosphate. The buffering activity of proteins present in cows’ milk also might allow the formation of very stable complexes of calcium phosphate. Other anticariogenic properties in cows’ milk include antibacterial enzymes, vitamin D and fluoride. (4, 5)

Infant formulas, on the other hand, have a high potential for inducing caries due to their high carbohydrate variability. The cariogenic potential of breast milk is inconclusive. Breast milk has been found to contain more lactose (8.3%) than cows’ milk (4.9%). A higher breast milk lactose concentration and the possibility that lactose fermentation of cows’ milk is slower than in breast milk, may make breast milk caries risk slightly higher. Some evidence indicates that breastfeeding for over 1 year during the night after tooth eruption might be associated with ECC, however other investigations showed no relationship between prevalence of caries and breastfeeding. Regardless of the type of milk consumed, sufficient dental care and cleaning after drinking milk/formula and breastfeeding can help prevent ECC. Avoiding inappropriate dietary practices, such as frequent juice consumption or snacking on highly cariogenic foods also remain important ECC preventive practices. (4, 5)

Table 1. Cariogenic Potential of Children's Foods and Snacks		
Noncariogenic	**Low Cariogenicity**	**High Cariogenicity**
Cheese Chicken Cottage cheese Eggs Flavored club soda Nuts and seeds* Plain cow's milk (unflavored) Plain yogurt Popcorn* Seltzer Vegetables	Flavored milk Fresh fruits Whole grain products	Breakfast bars Cake Candies** Cookies Doughnuts Fruit juice Granola bars Pretzels Dried fruits Soda crackers Sweetened beverages Sweetened dry cereals
*Not appropriate for infants and toddlers due to potential choking problems. **Sticky candy and/or slowly eaten candy are extremely cariogenic.		
Adapted from: Faine, MP. Nutrition and oral health. In: Proceedings of Promoting Oral Health of Children with Neurodevelopmental Disabilities and Other Special Health Care Needs. May 4-5, 2001. Seattle, WA.		

Women

Maternal periodontal disease and dental caries may impact pregnancy outcome, and the offspring's risk of developing early and severe dental caries. Periodontal disease and caries may also increase the women's risk of atherosclerosis, rheumatoid arthritis and diabetes. These oral health problems are highly prevalent in women of childbearing age, particularly among low-income women and members of racial and ethnic minority groups. Socioeconomic factors, lack of resources to pay for care, barriers to access care, lack of public understanding of the importance of oral health and effective self-care practices all represent underlying reasons cited for observed inadequacies in oral health. (6)

Maternal periodontal disease, a chronic infection of the gingiva (gums) and supporting tooth structures, has been associated with preterm birth, low birth weight and development of preeclampsia (6, 7). Studies indicate that periodontal infection can result in placental-fetal exposure and, when coupled with a fetal inflammatory response, can lead to preterm delivery (7). Additionally, in a cohort of 164 young, minority, pregnant and postpartum women, the preterm/low birth weight rate was 5.4% lower among women who received periodontal treatment than those who did not receive treatment (7). In a case-control study, researchers found that preeclamptic patients were 3.5 times more likely to have periodontal disease than normotensive patients (6). (See nutrition risk criterion ***304 History of Preeclampsia*** for more information.)

Fluoride and Fluorosis

Use of fluorides for the prevention and control of caries is documented to be both safe and highly effective. Fluoride, a naturally occurring substance, has several caries-protective mechanisms of action, including enamel remineralization and altering bacterial metabolism to help prevent caries. Excessive intake of fluoride can cause dental fluorosis which is a change in the appearance of the tooth's enamel. In the U.S., fluorosis appears mostly in the very mild or mild form – as barely visible lacy white markings or spots. The severe form of dental fluorosis, staining and pitting of the tooth surface, is rare in the U.S. The CDC reports that 32% of American children have some form of dental fluorosis, with 2.45% of children having the moderate to severe stages. (8, 9, 10, 11)

Parents and caregivers may have questions and concerns about fluoride content in water supplies and in infant formula. Fluoridated water can be found in communities that supplement tap water with fluoride and it may also be found in well water. The CDC's *My Water's Fluoride* website: http://apps.nccd.cdc.gov/MWF/Index.asp, allows consumers in currently participating States to learn the fluoridation status of their water system.

All formula, including powdered, concentrate and ready-to-feed, contain fluoride, but most infant formula manufacturers ensure low levels of fluoride (8). WIC State and local agencies should refer caregivers of formula fed infants with questions regarding the use of fluoridated vs. non-fluoridated water to prepare infant formula to the infants' health care provider.

Dental Care and Anxiety

It is reported that 50% of the U.S. population does not seek regular dental care. Of the entire U.S. population, 8-15% has dental phobias. Dental fear can be directly learned from previous painful or negative experiences or indirectly learned from family, friends and the media. Negative portrayal of dentistry by these sources adds to an individual's anxiety. Anxiety and/or fear of dental procedures may prevent participants from seeking necessary dental care during high risk periods of the life cycle (e.g., pregnancy). Dental providers are learning to understand the causes of dental fear, have techniques to assess the level of fear and have modified treatments to accommodate patients with high anxiety levels. (12)

Oral Health Problems and Special Health Care Needs

The following special health care needs can increase the risk for oral health problems and can also make the overall effects of poor oral health more severe (13):

- **Prematurity and intrauterine malnutrition** – can have adverse effects on an individual's oral health. A study of infants who weighed <2000g at birth indicated more porous dental enamel and subsurface lesions. Infants with very low birth weights (<1500g) are more apt to have enamel defects of the primary teeth. Malnutrition in the first few months of life (when oral structures develop) can increase the risk for oral problems.
- **Gastroesophageal Reflux Disease (GERD)** – common among children with cerebral palsy, Down syndrome and other conditions. GERD can contribute to oral health problems. As acidic gastric contents are regurgitated, primary and permanent teeth can be eroded.
- **Failure to thrive and other problems with weight gain and growth** – frequent meals and snacks (which may contribute to caries

development) may be needed to maintain an adequate energy intake, or if mealtime is longer than usual, the demineralization period may exceed remineralization. Delayed weaning and children sipping on a bottle throughout the day, could also contribute to oral health problems.

- **Craniofacial malformations** – individuals with these malformations are at higher risk of developing oral problems. For example, children with cleft lip/palate disorders have more decayed, missing, and filed teeth than children without.
- **Compromised immune function** – individuals with AIDS or those who take immunosuppressive medications are more susceptible to oral infections such as candidiasis, viral infections, dental caries, and periodontal disease.
- **Down syndrome (Trisomy 21)** – individuals with Down syndrome often have delayed dental development*, may be missing permanent teeth, and may have under-developed teeth or teeth with thin enamel. In addition, the potential for eating problems and GERD make oral care for individuals with Down's especially important. (13)

*Delayed Tooth Eruption (DTE) is the emergence of a tooth into the oral cavity at a time that deviates significantly from norms established for different races, ethnicities, and sexes. Variation in the normal eruption of teeth is a common finding, but significant deviations from established norms should alert the clinician to further investigate the patient's health and development. Eruption depends on genetics, growth of the jaw, muscular action and other factors. DTE is seen in children with certain genetic disorders, particularly Down syndrome, and in children with general developmental delays that involve the oral musculature. Whenever DTE is generalized, the child should be examined for systemic diseases affecting eruption, such as endocrine disorders, organ failures, metabolic disorders, drugs and inherited disorders. (14) Additional information about tooth eruption is available at: http://www.ada.org/2930.aspx.

Dentate Status, Diet Quality and General Health

By the time individuals reach adulthood, the human mouth has progressed from 20 primary teeth to 32 permanent (adult) teeth (2). The extent to which tooth loss can adversely affect nutritional status is not completely known. However, diet quality tends to decline as the degree of dental impairment increases. Studies have shown that intake of vitamin A, fiber, calcium and other key nutrients decline as the number of teeth decline. In The Health Professionals study, participants without teeth had diets that contained fewer vegetables, less carotene and fiber, and more cholesterol, saturated fat, and calories than persons with 25 teeth or more (15). Despite the trend toward increased tooth retention throughout adult life in developed countries, 11% of adults aged 25 and older have lost all of their natural teeth. This number increases to 30% for people over age 65 and is even higher in those living in poverty. Loss of teeth is not a normal result of the aging process; the major cause of tooth loss is extractions resulting from dental caries and/or periodontal disease. (15)

To help prevent oral health problems from developing and ensure the best possible health and developmental outcomes, WIC staff can encourage participants and caregivers to:

Diet

- Breastfeed infants during the first year of life and beyond as mutually desired.

- Avoid having an infant/child sleep with a bottle. Any bottle taken to bed should contain only water. (See Risks 425.3 and 411.2)
- Gradually introduce a cup between 6 and 12 months of age, wean from the bottle by 12 months of age.
- Drink/provide only water and milk between meals.
- Limit sugary foods and drinks (if sweets are eaten, it's best to restrict to mealtimes.)
- Avoid carbonated beverages and juice drinks. (See Risk 425.2)
- Limit the intake of 100% fruit juice to no more than 4-6 ounces per day.
- Establish eating patterns that are consistent with the Dietary Guidelines for Americans and the infant feeding practice guidelines of the American Academy of Pediatrics.
- Consume/provide a varied, balanced diet during gestation and throughout childhood to set the stage for optimal oral health. (1, 3, 4, 15)

Oral Hygiene

- Wipe the gums of even a very small infant with a soft washcloth or soft toothbrush, even prior to tooth eruption, to establish a daily oral hygiene routine (17, 18).
- Brush teeth (including an infant's, as soon as teeth erupt) thoroughly twice daily (morning and evening) and floss at least once every day.
- Minimize saliva sharing activities (i.e., sharing a drinking cup and utensils). (1, 3, 4, 15)

Fluoride

- Use fluoride toothpaste approved by the American Dental Association ("pea-size" for 2-5 year olds and, "smear" for under the age of two and at moderate or high caries risk). (1)
- Rinse every night with an alcohol-free over-the-counter mouth rinse with 0.05% sodium fluoride (guidance for woman participant and caregiver only). (3)
- Contact the infant's (if formula fed) health care provider with questions regarding the use of local drinking water or bottled water to prepare infant formula. (3)
- Talk to the dentist about fluoride supplements. These may be of benefit in reducing dental decay for children living in fluoride-deficient areas (See Risk 411.11).
- Check if the public water systems have added fluoride at: http://apps.nccd.cdc.gov/MWF/Index.asp
- Access the following website for more information about fluoride: http://www.cdc.gov/fluoridation/safety.htm.

Referrals

- Establish a dental home within 6 months of eruption of the first tooth and no later than 12 months of age. (3)
- See a dentist for examinations (every 6 months) and/or restoration of all active decay as soon as possible. (WIC staff should provide dental referrals as necessary.)

Oral Health Resources/Handouts

- Summary of Pediatric Oral Health Anticipatory Guidance: http://www.aafp.org/afp/2004/1201/p2113.html#afp20041201p2113-t2.
- Table: Oral health and dietary management for mothers and children (see page 3 of pdf.) http://www.sciencedirect.com/science/article/pii/S0002822398000443.

- Oral Health Care During Pregnancy: A National Consensus Statement: http://www.mchoralhealth.org/materials/consensus_statement.html.
- Your Baby's Teeth: http://www.aafp.org/afp/2004/1201/p2121.html.
- Two Healthy Smiles: Tips to Keep You and Your Baby Healthy: http://www.mchoralhealth.org/PDFs/pregnancybrochure.pdf.
- Tips for Good Oral Health Care During Pregnancy: http://www.mchoralhealth.org/PDFs/Oral HealthPregnancyHandout.pdf.
- A Healthy Smile for Your Baby (English): http://www.mchoralhealth.org/PDFs/babybrochure.pdf.
- A Healthy Smile for Your Baby (Spanish): http://www.mchoralhealth.org/PDFs/babybrochure_sp.pdf.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Policy on Early Childhood Caries (ECC): Classifications, consequences and preventive strategies. Am Academy of Pediatric Dentistry (2011) 47-49.
2. ADA Dental 101: Taking care of your teeth and gums. American Dental Association, 2008: 1-28.
3. American Academy of Pediatrics Policy Statement: Oral health risk assessment timing and establishment of the dental home. Pediatrics, (2003) Vol 11, No 5: 1113-1116.
4. Douglass BDS, et al. A practical guide to infant oral health. America Family Physician, (2004) Vol 70, No11: 2113-2120.
5. Rocha Peres RC, et al. Cariogenic potential of cows' human and infant formula milks and effect of fluoride supplementation. British Journal of Nutrition (2009), 101, 376-382.
6. Boggess K, Edelstein B. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. Matern Child Health J. (2006) 10: S169-S174.
7. Bobetsis Y, et al. Exploring the relationship between periodontal disease and pregnancy complications. J Am Dent Assoc. Vol 137, Suppl 2, 7S-135, 2006.
8. Review Council, Guideline on Fluoride Therapy. American Academy of Pediatric Dentistry. Reference Manual 33/No6. 11/12 153-156.
9. J Berg, et al. Evidence-based clinical recommendations regarding fluoride intake from reconstituted infant formula and enamel fluorosis: a report of the American Dental Association Council on Scientific Affairs. JADA, Jan 2011 Vol 142 79-87.
10. Position of the Academy of Nutrition and Dietetics: the impact of fluoride on health. Journal of the Academy of Nutrition and Dietetics, Sept 2012, Vol 112, No. 9 1443-1453.
11. HHS Proposed Guidelines on Fluoride in Drinking Water, Commentary by Dr. Howard Koh, Jan 2011.
12. Krochak M, Rubin JG. An overview of the treatment of anxious and phobic dental patients. Compend Contin Educ Denta, Vol XIV, No 5: 604-614.
13. Ogata B, Trahms C. For children with special health care needs. Nutrition Focus Nov/Dec 2003 Vol 18, No 6.

14. Suri L, et al. Delayed Tooth Eruption: Pathogenesis, diagnosis, and treatment. A Literature Review Am J of Orthodontics, (Oct 2004) 432-445.
15. Palmer CA. Diet and nutrition in oral health, 2nd ed. Pearson Education Inc. 2007. Chapter 15, 315-330.
16. Policy Statement: Preventive oral health intervention for pediatricians. Pediatrics (2008). Vol 122, No 6: 1387-1394.
17. American Academy of Pediatrics. A Pediatric Guide to Oral Health Flip Chart and Reference Guide. 2011. Copyright 2012.
18. Holt K, et al. Bright Futures Nutrition, 3rd Edition. American Academy of Pediatrics, 2011.

Nutrition Risk Criteria **382 FETAL ALCOHOL SYNDROME**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD, SA
CHILDREN	3	RD, SA

Definition

Fetal Alcohol Syndrome (FAS) is based on the presence of retarded growth, a pattern of facial abnormalities, and abnormalities of the central nervous system, including mental retardation.

Presence of FAS diagnosed by a physician as self-reported by applicant/participant/caregiver; or as reported or documented by a physician, or someone working under physician's orders.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

FAS is a combination of permanent, irreversible birth defects attributable solely to alcohol consumption by the mother during pregnancy. There is no known cure; it can only be prevented. Symptoms of FAS may include failure to thrive, a pattern of poor growth throughout childhood and poor ability to suck (for infants). Babies with FAS are often irritable and have difficulty feeding and sleeping.

Lower levels of alcohol use may produce Fetal Alcohol Effects (FAE) or Alcohol Related Birth Defects (ARBD) that can include mental deficit, behavioral problems, and milder abnormal physiological manifestations. FAE and ARBD are generally less severe than FAS and their effects are widely variable. Therefore, FAE and ARBD in and of themselves are not considered risks, whereas the risk of FAS is unquestionable.

Identification of FAS is an opportunity to anticipate and act upon the nutritional and educational needs of the child. WIC can provide nutritional foods to help counter the continuing poor growth and undifferentiated malabsorption that appears to be present with FAS. WIC can help caregivers acknowledge that children with FAS often grow steadily but slower than their peers. WIC can also educate the caregiver on feeding, increased calorie needs and maintaining optimal nutritional status of the child.

Alcohol abuse is highly concentrated in some families. Drinking, particularly abusive drinking, is often found in families that suffer from a multitude of other social problems. A substantial number of FAS children come from families, either immediate or extended, where alcohol abuse is common, even normative. This frequently results in changes of caregivers or foster placements. New caregivers need to be educated on the special and continuing nutritional needs of the child.

The physical, social, and psychological stresses and the birth of a new baby, particularly one with special needs, places an extra burden upon the recovering woman. This puts the child at risk for poor nutrition and neglect (e.g., the caregiver may forget to prepare food or be unable to adequately provide all the foods necessary for the optimal growth and development of the infant or child.)

WIC can provide supplemental foods, nutrition education and referral to medical and social services which can monitor and provide assistance to the family.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…" Should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Clarren, S.K., and Smith, D.W.: The Fatal Alcohol Syndrome; New England Journal of Medicine; May 11, 1978; 298:1063-1067.
2. Jones, K.L., Smith, D.W., Ulleland, C.N., and Streissguth, A.P.: Pattern of Malformation in Offspring of Chronic Alcoholic Mothers. Lancet; June 9, 1973; 815:1267-1271.
3. Masis B., M.D., May, A: A Comprehensive Local Program for the Prevention of Fetal Alcohol Syndrome, Public Health Reports; September-October 1991; 106: 5; pp. 484-489.
4. Lujan, C.C., BeBruyn, L., May, P.A., and Bird, M.E.: Profile of Abuse and Neglected Indian Children in the Southwest; Child Abuse Negligent; 1989; 34: 449-461.
5. Institute of Medicine: Fetal Alcohol Syndrome, Diagnosis, Epidemiology, Prevention and Treatment; 1996.
6. Weiner, L., Morse, B.A., and Garrido, P.: FAS/FAE Focusing Prevention on Women at Risk; International Journal of the Addictions; 1989; 24:385-395.

Category, Priority and Referral	Category	Priority	Referral
	INFANTS	1	RD, SA

Definition

Neonatal abstinence syndrome (NAS) is a drug withdrawal syndrome that occurs among drug-exposed (primarily opioid-exposed) infants as a result of the mother's use of drugs during pregnancy (1). NAS is a combination of physiologic and neurologic symptoms that can be identified immediately after birth and can last up to 6 months after birth (2, 3).

This condition must be present within the first 6 months of birth and diagnosed, documented, or reported by a physician or someone working under a physician's orders, or as self-reported by the infant's caregiver. See the clarification section for more information about self-reporting a diagnosis.

Required Documentation

Documentation of source of information (i.e. self-report of diagnosis by physician or written documentation by physician).

Justification

Neonatal abstinence syndrome occurs when an infant is born dependent on prescription or illicit drugs the mother was taking during pregnancy. NAS is a combination of withdrawal symptoms that involve multiple bodily systems. It is most commonly associated with chronic opioid exposure during fetal development; however, can also result from chronic intrauterine exposure to other substances including:
benzodiazepines, barbiturates, selective serotonin reuptake inhibitors and ethanol (3). Although these non-opioid substances can lead to NAS, these infants typically respond well to non-pharmacological methods of intervention (4).

Withdrawal in the newborn varies based on the type of substance, does, and timing of exposure (4). Opioid is a general term for a variety of illicit and prescription drugs that decrease pain. Prescription opioid pain relievers include oxycodone, hydrocodone, codeine, morphine, and fentanyl. Opioids are water soluble and are, therefore, able to move easily across the placenta to the infant. This transfer of opioids increases as gestational age increases (3).

Heroin is an illegal opioid that is synthesized from morphine and can be injected, inhaled, or smoked. About 23% of individuals who use heroin become dependent (5). Furthermore, those who take any form of opioid, including prescription opioids as directed for chronic pain, can become addicted. Due to the risk of the transmission of infectious diseases such as Human Immunodeficiency Virus (HIV) and Hepatitis C, women who become pregnant while using illicit opioids, such as heroin, are often put on opioid maintenance therapy. Opioid maintenance therapy involves the prescribed use of either methadone or buprenorphine. These prescribed opioids can still lead to NAS; however, since they are not injected, they decrease the risk of the mother contracting blood borne infectious diseases. Opioid maintenance therapy can also help protect the fetus from repeated opioid withdrawal in utero (6).

The incidence of NAS has increased from 1.2 to 3.39 per 1,000 live births from 2000 to 2009 in the United States. This increased incidence is due to an increase in antepartum opioid use from 1.19 to 5.63 per 1,000 live births in the same period (7). In another study, it was reported that 5.9% of all women who were pregnant in 2012 reported some illicit drug use during pregnancy (4). Infants born with NAS are often premature, have low birth weights, and are growth restricted (see risk #142 *Preterm or Early Term Delivery*, #141 *Low Birth Weight,* and #151 *Small for Gestational Age* for more information about these conditions). (3) In addition to the concerns of exposure to substances in utero, additional factors, including social, nutritional, physical, and mental health problems can also contribute to the health status of the infant. An increased risk of certain birth defects has also been associated with early pregnancy opioid use (8). These birth defects include: spina bifida, hydrocephaly, glaucoma, gastroschisis, and heart defects (9).

Neonatal Abstinence Syndrome Symptoms

Symptoms of NAS generally involve the central nervous system, autonomic nervous system, and the gastrointestinal tract (3). The severity of the infant's symptoms is commonly assessed using the Modified Finnegan Score Sheet. The Modified Finnegan Score Sheet consists of 21 symptoms that are associated with NAS. Following the determination of a baseline score, infants are assessed every 4 hours unless the severity of the symptoms requires more frequent monitoring (10). The following list includes symptoms associated with NAS (1, 6):

- Loud, high-pitched crying
- Sweating
- Yawning
- Sleep disturbances
- Feeding difficulties
- Poor weight gain
- Excessive sucking
- Regurgitation
- Diarrhea

Neonatal Abstinence Syndrome Treatment

Infants with NAS typically have longer hospital stays, can experience serious complications, and have costly treatment (2). The first treatment option for infants with NAS is to manage symptoms without medication by rooming in with the mother, encouraging skin-to-skin contact, swaddling, having a calm environment, avoiding overstimulation, and supporting breastfeeding (11). Infants who are at risk for NAS and who room-in with their mothers are not only at a lower risk of needing pharmacological treatment for NAS, but they also have a shortened hospital stay (12). If withdrawal is severe or if the initial treatment is not successful in managing symptoms of NAS, medications such as morphine, methadone, phenobarbital or clonidine may be used. An infant given these medications may have side effects that could include: slow or shallow breathing, slow heart rate, difficulty waking-up, excessive sleepiness, constipation, and fewer wet diapers (11).

Nutritional Considerations for Neonatal Abstinence Syndrome

The timing and type of feedings play an important role in the management of NAS symptoms. Infants with NAS may have impaired feeding behaviors such as excessive sucking, regurgitation, diarrhea and poor feeding that is characterized by fussiness, crying, and sleepiness (13, 14). Infants with NAS have higher caloric requirements due to their energy expenditure. This combined with the impaired feeding behaviors may result in difficulty with weight gain (14). The American Academy of Pediatrics (AAP) recommends breastfeeding if not contraindicated (15), The AAP also recommends that infants with NAS be fed frequent small volumes of human milk or high calorie formula, as needed, in a quiet and calm environment, to aid the infant in tolerating feedings and improving digestion and to allow for adequate growth (11, 15).

The Academy of Breastfeeding Medicine recommends breastfeeding for women who are on a prescribed stable dose of methadone maintenance because the concentrations of methadone in human milk are low (16). Studies have indicated that, although the amount of methadone in human milk is dependent on the mother's dose, the methadone transferred in human milk averages less than 2.8% of the maternal dose (17). Breastfeeding has been found to provide protection against the development of NAS symptoms and lessen the severity of symptoms, which would decrease the need for pharmacological intervention for the infant (18, 19, 20). The amount of methadone that is in human milk is small and therefore, it is thought that breastfeeding, and not the methadone in human milk, is responsible for its protective impact against NAS (18). Gradual weaning, when mutually desired by the mother and infant, is recommended for breastfeeding women who are being treated for opioid addiction. Gradual weaning (rather than an abrupt stop to breastfeeding) decreases the risk of the infant developing NAS (11, 17).

Implications for WIC Nutrition Services

NAS can be a difficult subject to talk about with WIC participants due to the stigma of addiction. In the WIC clinic, caregivers may not be forthcoming with the infant's diagnosis of NAS and an addiction history of the mother may not be available at the initial assessment. WIC staff can assist caregivers by:

- Educating to recognize infant hunger cues.
- Reviewing feeding frequency and/or formula type and amount to help manage gastrointestinal symptoms of NAS.
- Providing growth monitoring to assess adequate weight gain.
- Encouraging supportive interventions to include:
    - Skin-to-skin contact
    - Swaddling
    - Quiet environment with little stimulation
- Encouraging breastfeeding unless medically contraindicated.
- Providing referrals for support services such as drug and alcohol counseling, parenting support, and medical evaluations.
- Encouraging mothers who are on medication-assisted therapy (e.g., methadone or buprenorphine) and who are breastfeeding, to speak with their health care provider if they have questions about the timing and dose of their medication.
- Educating mothers who are on medication-assisted therapy and who are breastfeeding on the importance of gradual weaning when mutually desired by the mother and infant.

Clarification

Self-reporting of a diagnosis by a medical professional should not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has…") should prompt the CPA to validate the presence of the condition by asking more pointed questions related to that diagnosis.

References

1. Centers for Disease Control and Prevention [Internet]. Georgia: [cited 2017 May 19]. The Problem of Neonatal Abstinence Syndrome; [about 3 screens]. Available from: http://www.ncbi.nlm.nih.gov/pubmedhealth/PMHT0024264/ .
2. Centers for Disease Control and Prevention. Infant and maternal characteristics in neonatal abstinence syndrome-selected hospitals in Florida, 2010-2011. MMWR. 2015 Mar 6.
3. Kocherlakota, P. Neonatal Abstinence Syndrome. Pediatrics. 2014: 134(2):e547-e561.
4. Wiles JR, Isemann B, Ward LP, Vinks AA, Akinbi H. Current management of neonatal abstinence syndrome secondary to intrauterine opioid exposure. J Pediatr. 2014;165(3):440-446.
5. National Institute on Drug Abuse [Internet]. Maryland: [updated 2014 Oct; cited 2016 Feb 3]. DrugFacts: Heroin; [about 4 screens]. Available from: https://www.drugabuse.gov/publications/drugfacts/heroin .
6. Kaltenbach K, Holbrook A, Coyle MG, Heil SH, Salisbury A, Stine S, Martin P, Jones H. Addiction. 2012; 107: 45-52.
7. Bagley SM, Wachman EM, Holland E, Brogly SB. Addiction Science and Clinical Practice. 2014; 9:19.
8. Broussard CS, Rasmussen SA, Reefhuis J, Friedman JM, Jann MW, Riehle-Colarusso T, Honein MA. Maternal treatment with opioid analgesics and risk for birth defects. Am J of Obstetrics and Gynecology. 2011;204(4):314e1-314.e11.
9. Centers for Disease Control and Prevention [Internet]. Georgia: [updated 2014 Oct 22; cited 2016 Feb 3]. Key findings: maternal treatment with opioid analgesics and risk for birth defects. [about 4 screens]. Available from: http://www.cdc.gov/ncbddd/birthdefects/features/birthdefects-opioid-analgesics-keyfindings.html .
10. University of Iowa Children's Hospital [Internet]. Iowa: [updated 2013 Feb 2; cited 2016 Mar 31]. Identifying neonatal abstinence syndrome (NAS) and treatment guidelines. Available from: http://www.uichildrens.org/uploadedFiles/UIChildrens/Health_Professionals/Iowa_Neonatology_Handbook/Pharmacology/Neonatal%20Abstinence%20Syndrome%20Treatment%20Guidelines%20Feb2013%20revision.pdf .
11. Wisconsin Association for Perinatal Care [Internet]. Wisconsin: [updated June 2014; cited 2016 Feb 12]. Assessment and Intervention in the Home: Women and Infants Affected by Opioids. [about 2 screens]. Available from: http://www.perinatalweb.org/assets/cms/uploads/files/Opioid%20Tip%20Sheet_v4.pdf .
12. McKnight S, Coo H, Davies G, Holmes B, Newman A, Newton L, Dow K. Rooming-in for infants at risk of neonatal abstinence syndrome. Am J Perinatol. 2015 Nov 20.

13. Maguire DJ, Rowe MA, Spring H, Elliott AF. Patterns of disruptive feeding behaviors in infants with neonatal abstinence syndrome. Adv Neonatal Care. 2015; 15(6): 429-439.
14. MacMullen NJ, Dulski LA, Blobaum,P. Evidence-based intervention for neonatal abstinence syndrome. J Ped Nurs. 2014; 40(4): 165-203.
15. American Academy of Pediatrics, Committee on Drugs and the Committee on Fetus and Newborn. Clinical report: neonatal drug withdrawal. Pediatrics. 2012; 129: e540-e56.
16. Reece-Stremtan S, Marinelli KA. ABM Clinical Protocol #21: Guidelines for breastfeeding and substance use or substance use disorder. Breastfeeding Medicine. 2015; 10(3): 135-141.
17. Hale TW, Rowe HE. Medications & mother's milk. Hale Publishing. 2014.
18. Liu A, Juarez J, Nair A, Nanan R. Feeding modalities and the onset of the neonatal abstinence syndrome. Frontiers in Pediatrics. 2015; 3(14).
19. Pritham UA. Breastfeeding promotion for management of neonatal abstinence syndrome. J Obstet Gynecol Neonatal Nurs. 2013; 42(5): 517-526.
20. Welle-Strand GK, Skurtveit S, Jansson LM, Bakstad B, Bjarko L, Ravndal E. Breastfeeding reduces the need for withdrawal treatment in opioid-exposed infants. Acta Paediatr. 2013; 102(11): 1060-1066.

Additional Reference:

Food and Nutrition Service, U.S. Department of Agriculture. Substance use prevention: screening, education, and referral resource guide for local WIC agencies. 2013; FNS-276 revised. [cited 2016 Jul 12]. Available from: https://wicworks.fns.usda.gov/wicworks/Topics/ResourceManual.pdf .

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	-
BREASTFEEDING WOMEN	4	-
NON-BREASTFEEDING WOMEN	6	-
CHILDREN ≥ 2 YEARS OF AGE	5	-

Definition

Women and children two years of age and older who meet the eligibility requirements of income, categorical, and residency status may be presumed to be at nutrition risk based on *failure to meet Dietary Guidelines for Americans [Dietary Guidelines]* (1). For this criterion, *failure to meet Dietary Guidelines* risk criterion is defined as consuming fewer than the recommended number of servings from one or more of the basic food groups (grains, fruits, vegetables, milk products, and meat or beans).

The *failure to meet Dietary Guidelines for Americans* risk criterion can only be assigned when a complete nutrition assessment has been completed and no other risk criteria have been identified. This includes assessing for for risk 425 Inappropriate Nutrition Practices for Children or risk 427 Inappropriate Nutrition Practices for Women.

Required Documentation

Complete nutrition assessment

Justification

The 1996 Institute of Medicine (IOM) report, *WIC Nutrition Risk Criteria: A Scientific Assessment* raised questions about the quality of traditional dietary assessment methods (e.g., 24-hour recall and food frequency questionnaires) and recommended further research in the development and validation of diet assessment methodologies (2). In response to the 1996 IOM report, the Food and Nutrition Service (FNS) commissioned the IOM to review the use of various dietary assessment tools and to make recommendations for assessing inadequate diet or inappropriate dietary patterns, especially in the category of *failure to meet Dietary Guidelines* (wee Clarification) (3).

The IOM Committee on Dietary Risk Assessment in the WIC Program approached this task by using the Food Guide Pyramid* recommended number of servings, based on energy needs, as cut-off points for each of the five basic food groups to determine if individuals were meeting the *Dietary Guidelines.* As a result of the review of the cut-off points for food groups and dietary assessment methods, the IOM published the 2002 report, *Dietary Risk Assessment in the WIC Program*. The IOM Committee's findings related to dietary risk, the summary evidence, and the Committee's concluding recommendation are provided below. (4)

IOM Committee Findings Related to Dietary Risk (4) (For more information, refer to the specific pages listed.)

- A dietary risk criterion that uses the WIC applicant's usual intake of the five basic Pyramid* food groups as the indicator and the recommended

number of servings based on energy needs as the cut-off points is consistent with *failure to meet Dietary Guidelines.* (page 130)

- Nearly all U.S. women and children usually consume fewer than the recommended number of servings specified by the Food Guide Pyramid and, therefore, would be at dietary risk based on the criterion *failure to meet Dietary Guidelines*. (page 130)
- Even research-quality dietary assessment methods are not sufficiently accurate or precise to distinguish an individual's eligibility status using criteria based on the Food Guide Pyramid* or on nutrient intake. (page 131)

Summary Evidence Supporting a Presumed Dietary Risk Criterion (4) (For more information, refer to the specific page listed.)

- Less than 1 percent of all women meet recommendations for all five Pyramid* groups. (page 127)
- Less than 1 percent of children ages 2 to 5 years meet recommendations for all five Pyramid* groups. (page 127)
- The percentage of women consuming fruit during 3 days of intake increases with increasing income level. (page 127)
- Members of low-income households are less likely to meet recommendations than are more affluent households. (page 127)
- Food-insecure mothers are less likely to meet recommendations for fruit and vegetable intake than are food-secure mothers. (page 127)
- The percentage of children meeting recommendations for fat and saturated fat as a percentage of food energy increases with increasing income level. (page 127)
- Low-income individuals and African Americans have lower mean Healthy Eating Index scores than do other income and racial/ethnic groups. (page 127)

**The Food Guide Pyramid was the Dietary Guidelines icon at the time the 2002 IOM Committee on Dietary Risk Assessment in the WIC Program conducted the review. The Dietary Guidelines icon has been changed to MyPlate. Although the icon has changed, the Findings and the Supporting Research are still applicable to this criterion. Please see Clarification for more information.*

Summary Evidence Suggesting that Dietary Assessment Methods are Not Sufficient to Determine a WIC Applicant's Dietary Risk (4) (For more information, refer to the specific page listed.)

- 24-hour diet recalls and food records are not good measures of an individual's usual intake unless a number of independent days are observed. (page 61)
- On average, 24-hour diet recalls and food records tend to underestimate usual intake–energy intake in particular. (page 61)
- Food Frequency Questionnaires and diet histories tend to overestimate mean energy intakes. (page 61)

IOM Committee Concluding Recommendation (4) (For more information, refer to the specific page listed.)

"In summary, evidence exists to conclude that nearly all low-income women in the childbearing years and children ages 2 to 5 years are at dietary risk, are vulnerable to nutrition insults, and may benefit from WIC's services. Further, due

to the complex nature of dietary patterns, it is unlikely that a tool will be developed to fulfill its intended purpose with WIC, i.e., to classify individuals accurately with respect to their true dietary risk. Thus, any tools adopted would result in misclassification of the eligibility status of some, potentially many, individuals. By presuming that all who meet the Program's categorical and income eligibility requirements are at dietary risk, WIC retains its potential for preventing and correcting nutrition-related problems while avoiding serious misclassification errors that could lead to denial of services to eligible individuals." (page 135)

Implications for WIC Nutrition Services

As indicated in the 2002 IOM report, most Americans (including most WIC participants) fail to adhere to the *Dietary Guidelines*. Through participant-centered counseling, WIC staff can:

- Guide the participant in choosing healthy foods and age-appropriate physical activities as recommended in the *Dietary Guidelines.*
- Reinforce positive lifestyle behaviors that lead to positive health outcomes.
- Discuss nutrition-related topics of interest to the participant such as food shopping, meal preparation, feeding relationships, and family meals.
- Refer participants, as appropriate, to the Supplemental Nutrition Assistance Program (SNAP), community food banks and other available nutrition assistance programs.

Clarification

The recommendation and findings of the IOM Committee were developed using the *2000 Dietary Guidelines* as the standard for a healthy diet. Subsequent to the 2002 IOM report, the *Dietary Guidelines* have been updated with the release of the *2005* and *2010 Dietary Guidelines*. Although the subsequent editions of the *Dietary Guidelines* are different from the 2000 edition, there is no evidence to suggest that the 2002 IOM recommendation and findings are invalid or inaccurate. The fact remains that diet assessment methodologies may not reflect usual intakes and therefore are insufficient to determine an individual's eligibility status. In addition, future research will be necessary to determine if there is a change in the IOM finding that nearly all Americans fail to consume the number of servings from the basic food groups as recommended in the *Dietary Guidelines*.

References

1. United States Department of Agriculture and the United States Department of Health and Human Services. Dietary Guidelines for Americans, 7th Edition, 2010. Available at: www.usda.gov/cnpp.
2. Institute of Medicine (IOM); Committee on Scientific Evaluation of WIC Nutrition Risk Criteria. WIC nutrition risk criteria: A scientific assessment. Washington, DC: National Academy Press; 1996.
3. United States Department of Agriculture and the United States Department of Health and Human Services. Dietary Guidelines for Americans, 5th Edition, 2000. Available at: www.usda.gov/cnpp
4. Institute of Medicine (IOM); Committee on Dietary Risk Assessment in the WIC Program. Dietary risk assessment in the WIC program. Washington, DC: National Academy Press; 2002. Available at: http://www.iom.edu/Reports/2002/Dietary-Risk-Assessment-in-the-WIC-Program.aspx.

Nutrition Risk Criteria **411 INAPPROPRIATE NUTRITION PRACTICES (INFANTS)**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	INFANTS	4	-

Definition

Routine use of feeding practices that may result in impaired nutrient status, disease, or health problems. These practices, with examples, are outlined below. Refer to "Justification and References" for this criterion.

411.1 Routinely using a substitute(s) for breast milk or for FDA approved iron-fortified formula as the primary nutrient source during the first year of life.

Examples of substitutes:

- Low iron formula without iron supplementation;
- Cow's milk, goat's milk, or sheep's milk (whole, reduced fat, low-fat, skim), canned evaporated or sweetened condensed milk;
- Imitation or substitute milks (such as rice- or soy-based beverages, non-dairy creamer), or other "homemade concoctions."

411.2 Routinely using nursing bottles or cups improperly.

- Using a bottle to feed fruit juice.
- Feeding any sugar-containing fluids, such as soda/soft drinks, gelatin water, corn syrup solutions, sweetened tea.
- Allowing the infant to fall asleep or be put to bed with a bottle at naps or bedtime.
- Allowing the infant to use the bottle without restriction (e.g., walking around with a bottle) or as a pacifier.
- Propping the bottle when feeding.
- Allowing an infant to carry around and drink throughout the day from a covered or training cup.
- Adding any food (cereal or other solid foods) to the infant's bottle.

411.3 Routinely offering complementary foods* or other substances that are inappropriate in type or timing.

- Examples of inappropriate complementary foods:
- Adding sweet agents such as sugar, honey, or syrups to any beverage (including water) or prepared food, or used on a pacifier.
- Introducing any food other than breast milk or iron-fortified infant formula before 6 months of age.

*Complementary foods are any foods or beverages other than breast milk or infant formula.

411.4 Routinely using feeding practices that disregard the developmental needs or stage of the infant.

- Inability to recognize, insensitivity to, or disregarding the infant's cues for hunger and satiety (e.g., forcing an infant to eat a certain type and/or amount of food or beverage or ignoring an infant's hunger cues).
- Feeding foods of inappropriate consistency, size, or shape that put infants at risk of choking.
- Not supporting an infant's need for growing independence with self-feeding (e.g., solely spoon-feeding an infant who is able and ready to finger-feed and/or try self-feeding with appropriate utensils).

- Feeding an infant food with inappropriate textures based on his/her developmental stage (e.g., feeding primarily pureed or liquid foods when the infant is ready and capable of eating mashed, chopped or appropriate finger foods).

411.5 Feeding foods to an infant that could be contaminated with harmful microorganisms or toxins.

Examples of potentially harmful foods

- Unpasteurized fruit or vegetable juice;
- Unpasteurized dairy products or soft cheeses such as feta, Brie, Camembert, blue-veined, and Mexican-style cheese;
- Honey (added to liquids or solid foods, used in cooking, as part of processed foods, on a pacifier, etc.);
- Raw or undercooked meat, fish, poultry, or eggs;
- Raw vegetable sprouts (alfalfa, clover, bean, and radish);
- Deli meats, hot dogs, and processed meats (avoid unless heated until steaming hot).
- Donor breast milk acquired directly from individuals or the Internet.

411.6 Routinely feeding inappropriately diluted formula.

- Failure to follow manufacturer's dilution instructions (to include stretching formula for household economic reasons).
- Failure to follow specific instructions accompanying a prescription.

411.7 Routinely limiting the frequency of nursing of the exclusively breastfed infant when breast milk is the sole source of nutrients.

Examples of inappropriate frequency of nursing:

- Scheduled feedings instead of demand feedings;
- Less than 8 feedings in 24 hours if less than 2 months of age.

411.8 Routinely feeding a diet very low in calories and/or essential nutrients.

Examples:

- Strict vegan diet;
- Macrobiotic diet;
- Other diets very low in calories and/or essential nutrients.

411.9 Routinely using inappropriate sanitation in preparation, handling, and/or storage of expressed breast milk or formula.

Limited or no access to a:

- Safe water supply (documented by appropriate officials e.g., municipal or health department authorities);
- Heat source for sterilization; and/or
- Refrigerator or freezer for storage.

Failure to prepare, handle, and store bottles, storage containers or breast pumps properly; examples include:

Breast Milk

- Thawing/heating in a microwave.
- Refreezing.
- Adding freshly expressed unrefrigerated breast milk to frozen breast milk.
- Adding freshly pumped chilled breast milk to frozen breast milk in an amount that is greater than the amount of frozen breast milk.
- Feeding thawed refrigerated breast milk more than 24 hours after it was thawed.

- Saving breast milk from a used bottled for another feeding.
- Failure to clean breast pump per manufacturer's instruction.
- Feeding donor breast milk acquired directly from individuals or the Internet.

Formula

- Failure to prepare and/or store formula per manufacturer's or physician instructions.
- Storing at room temperature for more than 1 hour.
- Using formula in a bottle one hour after the start of a feeding.
- Saving formula from a used bottle for another feeding.
- Failure to clean baby bottle properly.

411.10 Feeding dietary supplements with potentially harmful consequences.

Examples of dietary supplements, which when fed in excess of recommended dosage, may be toxic or have harmful consequences:

- Single or multi-vitamins;
- Mineral supplements;
- Herbal or botanical supplements/remedies/teas.

411.11 Routinely not providing dietary supplements recognized as essential by national public health policy when an infant's diet alone cannot meet nutrient requirements.

- Infants who are 6 months of age or older who are ingesting less than 0.25 mg of fluoride daily when the water supply contains less than 0.3 ppm fluoride
- Infants who are exclusively breastfed, or who are ingesting less than 1 liter (or 1 quart) per day of vitamin D-fortified formula, and are not taking a supplement of 400 IU of vitamin D

Required Documentation

Document which behaviors are present.

Justification

411.1 Routinely using a substitute(s) for breast milk or for FDA approved iron-fortified formula as the primary nutrient source during the first year of life.

During the first year of life, breastfeeding is the preferred method of infant feeding. The American Academy of Pediatrics (AAP) recommends breast milk for the first 12 months of life because of its acknowledged benefits to infant nutrition, gastrointestinal function, host defense, and psychological well-being (1). In addition, the AAP gas established exclusive breastfeeding as the standard against which all alternative feeding methods must be measured with regard to growth, health, development, and all other short and long-term outcomes for children (2). For infants fed infant formula, iron-fortified formula is generally recommended as a substitute for breastfeeding (1-5). Rapid growth and increased physical activity significantly increase the need for iron and utilizes iron stores (1). Body stores are insufficient to meet the increased iron needs making it necessary for the infant to receive a dependable source of iron to prevent iron deficiency anemia (1). Iron deficiency anemia is associated with cognitive and psychomotor impairments that may be irreversible, and with decreased immune function, apathy, short attention span, and irritability (1, 6). Feeding of low-iron infant formula can compromise an infant's iron stores and lead to iron deficiency anemia. Cow's milk has insufficient and inappropriate

amounts of nutrients and can cause occult blood loss that can lead to iron deficiency, stress on the kidneys from a high renal solute load, and allergic reactions (1, 4, 6-9). Sweetened condensed milk has an abundance of sugar that displaces other nutrients or causes over consumption of calories (10). Homemade formulas prepared with canned evaporated milk do not contain optimal kinds and amounts of nutrients infants need (1, 6, 9, 10). Goat's milk, sheep's milk, imitation milks, and substitute milks do not contain nutrients in amounts appropriate for infants (1, 4, 6, 11, 12).

411.2 Routinely using nursing bottles or cups improperly.

Dental caries is a major health problem in U.S. preschool children, especially in low-income populations (13). Eating and feeding habits that affect tooth decay and are started during infancy may continue into early childhood. Most implicated in this disease process is prolonged use of baby bottles during the day or night, containing fermentable sugars, (e.g., fruit juice, soda, and other sweetened drinks), pacifiers dipped in sweet agents such as sugar, honey or syrups, or other high frequency sugar exposures (14). The AAP and the American Academy of Pedodontics recommended that juice should be offered to infants ($\geq$ 6 months of age) in a cup, not a bottle, and that infants not be put to bed with a bottle in their mouths (15,16). While sleeping with a bottle in his or her mouth, an infant's swallowing and salivary flow decreases, thus creating a pooling of liquid around the teeth (17). The practice of allowing infants to carry or drink from a bottle or training cup of juice for periods throughout the day leads to excessive exposure of the teeth to fermentable carbohydrates, which promotes the development of dental caries (15).

Allowing infants to sleep with a nursing bottle containing fermentable carbohydrates or to use it unsupervised during waking hours provides an almost constant supply of carbohydrates and sugars (1). This leads to rapid demineralization of tooth enamel and an increase in the risk of dental caries due to prolonged contact between cariogenic bacteria on the susceptible tooth surface and the sugars in the consumed liquid (1, 18). The sugars in the liquid pool around the infant's teeth and gums feed the bacteria there, and decay is the result (19). The process may start before the teeth are even fully erupted. Upper incisors (upper front teeth) are particularly vulnerable; the lower incisors are generally protected by the tongue (19). The damage begins as white lesions and progresses to brown or black discoloration typical of caries (19). When early childhood caries is severe, the decayed crowns may break off and the permanent teeth developing below may be damaged (19). Undiagnosed dental caries and other oral pain may contribute to feeding problems and failure to thrive in young children (19).

Unrestricted use of a bottle containing fermentable carbohydrates is a risk because the more times an infant consumes solid or liquid food, the higher the caries risk (1). Feeding behaviors such as unrestricted use of the bottle and frequent snacking can be habit forming in later infancy and may carry over into toddler-hood. Frequent cariogenic snacks eaten between meals place the toddler at high risk for caries development; this includes the habit of continually sipping from cups (or bottles) containing cariogenic liquids (juice, milk, soda, or sweetened liquid) (19). If inappropriate use of the bottle persists, the child is at risk of toothaches, costly dental treatment, loss of primary teeth, and developmental lags on eating and chewing. If this continues beyond the usual weaning period, there is a risk of decay to permanent teeth.

Propping the bottle deprives infants of vital human contact and nurturing which makes them feel secure. It can also cause ear infections because of fluid entering the middle ear and not draining properly; choking from liquid flowing into the lungs; and tooth decay from prolonged exposure to carbohydrate-containing liquids (20).

Adding solid food to a nursing bottle results in force-feeding, inappropriately increases the energy and nutrient composition of the formula, deprives the infant of experiences important in the development of feeding behavior, and could cause an infant to choke (1, 11, 21, 22).

411.3 Routinely offering complementary foods or other substances that are inappropriate in type of timing.

Infants, especially those living in poverty, are at high risk for developing early childhood caries (13). Most implicated in this disease process are: prolonged use of baby bottles containing fermentable sugars, (e.g., fruit juice, soda, and other sweetened drinks) during the day or night; pacifiers dipped in sweet agents such as sugar, honey or syrups, or other high frequency sugar exposures (14).

The AAP recommends exclusive breastfeeding through 6 months of age (1). Feeding solid foods too early (i.e., before 6 months of age) by, for example, adding dilute cereal or other solid foods to bottles deprives infants of the opportunity to learn to feed themselves (4, 11, 23). The major objection to the introduction of solids before age 6 months of age is based on the possibility that it may interfere with establishing sound eating habits and may contribute to overfeeding (6, 24). In early infancy, the infant possesses an extrusion reflex that enables him/her to swallow only liquid foods (1, 13, 25). The extrusion reflex is normally diminished by 6 months of age (1). Breast milk or iron-fortified infant formula is all the infant needs. Gastric secretions, digestive capacity, renal capacity and enzymatic secretions are low, which makes digestion of solids inefficient and potentially harmful (6, 24, 25). Furthermore, there is the potential for antigens to be developed against solid foods, due to the undigested proteins that may permeate the gut; however, the potential for developing allergic reactions may primarily be in infants with a strong family history of atopy (6, 24). If solid foods are introduced before the infant is developmentally ready, breast milk or iron-fortified formula necessary for optimum growth is displaced (1, 25). Around 6 months of age, the infant is developmentally ready for solid foods when: the infant is better able to express certain feeding cues such as turning head to indicate satiation; oral and gross motor skills begin to develop that help the infant to take solid foods; the extrusion reflex disappears; and the infant begins to sit upright and maintain balance with little or no support (1, 6, 24, 25).

The AAP advises against giving fruit juice to infants younger than 6 months since it offers no nutritional benefit at this age (1). Offering juice before solid foods are introduced into the diet could risk having juice replace breast milk or infant formula in the diet (15). This can result in reduced intake of protein, fat, vitamins, and minerals such as iron, calcium, and zinc (26). It is prudent to give juice only to infants who can drink from a cup (15).

411.4 Routinely using feeding practices that disregard the developmental needs or stage of the infant.

Infants held to rigid feeding schedules are often underfed or overfed. Caregivers insensitive to signs of hunger and satiety, or who over-manage feeding may inappropriately restrict or encourage excessive intake. Findings show that these

practices may promote negative or unpleasant associations with eating that may continue into later life, and may also contribute to obesity. Infrequent breastfeeding can result in lactation insufficiency and infant failure-to-thrive. Infants should be fed foods with a texture appropriate to their developmental level. (4, 6, 11, 13, 23)

411.5 Feeding foods to an infant that could be contaminated with harmful microorganisms or toxins.

Only pasteurized juice is safe for infants, children, and adolescents (15). Pasteurized fruit juices are free of microorganisms (15). Unpasteurized juice may contain pathogens, such as Escherichia coli, Salmonella, and Cryptosporidium organisms (15, 27). These organisms can cause serious disease, such as hemolytic-uremic syndrome, and should never be fed to infants and children (15). Unpasteurized juice must contain a warning on the label that the product may contain harmful bacteria (15, 28). Infants or young children should not eat raw or unpasteurized milk or cheeses (1)—unpasteurized dairy products could contain harmful bacteria, such as Brucella species, that could cause young children to contract a dangerous food borne illness. The AAP also recommends that young children should not eat soft cheeses such as feta, Brie, Camembert, blue-veined, and Mexican-style cheese—these foods could contain Listeria bacteria (1). Hard cheeses, processed cheeses, cream cheese, cottage cheese, and yogurt need not be avoided (1).

Honey has been implicated as the primary food source of *Clostridium botulinum* during infancy. These spores are extremely resistant to heat, including pasteurization, and are not destroyed by present methods of processing honey. Botulism in infancy is caused by ingestion of the spores, which germinate into the toxin in the lumen of the bowel (10, 11, 29, 30).

Infants or young children should not eat raw or undercooked meat or poultry, raw fish or shellfish, including oysters, clams, mussels, and scallops—these foods may contain harmful bacteria or parasites that could cause children to contract a dangerous food borne illness (1).

According to the AAP, to prevent food borne illness, the foods listed below should not be fed to infants or young children (1). All of the foods have been implicated in selected outbreaks of food-borne illness, including in children. Background information regarding foods that could be contaminated with harmful microorganisms is also included below (1):

- Raw vegetable sprouts (alfalfa, clover, bean, and radish)--can cause potentially dangerous Salmonella and E. coli O157 infection. Sprouts grown under clean conditions in the home also present a risk because bacteria may be present in seeds. Cook sprouts to significantly reduce the risk of illness (31).
- Deli meats, hot dogs, and processed meats (avoid unless heated until steaming hot) --These foods have been found to be contaminated with *Listeria monocytogenes*; if adequately cooked, this bacteria is destroyed.

Please see section 411.9 below under “Breast Milk”, for information related to the use of donor human milk acquired via the internet or directly from an individual.

411.6 Routinely feeding inappropriately diluted formula.

Over-dilution can result in water intoxication resulting in hyponatremia; irritability; coma; inadequate nutrient intake; failure to thrive; and/or poor growth (1, 4, 6, 11, 32). Under-dilution of formula increases calories, protein, and solutes presented to the kidney for excretion, and can result in hypernatremia, tetany, and obesity (4, 6, 11, 32).

Dehydration and metabolic acidosis can occur with under-dilution of formula (4, 6, 11, 32). Powdered formulas vary in density so manufacturers' scoops are formula-specific to assure correct dilution (6). One clue for staff to identify incorrect formula preparation is to determine if the parent/caregiver is using the correct manufacturer's scoop to prepare the formula.

411.7 Routinely limiting the frequency of nursing of the exclusively breastfed infant when breast milk is the sole source of nutrients.

Exclusive breastfeeding provides ideal nutrition to an infant and is sufficient to support optimal growth and development in the first 6 months of life (5). Human breast colostrum and milk have been studied extensively. They are composed of a mixture of nutritive components and other bioactive factors that are easy to digest and absorb and have strong physiologic effects upon the infant, and their composition changes over time to meet the infant's changing nutritional needs (1).

Frequent breastfeeding is critical to the establishment and maintenance of an adequate milk supply for the infant (5, 33-37). Inadequate frequency of breastfeeding may lead to lactation failure in the mother and dehydration, poor weight gain, diarrhea, and vomiting, illness, and malnourishment in the infant (5, 35, 38-43). Exclusive breastfeeding protects infants from early exposure to contaminated foods and liquids (41). Infants who receive breast milk more than infant formulas, have a lower risk of being overweight in childhood and adolescence (44, 45). In addition, a summary report of several primary studies and meta-analyses has reported that a history of breastfeeding is associated with a reduction in the risk of otitis media, gastroenteritis, hospitalization for lower respiratory tract infections, atopic dermatitis, sudden infant death syndrome, childhood asthma, childhood leukemia, and type 1 and 2 diabetes (46).

411.8 Routinely feeding a diet very low in calories and/or essential nutrients.

Highly restrictive diets prevent adequate intake of nutrients, interfere with growth and development, and may lead to other adverse physiological effects (4). Infants older than 6 months are potentially at the greatest risk for overt deficiency states related to inappropriate restrictions of the diet, although deficiencies of vitamins B12 and essential fatty acids may appear earlier (1, 47, 48). Infants are particularly vulnerable during the weaning period if fed a macrobiotic diet and may experience psychomotor delay in some instances (1, 49, 50). Well-balanced vegetarian diets with dairy products and eggs are generally associated with good health. However, strict vegan diets may be inadequate in calories, vitamin B12, vitamin D, calcium, iron, protein and essential amino acids needed for growth and development (51). The more limited the diet, the greater the health risk. Given the health and nutrition risks associated with highly restrictive diets, WIC can help the parent to assure that the infant consumes an adequate diet to optimize health during critical periods of growth as well as for the long term.

411.9 Routinely using inappropriate sanitation in preparation, handling,

and storage of expressed breast milk or formula.

Lack of sanitation in the preparation, handling and storage of expressed breast milk or formula may cause gastrointestinal infection. The water used to prepare concentrated or powdered infant formula and prepare bottles and nipples (for formula and breast milk) must be safe for consumption. Water contaminated with toxic substances (such as nitrates, lead, or pesticides) poses a hazard to an infant's health and should NOT be used (10). In addition, a heat source is necessary to sterilize bottles and other items used in the storage of both breast milk and formula. Adequate refrigeration (40 Degrees Fahrenheit or below) is necessary to safely store breast milk and prepared formula (10).

Breast Milk

Published guidelines on the handling and storage of breast milk may differ among pediatric nutrition authorities (1, 10, 52-55). However, the following breast milk feeding, handling, and storage practices that are considered inappropriate and unsafe (10, 52, 56-59):

- Thawing frozen breast milk in the microwave oven;
- Refreezing breast milk;
- Adding freshly expressed unrefrigerated breast milk to already frozen milk in a storage container*****
- Feeding previously frozen breast milk thawed in the refrigerator that has been refrigerated for more than 24 hours
- Saving breast milk from a used bottle for use at a subsequent feeding
- Failure to clean a breast pump per manufacturer's instruction
- Feeding donor breast milk acquired directly from individuals or the internet.

Another consideration when recommending length of storage time is its effect on protective properties in breast milk. There is evidence that after 48 hours of refrigeration, breast milk significantly loses important antibacterial and antioxidant properties (60). These properties of breast milk are specifically important for the prevention of necrotizing enterocolitis, retinopathy, and bronchopulmonary dysplasia of premature infants (5). Although some properties may be reduced with longer refrigerated storage, this does not diminish the overall superiority of breast milk over formula, as formula does not contain these protective properties or many of the other benefits of breast milk.

Participant circumstances (e.g., adequate refrigeration, safe water, heat source), the health of the infant and health care provider directions need to be considered when recommending the length of time breast milk should be stored.

If the breastfeeding mother uses a breast pump, it is essential for her to fully understand the importance of and the specific manufacturer's instructions for cleaning the breast pump. Improper cleaning of breast pumps and pump parts can increase the risk of expressed breast milk contamination (58).

With increased awareness of the benefits and efforts to promote breastfeeding, more mothers are choosing to breastfeed, as evidenced by data from CDC in the Breastfeeding Report Card (61). But in situations such as illness, physical inability to produce breast milk, decisions to not breastfeed, or adoptive parents seeking breast milk, the desire to provide breast milk may prompt parents/caregivers to turn to alternate methods of obtaining breast milk to feed

their infant. Since the cost of banked breast milk can be prohibitive for WIC clients, these mothers may turn to informal milk sharing from known sources such as friends or relatives, or from unknown sources such as internet sites or other advertisements.

A study that evaluated breast milk shared via the internet concluded that there was a high overall rate of bacterial growth and contamination, which suggests poor collection, storage, and shipping practices (62). In another study, researchers looked at current and past infection among potential donors to a breast milk bank. It was revealed that a minimum of 3% of potential donors had positive serology for disease conditions such as syphilis, HIV, hepatitis B, hepatitis C, HTLV-1 or HTLV-2 (63). It was concluded that if these relatively low risk potential donors tested positive then the untested or unscreened women of donor human milk may present a significant health risk (63).

Although sharing breast milk between those with an excess milk supply and those seeking milk for their infant may be growing in popularity (often facilitated by web sites established to link providers and recipients), both the AAP and the Food and Drug Administration (FDA) recommend against feeding infants breast milk obtained directly from individuals or through the internet (59, 64). Obtaining donor breast milk via these means is discouraged due to the lack of adequate screening for infectious diseases and the risk of contamination (59).

The FDA suggests that a decision to give donor breast milk should be made in consultation with the infant's health care provider and only screened donor breast milk should be used. Also, caregivers should consult with the infant's health care provider on where to obtain screened donor breast milk (59). Due to the lack of Federal guidelines and standards pertaining to the operation, quality, and safety of breast milk banks and potential liability concerns, the U.S. Department of Agriculture, Food and Nutrition Service does not authorize banked breast milk as an allowable substitute for WIC-eligible formulas (see WIC Policy Memorandum 2000-2: *Use of Banked Human Breast Milk in the WIC Program*).

Formula

Formula must be properly prepared in a sanitary manner to be safe for consumption. Furthermore, prepared infant formula is a perishable food, and must be handled and stored properly to be safe for consumption (4, 10).

Most babies who are hospitalized for vomiting and diarrhea are bottle fed. This has often been attributed to the improper handling of formula rather than sensitivities to the formula. In rare cases, the contaminated powdered formulas may cause infections in preterm or immune compromised infants. To reduce the risk of infection in infants it is important that formulas be carefully prepared and handled. All formula should be prepared according to the manufacturer's instruction on the label, or those given by the health care provider.

Manufacturers' instructions vary, depending on the product, in the length of time it is considered safe to store prepared infant formula without refrigeration before bacterial growth accelerates to an extent that the infant is placed at risk (1). Published guidelines on the handling and storage of infant formula indicate that it is unsafe to use prepared formula which (1):

- Has been held at room temperature longer than 1 hour or longer than recommended by the manufacturer

- Has been held in the refrigerator longer than the safe storage time indicated by the manufacturer
- Remains in a bottle one hour after the start of feeding
- Remains in a bottle from an earlier feeding
- Is fed using improperly cleaned baby bottles

411.10 Feeding dietary supplements with potentially harmful consequences.

An infant consuming inappropriate or excessive amounts of single or multivitamin or mineral or herbal remedy not prescribed by a physician is at risk for a variety of adverse effects including harmful nutrient interactions, toxicity, and teratogenicity (1, 65). While some herbal teas may be safe, some have undesirable effects, particularly on infants who are fed herbal teas or who receive breast milk from mothers who have ingested herbal teas (66). Examples of teas with potentially harmful effects to infants and children include: licorice, comfrey leaves, sassafras, senna, buckhorn bark, cinnamon, wormwood, woodruff, valerian, foxglove, pokeroot or pokeweed, periwinkle, nutmeg, catnip, hydrangea, juniper, Mormon tea, thorn apple, yohimbe bark, lobelia, oleander, Maté, kola nut or gotu cola, and chamomile (66-68). Like drugs, herbal or botanical preparations have chemical and biological activity, may have side effects, and may interact with certain medications--these interactions can cause problems and can even be dangerous (69). Botanical supplements are not necessarily safe because the safety of a botanical depends on many things, such as its chemical makeup, how it works in the body, how it is prepared, and the dose used (69).

411.11 Routinely not providing dietary supplements recognized as essential by national public health policy when an infant's diet alone cannot meet nutrient requirements.

Depending on an infant's specific needs and environmental circumstances, certain dietary supplements may be recommended by the infant's health care provider to ensure health. For example, fluoride supplements may be of benefit in reducing dental decay for children living in fluoride-deficient areas (1, 70).

To prevent rickets and vitamin D deficiency in healthy infants and children, the AAP recommends a supplement of 400 IU per day for the following (5, 71).

- All breastfed and partially breastfed infants unless they are weaned to at least 1 liter (or 1 quart) per day of vitamin D-fortified formula.
- All non-breastfed infants who are ingesting less than 1 liter (or 1 quart) per day of vitamin D-fortified formula.

References

1. Committee on Nutrition, American Academy of Pediatrics. Pediatrics nutrition handbook. 7th Ed. Elk Grove Village, III: American Academy of Pediatrics, 2009.
2. American Academy of Pediatrics. Policy Statement Breastfeeding and the Use of Human Milk. Organizational Principles to Guide and Define the Child Health Care System and/or Improve the Health of all Children. Section on Breastfeeding. (Downloaded from pediatrics.aappublications.org accessed on January 6, 2015.)

3. American Academy of Pediatrics, Committee on Nutrition. Iron fortification of infant formula. Pediatrics 1999; 104:119-123.
4. Institute of Medicine. WIC nutrition risk criteria: a scientific assessment. National Academy Press, Washington D.C.; 1996.
5. American Academy of Pediatrics, Section on Breastfeeding: Breastfeeding and the use of human milk. Pediatrics 2005 Feb; 115(2):496-506.
6. Fomon SJ. Nutrition of normal infants. St. Louis: Mosby, 1993.
7. Whitney EN, Rolfes SR. Understanding nutrition. 9th Ed. Wadsworth: Thomson Learning, 2002: p. 241.
8. American Academy of Pediatrics, Committee on Nutrition. The use of whole cow's milk in infancy. Pediatrics 1992; 89(6):1105-1109.
9. Friel JK, et al. Eighteen-month follow-up of infants fed evaporated milk formula. Canadian Journal of Public Health. Revue Canadienne de Sante Publique, 90.4 (Jul-Aug 1999): 240-3. Abstract.
10. United States Department of Agriculture, Food and Nutrition Service. Infant nutrition and feeding, a guide for use in the WIC and CSF programs. Alexandria, VA: Special Supplemental Nutrition Programs, revised 2008. [FNS-288].
11. Trahms CM, Pipes PL, editors. Nutrition in Infancy and Childhood. WCB/McGraw-Hill; 1997.
12. Bellioni-Businco B, Paganelli R, Lucenti P, Giampietro PG, Perborn H, Businco L. Allergenicity of goat's milk in children with cow's milk allergy. J. Allergy Clin. Immunol. 1999; 103:1191-1194.
13. Tang J, Altman DS, Robertson D, O"Sullivan DM, Douglas JM, Tinanoff N. Dental caries prevalence and treatment levels in Arizona preschool children. Public Health Rep. 1997; 112:319-29.
14. Tinaoff N and Palmer CA. Dietary determinants of dental carries and dietary recommendations for preschool children. J. Public Health Dent. 2000; 60(3):197-206.
15. American Academy of Pediatrics, Committee on Nutrition. The use and misuse of fruit juice in pediatrics. Pediatrics 2001; 107:1210-1213.
16. American Academy of Pediatrics and American Academy of Pedodontics. Juice in ready-to-use bottles and nursing bottle carries. AAP News. 1978; 29(1):11.
17. Samour PQ, Helm KK, Lang CE. Handbook of pediatric nutrition. 2nd Ed. Gaithersburg, MD: Aspen Publishers, Inc.; 1999.
18. American Academy of Pediatric Dentistry. Baby Bottle Tooth Decay/Early Childhood Caries. Pediatr Dent 2000-2001 (revised May 1996); 2001 Mar-Apr; 23(2):18.
19. Fitzsimons D, Dwyer JT, Palmer C, Boyd LD. Nutrition and oral health guidelines for pregnant women, infants, and children. J. Am. Diet. Assoc. Feb 1998; 98(2):182-6.
20. Shelov SD. Caring for your baby and young child: birth to age 5. Elk Grove Village, IL: American Academy of Pediatrics; 1998.
21. Satter E. Child of mine: Feeding with love and good sense. Palo Alto (CA): Bull Publishing Company; 2000.
22. Tamborlane WV, editor. The Yale guide to children's nutrition. Connecticut: Yale University; 1997.
23. Williams CP, editor. Peidatric manual of clinical dietetics. Chicago: American Dietetic Association; 1998.
24. Fomon SJ. Feeding normal infants: rationale for recommendations. J. Am. Diet. Assoc. 2001; 101:1002-1005.
25. Rolfes, Debruyne, Whitney. Life span nutrition: conception through life; 1990; pp. 231-237.

26. Gibson SA. Non-milk extrinsic sugars in the diets of pre-school children: association with intakes of micronutrients, energy, fat and NSP. Br. J. Nutr. 1997; 78:367-378.
27. Parish ME. Public health and non-pasteurized fruit juices. Crit. Rev. Microbiol. 1997; 23:109-119.
28. Food Labeling. Warning and Notice Statement: Labeling of juice products; Final Rule. 63 Federal Register 37029-37056 (1998) (codified at 21 CFR §101, 120).
29. Botulism Fact Sheet [electronic file]. Atlanta (GA): Centers for Disease Control and Prevention; 1995.
30. Centers for Disease Control and Prevention (US). Botulism in the United States, 1899-1996. Atlanta (GA): Centers for Disease Control and Prevention; 1998.
31. Food and Drug Administration. Updates: Avoid raw sprouts to reduce food poisoning risk, agency advises. FDA Consumer magazine, September-October 1999. Available from: http://www.fda.gov/fdac/departs/1999/599_upd.html.
32. Fein SB, Falci CD. Infant formula preparation, handling, and related practices in the United States. J. Am. Diet. Assoc. 1999. 99:1234-1240.
33. Biancuzzo M. Breastfeeding the newborn: clinical strategies for nurses. St. Louis, MO; Morby, 1999, Pages 103-104.
34. Mochbracher N, Stock J. The Breastfeeding answer book (Revised edition). La Leche League International, 1997, Pages 20-23.
35. Eiger MS, Olds SW. The complete book of breastfeed. New York: Workman Publishing: 1999, p.88, 112-114.
36. Rosenthal MS. The breastfeeding sourcebook. Los Angeles: Lowell House; 1996, p. 157.
37. Sears M, Sears W. The breastfeeding book. Boston: Little, Brown and Company; 2000, p. 108-110.
38. Johnson DB. Nutrition in infancy:-evolving views on recommendations. Nutrition Today 1998; 32: 63-68.
39. Mark DH. Breastfeeding and infant illness: a dose-response relationship. J Amer Med Assoc 1990; 281: 1154.
40. Muztagh M. Optimal breastfeeding duration. J. Am. Diet. Assoc. 1997;97: 1252-1255.
41. Raiser J, Alexander C, O'Campo P. Breastfeeding and Infant Illness; a dose-response relationship? Am. J. Pub. Health 1999; 89: 25-30.
42. Scariest PD, Grummer-Strawn LM, Fein SB. A longitudinal analysis of infant mortality and the extent of breastfeeding in the United States. Pediatrics 1997; 99-6.
43. Story M, Hoyt K, Sofka D. Bright futures in practice. National Center for Education in Maternal and Child Health. Arlington: Georgetown University; 2000, p. 25.
44. Gilman MW, Rifas-Shiman SL, Camrgo CA Jr, Berkey CS, Frazier AL, Rockett HR, Field AE, Colditz GA. Risk of overweight among adolescents who were breastfed infants. J. Amer. Med. Assoc. 2001; 285(19): 2461-7.
45. Von Kries R, Koletzko B, Sauerwald T, von Mutius E, Barnert D, Grunert V, Von Voss H. Breastfeeding and obesity: cross-sectional study. Br. Med. J. 1999: 319(7203):147-50.
46. Ip S, Chung M, Ramen G, Trikalinos TA, Lau J. A Summary of the Agency for Healthcare Research and Quality's Evidence Report on Breastfeeding in Developed Countries. Breastfeed Med. 2009;4(suppl 1):S17-S30.
47. Sanders TA, Reddy S. Vegetarian diets and children. Am. J. Clin. Nutr. 1994; 59(suppl):1176S-1181S.

48. Sanders TA. Essential fatty acid requirements of vegetarians in pregnancy, lactation and infancy. Am. J. Clin. Nutr. 1999; 70:5555-5595.
49. Sanders TA. Vegetarian diets and children. Pediatric Clin. North Am. 1995; 42-955-965.
50. Dagnelie PC, Vergote FJ, van Staveren, WA, et al. High prevalence of rickets in infants on macrobiotic diets. Am. J. Clin. Nutr. 1990;51:202-208.
51. Duyff RL. American Dietetic Association. The American Dietetic Association's complete food and nutrition guide. Minneapolis, MN: Chronimed Pub; 1996.
52. American Academy of Pediatrics: A woman's guide to breastfeeding. 1999, pp. 13-14.
53. United States Department of Agriculture (USDA), Food and Nutrition Service. Breastfed Babies Welcome Here [Program Aid 1516]. Alexandria, VA: USDA, 1995, pp. 12-15.
54. Lawrence RA. Breastfeeding: a guide for the medical profession. 5th edition. St. Louis, MO: Mosby, 1999, pp. 677-710.
55. Academy of Breastfeeding Medicine Protocol Committee. ABM clinical protocol #8: human milk storage information for home use for full-term infants; Revision #1. Academy of Breastfeeding; March 2010. Available from: http://www.bfmed.org/Media/Files/Protocols/Protocol%208%20-%20English%20revised%202010.pdf.
56. Duke, CS. Common concerns when storing human milk. New Beginnings; July-August 1998; 15(4), p. 109.
57. Neifert, M. Dr. Mom's guide to breastfeeding. 1998; New York, NY: Plume, pp. 305-306.
58. Jones F, Tully, MR. Best practice for expressing, storing, and handling human milk in hospitals, homes and child care settings. Raleigh, NC: Human Milk Banking Association of North America Inc.; 2006.
59. US Food and Drug Administration. Use of donor human milk [Internet]. Washington DC: [updated 2015 Aug 7; cited 2016 Aug 17]. Available from: http://www.fda.gov/ScienceResearch/SpecialTopics/PediatricTherapeuticsResearch/ucm235203.htm .
60. Lee JW, Davis JM. Future applications of antioxidants in premature infants. Vurr Opin Pediatr. 2011 Apr; 23(2):161-6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21150443.
61. Centers for Disease Control and Prevention. Breastfeeding Report Card 2014 [Internet]. Georgia: 2014 [cited 2016 Aug 17]. Available from: http://www.cdc.gov/breastfeeding/pdf/2014breastfeedingreportcard.pdf .
62. American Academy of Pediatrics. Microbial contamination of human milk purchased via the internet. Pediatrics. Originally published online October 21, 2013; DOI: 10.1542/peds. 2013-1687.
63. Cohen RS, Xiong SC, Sakamoto P. Retrospective review of serological testing of potential human donors. Arch Dis Child Fetal Neonatal Ed 2010 95: F118-F120. doi: 10.1136/adc.2008.1564711997;100:1035-1038.
64. American Academy of Pediatrics. Breastfeeding and the use of human milk. Pediatrics. Feb 2005; American Academy of Pediatrics. Volume 115/Issue2, Revised 100(6):1035. Available from: http://pediatrics.aappublications.org/content/115/2/496 .
65. Anderson JV, Van Nierop MR. Basic nutrition facts a nutrition reference. Lansing, MI: Michigan Department of Public Health; 1989.
66. Lawrence RA. Breastfeeding: a guide for the medical profession. 5th edition. St. Louis, MO: Mosby, 1999, pp. 371-377.

67. Siegel RK. Herbal intoxication: psychoactive effects from herbal cigarettes, tea and capsules. JAMA 236:473, 1976.
68. Ridker PM. Toxic effects of herbal teas. Arch Environ Health 42(3):133-6, 1987.
69. Office of Dietary Supplements, National Institutes of Health (NIH). Botanical dietary supplements: background information. NIH web page, last updated 7/7/2004. Available from: http://ods.od.nih.gov/factsheets/BotanicalBackground.asp.
70. American Academy of Pediatric Dentistry. Fluoride. Pediatric Dent. 1999; 21:40.
71. American Academy of Pediatrics, Section on Breastfeeding and Committee on Nutrition. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. Pediatrics 2008; http://pediatrics.aappublications.org/content/122/5/1142.full.

Nutrition Risk Criteria **425 INAPPROPRIATE NUTRITION PRACTICES (CHILD)**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	CHILDREN	5	-

Definition

FORMERLY QUESTIONABLE CHILD FEEDING

Routine use of feeding practices that may result in impaired nutrient status, disease, or health problems. These practices, with examples, are outlined below. Refer to "Justification and References" for this criterion.

425.1 Routinely feeding inappropriate beverages as the primary milk source.
Examples of inappropriate beverages as primary milk source:

- Non-fat or reduced-fat milks (between 12 and 24 months of age, unless allowed by State agency policy for a child for whom overweight or obesity is a concern) or sweetened condensed milk
- Goat's milk, sheep's milk, imitation or substitute milks (that are unfortified or inadequately fortified), or other "homemade concoctions."

425.2 Routinely feeding a child any sugar-containing fluids.
Examples of sugar-containing fluids:

- Soda/soft drinks
- Gelatin water
- Corn syrup solutions
- Sweetened tea

425.3 Routinely using nursing bottles, cups, or pacifiers improperly.

- Using a bottle to feed
  - fruit juice
  - diluted cereal or other solid foods
- Allowing the child to fall asleep or be put to bed with a bottle at naps or bedtime
- Allowing the child to use the bottle without restriction (e.g., walking around with a bottle) or as a pacifier
- Using a bottle for feeding or drinking beyond 14 months of age
- Using a pacifier dipped in sweet agents such as sugar, honey, or syrups
- Allowing a child to carry around and drink throughout the day from a covered or training cup

425.4 Routinely using feeding practices that disregard the developmental needs or stages of the child.

- Inability to recognize, insensitivity to, or disregarding the child's cues for hunger and satiety (e.g., forcing a child to eat a certain type and/or amount of food or beverage or ignoring a hungry child's requests for appropriate foods)
- Feeding foods of inappropriate consistency, size, or shape that put children at risk of choking
- Not supporting a child's need for growing independence with self-feeding (e.g., solely spoon-feeding a child who is able and ready to finger-feed and/or try self-feeding with appropriate utensils)

- Feeding a child food with an inappropriate texture based on his/her developmental stage (e.g., feeding primarily pureed or liquid food when the child is ready and capable of eating mashed, chopped or appropriate finger foods)

425. 5 Feeding foods to a child that could be contaminated with harmful microorganisms.
Examples of potentially harmful foods for a child:

- Unpasteurized fruit or vegetable juice
- Unpasteurized dairy products or soft cheeses such as feta, Brie, Camembert, blue-veined, and Mexican-style cheese
- Raw or undercooked meat, fish, poultry, or eggs
- Raw vegetable sprouts (alfalfa, clover, bean, and radish)
- Deli meats, hot dogs, and processed meats (avoid unless heated until steaming hot)

425.6 Routinely feeding a diet very low in calories and/or essential nutrients.
Examples:

- Vegan diet
- Macrobiotic diet
- Other diets very low in calories and/or essential nutrients

425.7 Feeding dietary supplements with potentially harmful consequences.
Examples of dietary supplements which when fed in excess of recommended dosage may be toxic or have harmful consequences:

- Single or multi-vitamins
- Mineral supplements
- Herbal or botanical supplements/remedies/teas

425.8 Routinely not providing dietary supplements recognized as essential by national public health policy when a child's diet alone cannot meet nutrient requirements.

- Providing children under 36 months of age less than 0.25 mg of fluoride daily when the water supply contains less than 0.3 ppm fluoride
- Providing children 36-60 months of age less than 0.50 mg of fluoride daily when the water supply contains less than 0.3 ppm fluoride
- Not providing 400 IU of vitamin D if a child consumes less than 1 liter (or 1 quart, approximately 4 cups) of vitamin D-fortified milk or formula

425.9 Routine ingestion of nonfood items (pica).
Examples of inappropriate nonfood items:

- Ashes
- Carpet fibers
- Cigarettes or cigarette butts
- Clay
- Dust
- Foam rubber
- Paint chips
- Soil
- Starch (laundry and cornstarch)

Required Documentation	Document which behaviors are present.
Justification	**425.1 Routinely feeding inappropriate beverages as the primary milk source.** Goat's milk, sheep's milk, imitation milks and substitute milks (that are unfortified or inadequately fortified) do not contain nutrients in amounts appropriate as a primary milk source for children (1-4). Non-fat and reduced-fat milks are not recommended for use with children from 1 to 2 years of age because of the lower calorie density compared with whole-fat products (1, 5). The low-calorie, low-fat content of these milks requires an increase in caloric intake to meet energy needs. Infants and children under two using reduced fat milks gain at a slower growth rate, lose body fat as evidenced by skin fold thickness, lose energy reserves, and are at risk of inadequate intake of essential fatty acids. Additionally, essential fatty acids are a critical component of infant and child brain development with deficits early in life leading to significantly altered brain structure and function (6-8). Similar malnourishment has been associated with negative health outcomes including, but not limited to, slower language development, poorer motor function, lower IQ, poorer school performance, and eyesight problems (9). WIC Regulations [7 CFR 246.10(e)], however, include the option for WIC State agencies to issue reduced-fat milk and/or reduced-fat yogurt to children (1 to 2 years of age) for whom overweight or obesity is a concern, as determined by the Competent Professional Authority (CPA) (Food Package Guidance, May 2014). This option is consistent with the American Academy of Pediatrics (AAP) recommendation in the clinical report: *Lipid Screening and Cardiovascular Health in Childhood* (10). The AAP identifies parental history of obesity, lipidemia, and cardiovascular disease as determinants for a child for whom overweight or obesity is a concern. WIC State agencies that choose to authorize reduced-fat milk and/or reduced fat yogurt for the 1 year old child must develop policy that defines the assessment criteria the CPA will use to determine if the child should be given reduced-fat dairy products. For example, a State agency may choose to use existing nutrition risk criteria: #114 *Overweight or At Risk of Overweight (Infants and Children)* and/or #115 *High Weight-for-Length (Infants and Children <24 Months of Age)* to identify children to receive reduced-fat milk. For more information about the required State agency policy for issuing reduced-fat milk to children 12 months to 2 years of age, please see Food and Nutrition Service, Food Package Guidance issued May 2014. **425.2 Routinely feeding a child any sugar-containing fluids.** Abundant epidemiologic evidence from groups who have consumed low quantities of sugar as well as from those who have consumed high quantities shows that sugar—especially sucrose—is the major dietary factor affecting the prevalence and progression of dental carries (11). Consumption of foods and beverages high in fermentable carbohydrates, such as sucrose, increases the risk of early childhood caries and tooth decay (11, 12). **425.3 Routinely using nursing bottles, cups, or pacifiers improperly.**

Dental caries is a major health problem in U. S. preschool children, especially in low-income populations (13). Most implicated in this rampant disease process is prolonged use of baby bottles during the day or night, containing fermentable sugars, (e.g., fruit juice, soda, and other sweetened drinks), pacifiers dipped in sweet agents such as sugar, honey or syrups, or other high frequency sugar exposures (11). Solid foods such as cereal should not be put into a bottle for feeding; this is a form of force-feeding (14) and does not encourage the child to eat the cereal in a more developmentally-appropriate way.

Additional justification for the examples include:

- The American Academy of Pediatrics (AAP) and the American Academy of Pedodontics recommend that children not be put to bed with a bottle in their mouth (15, 16). While sleeping with a bottle in his or her mouth, a child's swallowing and salivary flow decreases, thus creating a pooling of liquid around the teeth (17). Propping the bottle can cause: ear infections because of fluid entering the middle ear and not draining properly; choking from liquid flowing into the lungs; and tooth decay from prolonged exposure to carbohydrate-containing liquids (18).
- Pediatric dentists recommend that parents be encouraged to have infants drink from a cup as they approach their first birthday, and that infants are weaned from the bottle by 12-14 months of age (19).
- The practice of allowing children to carry or drink from a bottle or cup of juice for periods throughout the day leads to excessive exposure of the teeth to carbohydrate, which promotes the development of dental caries (15). Allowing toddlers to use a bottle or cup containing fermentable carbohydrates unsupervised during waking hours provides an almost constant supply of carbohydrates and sugars (1). This leads to rapid demineralization of tooth enamel and an increase in the risk of dental caries due to prolonged contact between cariogenic bacteria on the susceptible tooth surface and the sugars in the consumed liquid (1, 19). The sugars in the liquid pool around the child's teeth and gums feed the bacteria there and result in decay (20). The process may start before the teeth are even fully erupted. Upper incisors (upper front teeth) are particularly vulnerable; the lower incisors are generally protected by the tongue (20). The damage begins as white lesions and progresses to brown or black discoloration typical of caries (20). When early childhood caries are severe, the decayed crowns may break off and the permanent teeth developing below may be damaged (20). Undiagnosed dental caries and other oral pain may contribute to feeding problems and failure to thrive in young children (20). Use of a bottle or cup, containing fermentable carbohydrates, without restriction is a risk because the more times a child consumes solid or liquid food, the higher the caries risk (1). Cariogenic snacks eaten between meals place the toddler most at risk for caries development; this includes the habit of continually sipping from cups (or bottles) containing cariogenic liquids (juice, milk, soda, or sweetened liquid) (20). If inappropriate use of the bottle persists, the child is at risk of toothaches, costly dental treatment, loss of primary teeth, and developmental lags on eating and chewing. If this continues beyond the usual weaning period, there is a risk of decay to permanent teeth.

425.4 Routinely using feeding practices that disregard the developmental needs or stages of the child.

The interactions and communication between a caregiver and child during feeding and eating influence a child's ability to progress in eating skills and consume a nutritionally adequate diet. These interactions comprise the "feeding relationship" (14). A dysfunctional feeding relationship, which could be characterized by a caregiver misinterpreting, ignoring, or overruling a young child's innate capability to regulate food intake based on hunger, appetite and satiety, can result in poor dietary intake and impaired growth (21, 22). Parents who consistently attempt to control their children's food intake may give children few opportunities to learn to control their own food intake (23). This could result in inadequate or excessive food intake, future problems with food regulation, and problems with growth and nutritional status. Instead of using approaches such as bribery, rigid control, struggles, or short-order cooking to manage eating, a healthier approach is for parents to provide nutritious, safe foods at regular meals and snacks, allowing children to decide how much, if any, they eat (1, 22). Young children should be able to eat in a matter-of-fact way sufficient quantities of the foods that are given to them, just as they take care of other daily needs (3). Research indicates that restricting access to foods (i.e., high fat foods) may enhance the interest of 3- to 5-year-old children in those foods and increase their desire to obtain and consume those foods. Stringent parental controls on child eating have been found to potentiate children's preference for high-fat energy dense foods, limit children's acceptance of a variety of foods, and disrupt children's regulation of energy intake (24, 25). Forcing a child to clean his or her plate may lead to overeating or development of an aversion to certain foods (12). The toddler and preschooler are striving to be independent (12). Self-feeding is important even though physically they may not be able to handle feeding utensils or have good eye-hand coordination (12). Children should be able to manage the feeding process independently and with dispatch, without either unnecessary dawdling or hurried eating (3, 17). Self-feeding milestones include (1): During infancy, older infants progress from semisolid foods to thicker and lumpier foods to soft pieces to finger-feeding table food (14). By 15 months, children can manage a cup, although not without some spilling. At 16 to 17 months of age, well-defined wrist rotation develops, permitting the transfer of feed from the bowl to the child's mouth with less spilling. The ability to lift the elbow as the spoon is raised and to flex the wrist as the spoon reaches the mouth follows. At 18 to 24 months, they learn to tilt a cup by manipulation with the fingers. Despite these new skills, 2-year-old children often prefer using their fingers to using the spoon. Preschool children learn to eat a wider variety of textures and kinds of food (3, 12). However, the foods offered should be modified so that the child can chew and swallow the food without difficulty (3).

425.5 Feeding foods to a child that could be contaminated with harmful microorganisms.

According to the AAP, to prevent food-borne illness, the foods listed below should not be fed to young children or infants (1). All of the foods have been implicated in selected outbreaks of food-borne illness, including in children. Background information regarding foods that could be contaminated with harmful microorganisms is also included below:

- Unpasteurized fruit or vegetable juice—Only pasteurized juice is safe for infants, children, and adolescents (15). Pasteurized fruit juices are free of microorganisms (15). Unpasteurized juice may contain pathogens, such as Escherichia coli, Salmonella, and Cryptosporidium organisms (15, 26). These organisms can cause serious disease, such as hemolytic-uremic syndrome, and should never be fed to infants and children (15). Unpasteurized juice must contain a warning on the label that the product may contain harmful bacteria (15, 27).

- Unpasteurized dairy products or soft cheeses such as feta, Brie, Camembert, blue-veined, and Mexican-style cheese—Young children or infants should not eat raw or unpasteurized milk or cheeses (1). Unpasteurized dairy products could contain harmful bacteria, such as Brucella species, that could cause young children to contract a dangerous food borne illness. The American Academy of Pediatrics also recommends that young children should not eat soft cheeses such as feta, Brie, Camembert, blue-veined, and Mexican-style cheese—these foods could contain Listeria bacteria (hard cheeses, processed cheeses, cream cheese, cottage cheese, and yogurt need not be avoided) (1).
- Raw or undercooked meat, fish, poultry, or eggs—Young children or infants should not eat raw or undercooked meat or poultry; or raw fish or shellfish, including oysters, clams, mussels, and scallops (1). These foods may contain harmful bacteria or parasites that could cause children to contract a dangerous foodborne illness.
- Raw vegetable sprouts (alfalfa, clover, bean, and radish)—Sprouts can cause potentially dangerous Salmonella and E. coli O157 infection. Sprouts grown under clean conditions in the home also present a risk because bacteria may be present in seed. Cook sprouts to significantly reduce the risk of illness (28).
- Deli meats, hot dogs, and processed meats (avoid unless heated until steaming hot)—These foods have been found to be contaminated with *Listeria monocytogenes*; if adequately cooked, this bacteria is destroyed.

425.6 Routinely feeding a diet very low in calories and/or essential nutrients.

Highly restrictive diets prevent adequate intake of nutrients, interfere with growth and development, and may lead to other adverse physiological effects (29). Well-balanced vegetarian diets with dairy products and eggs are generally associated with good health. However, strict vegan diets may be inadequate in calories, vitamin B12, vitamin D, calcium, iron, protein and essential amino acids needed for growth and development (30). The more limited the diet, the greater the health risk. Given the health and nutrition risks associated with highly restrictive diets, WIC can help the parent to assure that the child consumes an adequate diet to optimize health during critical periods of growth as well as for the long term.

425.7 Feeding dietary supplements with potentially harmful consequences.

A child consuming inappropriate or excessive amounts of single or multivitamin or mineral or herbal remedy not prescribed by a physician is at risk for a variety of adverse effects including harmful nutrient interactions, toxicity, and teratogenicity (1, 31). Like drugs, herbal or botanical preparations have chemical and biological activity, may have side effects, and may interact with certain medications--these interactions can cause problems and can even be dangerous (32). Botanical supplements are not necessarily safe because the safety of a botanical depends on many things, such as its chemical makeup, how it works in the body, how it is prepared, and the dose used (32). While some herbal teas may be safe, some have undesirable effects, particularly on young children who are fed herbal teas or who receive breast milk from mothers who have ingested herbal teas (33). Examples of teas with potentially harmful effects to children include: licorice, comfrey leaves, sassafras, senna, buckhorn bark, cinnamon, wormwood, woodruff, valerian, foxglove, pokeroot or pokeweed, periwinkle, nutmeg, catnip, hydrangea, juniper, Mormon tea, thorn apple, yohimbe bark, lobelia, oleander, Maté, kola nut or gotu cola, and chamomile (33-35).

425.8 Routinely not providing dietary supplements recognized as essential by national public health policy when a child's diet alone cannot meet nutrient requirements.

Depending on a child's specific needs and environmental circumstances, certain dietary supplements may be recommended by the child's health care provider to ensure health. For example, fluoride supplements may be of benefit in reducing dental decay for children living in fluoride-deficient areas (1, 36). In addition, the AAP recommends that children who are ingesting less than 1 liter (1 quart) per day of vitamin D-fortified formula or milk, should receive a vitamin D supplement of 400 IU/day (37). Since 1 quart of milk is in excess of the recommended 2 cups of milk per day for preschool children (38), most children will require a vitamin D supplement.

425.9 Routine ingestion by child of nonfood items (pica).

Pica is the compulsive eating of nonnutritive substances and can have serious medical implications (38). Pica is observed most commonly in areas of low socioeconomic status and is more common in women (especially pregnant women) and in children (35). Pica has also been seen in children with obsessive-compulsive disorders, mental retardation, sickle cell disease (39-41). Complications of this disorder include iron-deficiency anemia, lead poisoning, intestinal obstruction, acute toxicity from soil contaminants, and helminthic infestations (39, 42, 43).

References

1. Committee on Nutrition, American Academy of Pediatrics. Pediatric nutrition handbook. 6th ed. Elk Grove Village, Ill: American Academy of Pediatrics, 2009.
2. American Academy of Pediatrics, Committee on Nutrition: Iron fortification of infant formula. Pediatrics 1999; 104:119-123.
3. Trahms CM, Pipes PL, editors. Nutrition in Infancy and Childhood. WCB/McGraw-Hill; 1997.
4. Bellioni-Businco B, Paganelli R, Lucenti P, Giampietro PG, Perborn H, Businco L. Allergenicity of goat's milk in children with cow's milk allergy. J Allergy Clin Immunol 1999;103:1191-1194.
5. Tamborlane, WV, editor. The Yale guide to children's nutrition. Connecticut: Yale University; 1997.
6. Hansen AE, Wiese HF, Boelsche AN, Haggard ME, Adam DJ, Davis H. Role of Linoleic Acid in Infant Nutrition: Clinical and Chemical Study of 428 Infants Fed on Milk Mixtures Varying in Kind and Amount of Fat. Pediatrics. 1963; 31(1), 171-19.
7. Uauy R, Castillo C. Lipid requirements of infants: implications for nutrient composition of fortified complementary foods. The Journal of Nutrition. 2003; 133(9), 2962S-2972S.
8. Innis SM. Dietary (n-3) fatty acids and brain development. The Journal of Nutrition. 2007; 137(4), 855-859.
9. Birch EE, Garfiedl S, Castaneda Y, Hughbanks-Wheaton D, Uauy R, Hoffman D. Visual acuity and cognitive outcomes at 4 years of age in a double-blind, randomized trial of long-chain polyunsaturated fatty acid-supplemented infant formula. Early human development. 2007; 83(5), 279-284.
10. Daniels, SR, Greer, FR. Lipid screening and cardiovascular health in childhood. Pediatrics. 2008; 122(1), 198-208.

11. Tinanoff N, Palmer CA. Dietary determinants of dental caries and dietary recommendations for preschool children. J Public Health Dent 2000;60(3):197-206.
12. Williams, CP, editor. Pediatric manual of clinical dietetics. Chicago: American Dietetic Association; 1998.
13. Tang J, Altman DS, Robertson D, O'Sullivan DM, Douglass JM, Tinanoff N. Dental caries prevalence and treatment levels in Arizona preschool children. Public Health Rep 1997;112:319-29.
14. Satter E. Child of mine: Feeding with love and good sense. Palo Alto (CA): Bull Publishing Company; 2000.
15. American Academy of Pediatrics Committee on Nutrition: The use and misuse of fruit juice in pediatrics. Pediatrics 2001;107:1210-1213.
16. American Academy of Pediatrics and American Academy of Pedodontics. Juice in ready-to-use bottles and nursing bottle carries. AAP News. 1978;29(1):11.
17. Samour PQ, Helm KK, Lang CE. Handbook of pediatric nutrition. 2nd ed. Gaithersburg, MD: Aspen Publishers, Inc.; 1999.
18. Shelov SD. Caring for your baby and young child: birth to age 5. Elk Grove Village, IL: American Academy of Pediatrics; 1998.
19. American Academy of Pediatric Dentistry. Baby Bottle Tooth Decay/Early Childhood Caries. Pediatr Dent 2000-2001 (revised May 1996); 2001 Mar-Apr;23(2):18
20. Fitzsimons D, Dwyer JT, Palmer C, Boyd LD. Nutrition and oral health guidelines for pregnant women, infants, and children. J Am Diet Assoc. Feb 1998;98(2):182-6.
21. Satter, E. Childhood feeding problems. Feelings and Their Medical Significance; Vol. 32, no. 2; Columbus, OH; Ross Laboratories; 1990.
22. Satter EM. The feeding relationship. J.Am.Diet.Assoc. 1986;86:352-6.
23. Johnson SL, Birch LL. Parents' and children's adiposity and eating style. Pediatrics 1994;94:653-61.
24. Olson RE. Is it wise to restrict fat in the diets of children? J Am Diet Assoc 2000 Jan;100(1):28-32.
25. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. Pediatrics 1998;101:539-549.
26. Parish ME. Public health and nonpasteurized fruit juices. Crit Rev Microbiol. 1997;23:109-119.
27. Food Labeling. Warning and Notice Statement: Labeling of Juice Products; Final Rule. 63 Federal Register 37029-37056 (1998) (codified at 21 CFR §101, 120).
28. Food and Drug Administration. Updates: Avoid Raw Sprouts to Reduce Food Poisoning Risk, Agency Advises. FDA Consumer magazine, September-October 1999 (see: http://www.fda.gov/fdac/departs/1999/599_upd.html)
29. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
30. Duyff RL. American Dietetic Association. The American Dietetic Association's complete food and nutrition guide. Minneapolis, MN: Chronimed Pub; 1996.
31. Anderson JV, Van Nierop MR. Basic nutrition facts a nutrition reference. Lansing, MI: Michigan Department of Public Health; 1989.
32. Office of Dietary Supplements, National Institutes of Health (NIH). Botanical Dietary Supplements: Background Information. NIH web page, last updated 7/7/2004, see: http://ods.od.nih.gov/factsheets/BotanicalBackground.asp
33. Lawrence, RA. Breastfeeding: a guide for the medical profession. 5th edition. St. Louis, MO: Mosby, 1999, pp. 371-377.
34. Siegel RK. Herbal intoxication: psychoactive effects from herbal cigarettes, tea and capsules. JAMA 236:473, 1976.

35. Ridker PM. Toxic effects of herbal teas. Arch Environ Health 42(3):133-6, 1987.
36. American Academy of Pediatric Dentistry. Fluoride. Pediatr Dent. 1999;21:40.
37. American Academy of Pediatrics Section on Breastfeeding and Committee On Nutrition, Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. Pediatrics 2008; [cited 2015 Feb 27]. Available from: www.pediatrics.org/cgi/doi/10.1542/peds.2008-1862.
38. U.S. Department of Agriculture, Center for Nutrition Policy and Promotion. Health and Nutrition Information for Preschoolers. [cited 2015 Feb 27]; Available from: http://www.choosemyplate.gov/preschoolers.html.
39. Rose EA, Porcerelli JH, Neale AV. Pica: common but commonly missed. J Am Board Fam Pract. 2000;13(5):353-8.
40. LeBlanc LA, Piazza CC, Krug MA. Comparing methods for maintaining the safety of a child with pica. Res Dev Disabil. 1997; 18(3):215-20.
41. Ivascu NS, et al. Characterization of pica prevalence among patients with sickle cell disease. Arch Pediatr Adoles Med 2001; 155(11):1243-7.
42. Calabrese EJ, et al. Soil ingestion: a concern for acute toxicity in children. Environ Health Perspect. 1997; 105(12):1354-8.
43. Wang PY, Skarsgard ED, Baker RJ. Carpet bezoar obstruction of the small intestine. J Pediatr Surg. 1996; 31(12):1691-3.

Nutrition Risk Criteria **427 INAPPROPRIATE NUTRITION PRACTICES (WOMEN)**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	-
BREASTFEEDING WOMEN	4	-
NON-BREASTFEEDING WOMEN	6	-

Definition

FORMERLY QUESTIONABLE DIET FOR WOMEN

Routine nutrition practices that may result in impaired nutrient status, disease, or health problems. These practices with examples are outlined below. Refer to "Justification and References" for this criterion.

427.1 Consuming dietary supplements with potentially harmful consequences.

Examples of dietary supplements which when ingested in excess of recommended dosages, may be toxic or have harmful consequences:

- Single or multiple vitamins
- Mineral supplements
- Herbal or botanical supplements/remedies/teas

427.2 Consuming a diet very low in calories and/or essential nutrients; or impaired caloric intake or absorption of essential nutrients following bariatric surgery.

- Strict vegan diet
- Low-carbohydrate, high-protein diet
- Macrobiotic diet
- Any other diet restricting calories and/or essential nutrients

427.3 Compulsively ingesting non-food items (pica).

Non-food items:

- Ashes
- Baking soda
- Burnt matches
- Carpet fibers
- Chalk
- Cigarettes
- Clay
- Dust
- Large quantities of ice and/or freezer frost
- Paint chips
- Soil
- Starch (laundry and cornstarch)

427.4 Inadequate vitamin/mineral supplementation recognized as essential by national public health policy.

- Consumption of less than 27 mg of iron as a supplement daily by pregnant woman
- Consumption of less than 150 mcg of supplemental iodine per day by pregnant and breastfeeding women
- Consumption of less than 400 mcg of folic acid from fortified foods and/or supplements daily by non-pregnant woman

427.5 Pregnant woman ingesting foods that could be contaminated with pathogenic microorganisms.

Potentially harmful foods:

- Raw fish or shellfish, including oysters, clams, mussels, and scallops
- Refrigerated smoked seafood, unless it is an ingredient in a cooked dish, such as a casserole
- Raw or undercooked meat or poultry
- Hot dogs, luncheon meats (cold cuts), fermented and dry sausage and other deli-style meat or poultry products unless reheated until steaming hot
- Refrigerated pâté or meat spreads
- Unpasteurized milk or foods containing unpasteurized milk
- Soft cheeses such as feta, Brie, Camembert, blue-veined cheeses and Mexican style cheese such as queso blanco, queso fresco, or Panela unless labeled as made with pasteurized milk
- Raw or undercooked eggs or foods containing raw or lightly cooked eggs including certain salad dressings, cookie and cake batters, sauces, and beverages such as unpasteurized eggnog
- Raw sprouts (alfalfa, clover, and radish)
- Unpasteurized fruit or vegetable juices

Required Documentation

Document which behaviors are present.

Justification

427.1 Consuming dietary supplements with potentially harmful consequences.

Women taking inappropriate or excessive amounts of dietary supplements such as, single or multivitamins or minerals, or botanical (including herbal) remedies or teas, are at risk for adverse effects such as harmful nutrient interactions, toxicity and teratogenicity (1, 2). Pregnant and lactating women are at higher risk secondary to the potential transference of harmful substances to their infant.

Most nutrient toxicities occur through excessive supplementation of particular nutrients, such as, vitamins A, B-6 and niacin, iron and selenium (3). Large doses of vitamin A may be teratogenic (4). Because of this risk, the Institute of Medicine recommends avoiding preformed vitamin A supplementation during the first trimester of pregnancy (4). Besides nutrient toxicities, nutrient-nutrient and drug-nutrient interactions may adversely affect health.

Many herbal and botanical remedies have cultural implications and are related to beliefs about pregnancy and breastfeeding. The incidence of herbal use in pregnancy ranges from 7% to 55% with echinacea and ginger being the most common (1). Some botanical (including herbal) teas may be safe; however, others have undesirable effects during pregnancy and breastfeeding. Herbal supplements such as, blue cohash and pennyroyal stimulate uterine contractions, which may increase the risk of miscarriage or premature labor (1, 5). The March of Dimes and the American Academy of Pediatrics recommend cautious use of tea mixtures because of the lack of safety testing in pregnant women (6).

427.2 Consuming a diet very low in calories and /or essential nutrients; or impaired caloric intake or absorption of essential nutrients following bariatric surgery.

Women consuming highly restrictive diets are at risk for primary nutrient deficiencies, especially during critical developmental periods such as pregnancy. Pregnant women who restrict their diets may increase the risk of birth defects, suboptimal fetal development and chronic health problems in their children. Examples of nutrients associated with negative health outcomes are:

- Low iron intake and maternal anemia and increased risk of preterm birth or low
  birth weight (7, 8).
- Low maternal vitamin D status and depressed infant vitamin D status (9)
- Low folic acid and NTD (10, 11, 12)

Low calorie intake during pregnancy may lead to inadequate prenatal weight gain, which is associated with infant intrauterine growth restriction (IUGR) (13) and birth defects (10, 11, 14). The pregnant adolescent who restricts her diet is of particular concern since her additional growth needs compete with the developing fetus and the physiological changes of pregnancy (14).

Strict vegan diets may be highly restrictive and result in nutrient deficiencies. Nutrients of potential concern that may require supplementation are:

- Riboflavin (15, 16)
- Iron (15)
- Zinc (15, 17)
- Vitamin B12 (15, 16, 18)
- Vitamin D (15, 16, 18)
- Calcium (15, 16, 18, 19)
- Selenium (16)

The pregnant adolescent who consumes a vegan diet is at even greater risk due to her higher nutritional needs (16, 18). The breastfeeding woman who chooses a vegan or macrobiotic diet increases her risk and her baby's risk for vitamin B12 deficiency (18). Severe vitamin B12 deficiency resulting in neurological damage has been reported in infants of vegetarian mothers (18).

With the epidemic of obesity, treatment by gastric bypass surgery has increased more than 600% in the last ten years and has created nutritional deficiencies not typically seen in obstetric or pediatric medical practices (20). Gastrointestinal surgery promotes weight loss by restricting food intake and, in some operations, interrupting the digestive process. Operations that only reduce stomach size are known as "restrictive operations" because they restrict the amount of food the stomach can hold. Examples of restrictive operations are adjustable gastric banding and vertical banded gastroplasty. These types of operations do not interfere with the normal digestive process (21).

Some operations combine stomach restriction with a partial bypass of the small intestine, these are known as malabsorptive operations. Examples of malabsorptive operations are Roux-en-y gastric bypass (RGB) and Biliopancreatic diversion (BPD). Malabsorptive operations carry a greater risk for nutritional deficiencies because the procedure causes food to bypass the duodenum and jejunum, where most of the iron and calcium are absorbed. Menstruating women may develop anemia because not enough iron and vitamin

B12 are absorbed. Decreased absorption of calcium may also contribute to osteoporosis and metabolic bone disease (21). A breastfeeding woman who has had gastric bypass surgery is at risk of vitamin B12 deficiency for herself and her infant (22).

427.3 Compulsively ingesting non-food items (pica).

Pica, the compulsive ingestion of non-food substances over a sustained period of time, is linked to lead poisoning and exposure to other toxicants, anemia, excess calories or displacement of nutrients, gastric and small bowel obstruction, as well as, parasitic infection (23). It may also contribute to nutrient deficiencies by either inhibiting absorption or displacing nutrient dense foods in the diet.

Poor pregnancy outcomes associated with pica-induced lead poisoning, include lower maternal hemoglobin level at delivery (24) and a smaller head circumference in the infant (25). Maternal transfer of lead via breastfeeding has been documented in infants and can result in a neuro-developmental insult depending on the blood lead level and the compounded exposure for the infant during pregnancy and breastfeeding (26, 27, 28).

427.4 Inadequate vitamin/mineral supplementation recognized as essential by national public health policy.

The Recommended Dietary Allowance (RDA) for pregnant women is 27 mg of iron per day (29). The Centers for Disease Control and Prevention recommends iron supplementation for all pregnant women to prevent iron deficiency (30); however, pregnant women should seek guidance from a qualified health care provider before taking dietary supplements (31).

During pregnancy and lactation the iodine requirement is sharply elevated. The RDA for iodine during pregnancy is 220 mcg and 290 mcg during lactation (29). Severe iodine deficiency during pregnancy can cause cretinism and adversely affect cognitive development in children (32). Even mild iodine deficiency may have adverse affects on the cognitive function of children (33). Since the 1970s, according to the 2001-2002 National Health and Nutrition Examination Surveys (NHANES), there has been a decrease of approximately 50% in adult urinary iodine values. For women of child bearing age, the median urinary iodine value decreased from 294 to 128 mcg per liter (34). The American Thyroid Association recommends that women receive prenatal vitamins containing 150 mcg of iodine daily during pregnancy and lactation (35). The iodine content of prenatal vitamins in the United States is not mandated, thus not all prenatal vitamins contain iodine (36). Pregnant and breastfeeding women should be advised to review the iodine content of their vitamins and discuss the adequacy of the iodine with their health care provider.

Non-pregnant women of childbearing age who do not consume adequate amounts of folic acid are at greater risk for functional folate deficiency, which has been proven to cause neural tube defects (NTDs), such as, spina bifida and anencephaly (37-40).

Folic acid consumed from fortified foods and/or a vitamin supplement in addition to folate found naturally in food reduces this risk (12). The terms “folic acid” and “folate” are used interchangeably, yet they have different meanings. Folic acid is the synthetic form used in vitamin supplements and fortified foods (12, 38, 39). Folate occurs naturally and is found in foods, such as dark green leafy vegetables, strawberries, and orange juice (12).

Studies show that consuming 400 mcg of folic acid daily interconceptionally can prevent 50 percent of neural tube defects (12). Because NTDs develop early in pregnancy (between the 17th and 30th day) and many pregnancies are not planned, it is important to have adequate intakes before pregnancy and throughout the childbearing years (14). NTDs often occur before women know they are pregnant. It is recommended that all women capable of becoming pregnant consume a multivitamin containing 400 mcg of folic acid daily (39-41). It is important that breastfeeding and non-breastfeeding women participating in the WIC Program know about folic acid and foods that contain folate to encourage preconceptional preventive practices (38).

427.5 Pregnant woman ingesting foods that could be contaminated with pathogenic microorganisms.

Foodborne illness is a serious public health problem (42). The causes include pathogenic microorganisms (bacteria, viruses, and parasites) and their toxins and chemical contamination. The symptoms are usually gastrointestinal in nature (vomiting, diarrhea, and abdominal pain), but neurological and “non-specific” symptoms may occur as well. Over the last 20 years, certain foods have been linked to outbreaks of food-borne illness. These foods include: milk (Campylobacter); shellfish (Norwalk-like viruses), unpasteurized apple cider (Escherichia coli O 157:H7); eggs (Salmonella); fish (ciguatera poisoning); raspberries (Cyclospora); strawberries (Hepatitis A virus); and ready-to-eat meats (Listeria monocytogenes).

Listeria monocytogenes can cause an illness called listeriosis. Listeriosis during pregnancy can result in premature delivery, miscarriage, fetal death, and severe illness or death of a newborn from the infection (43). Listeriosis can be transmitted to the fetus through the placenta even if the mother is not showing signs of illness.

Pregnant women are especially at risk for food-borne illness. For this reason, government agencies such as the Centers for Disease Control and Prevention, the USDA Food Safety and Inspection Service, and the Food and Drug Administration advise pregnant women and other high risk individuals not to eat foods as identified in the definition for this criterion (42, 43).

The CDC encourages health care professionals to provide anticipatory guidance, including the “four simple steps to food safety” of the Fight BAC campaign, to help reduce the incidence of food-borne illnesses.

References

1. Tiran D. The use of herbs by pregnant and childbearing women: a risk-benefit assessment. Complementary Therapies in Nursing and Midwifery. November 2003. 9(4):176-181.
2. Position of the American Dietetic Association: Nutrition and lifestyle for a healthy pregnancy outcome. J Am Diet Assoc. 2002 October;102(10):1479-1490.
3. Position of the American Dietetic Association: Food fortification and dietary supplements. J Am Diet Assoc. January 2001.
4. Langkamp-Henken B, Lukowski MJ, Turner RE, Voyles LM. High levels of retinol intake during the first trimester of pregnancy result from use of over-the-counter vitamin/mineral supplements. J Am Diet Assoc. September 2000.

5. March of Dimes (homepage on the Internet). New York: Herbal Supplements: heir Safety, a Concern for Health Care Providers. [cited May 26, 2004] Available from: http://www.marchofdimes.com
6. American Academy of Pediatrics, Committee on Nutrition. Pediatric Nutrition Handbook. 5th ed. Kleinman, Ronald, editor. Washington DC: American Academy of Pediatrics; 2004.
7. Recommendations to prevent and control iron deficiency in the United States. MMWR [serial on the Internet]. 1998 April [cited 2004 March 12]. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/00051880.htm.
8. Rasmussen, K. M. Is there a causal relationship between iron deficiency or iron-deficiency anemia and weight at birth, length of gestation and perinatal mortality? American Society for Nutritional Sciences. 2001;590S-603S.
9. Scanlon KS, editor. Vitamin D expert panel meeting; October 11-12, 2001; Atlanta, Georgia. Available from: http://www.cdc.gov/nccdphp/dnpa/nutrition/pdf/Vitamin D Expert Panel Meeting.pdf.
10. Carmichael SL, Shaw GM, Schaffer DM, Selvin S. Diet quality and risk of neural tube defects. Medical Hypotheses. 2003;60(3):351-355.
11. Shaw GM, Todoroff K, Carmichael SL, Schaffer DM, Selvin S. Lowered weight gain during pregnancy and risk of neural tube defects among offspring. Int J Epidemiol. 2001; 30:60-65.
12. American Academy of Pediatrics, Committee on Genetics. Folic acid for the prevention of neural tube defects. Pediatrics.1999; 104(2):325-327.
13. Strauss RS, Dietz WH. Low maternal weight gain in the second and third trimester increases the risk for intrauterine growth retardation. American Society for Nutritional Sciences. 1999; 988-993.
14. Scholl TO, Hediger ML, Ances IG. Maternal growth during pregnancy and decreased infant birth weight. Am J Clin Nutr. 1990;51:790-793.
15. Position of the American Dietetic Association and Dietitians of Canada: Vegetarian diets. J Am Diet Assoc. 2003; 103(6):748-765.
16. Larsson CL, Johansson GK. Dietary intake and nutritional status of young vegans and omnivores in Sweden. Am J Clin Nutr. 2002; 76:100-106.
17. Bakan R, Birmingham CL, Aeberhardt L, Goldner EM. Dietary zinc intake of vegetarian and nonvegetarian patients with anorexia nervosa. International Journal of Eating Disorders. 1993;13(2):229-233.
18. Specker, Bonny L., Nutritional concerns of lactating women consuming vegetarian diets. Am J Clin Nutr. 1994:59(suppl):1182-1186.
19. Heaney RP, Dowell MS, Rafferty K, Bierman J. Bioavailability of the calcium in fortified soy imitation milk, with some observation on method. Am J Clin Nutr. 2000;71:1166-1169.
20. Steinbrook, R. Surgery for severe obesity. N Engl J Med. 2004; 350(11):1075-9.
21. National Institute of Diabetes and Digestive and Kidney Diseases. Gastrointestinal Surgery for Severe Obesity. [cited August 18, 2004] Available from: http://www.niddk.nih.gov/health/nutrit/pubs/gastric/gastricsurgery.htm
22. Grange DK, Finlay JL. Nutritional vitamin B12 deficiency in a breastfed infant following maternal gastric bypass. Pediatr Hematol Oncol. 1994; 11(3):311-8.
23. Corbett RW, Ryan C, Weinrich SP. Pica in pregnancy: does it affect pregnancy outcomes? American Journal of Maternal and Child Nursing. 2003;28(3):183-189.
24. Rainville AJ. Pica practices of pregnant women are associated with lower maternal hemoglobin level at delivery. J Am Diet Assoc. 1998;98(3): 293-6.
25. Institute of Medicine: WIC Nutrition Risk Criteria: A Scientific Assessment; 1996; pp.270-272.

26. Gulson, Brian L., et. al., Relationships of lead in breast milk to lead in blood, urine, and diet of infant and mother. Environmental Health Perspectives. 1998:106(10): 667-674.
27. Ping-Jian L, Ye-Zhou S, Qian-Ying W, Li-Ya G, Yi-Land W. Transfer of lead via placenta and breast milk in human. Biomedical and Environmental Sciences. 2000; 13:85-89.
28. Canfield, RL, Henderson, C, Cory-Slecha, D, Cox, C, Jusko, T, Lanphear, B. Intellectual impairment in children with blood lead concentrations below 10 mcg per deciliter. N Engl J Med. 2003;348(16):1517-1526.
29. Institute of Medicine. Dietary reference intakes for vitamin A, vitamin K, arsenic, Boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. Food and Nutrition Board, Washington, DC: National Academy Press; 2001.
30. Centers for Disease Control and Prevention. Recommendations to prevent and control iron deficiency in the United States. MMWR 1998:RR-3.
31. U.S. National Library of Medicine and National Institutes of Health. Drugs and supplements: iron. Medline Plus. http://www.nlm.nih.gov/medlineplus/druginfo/natural/patient-iron.html#Satety. Accessed May 2009.
32. Zimmerman MB. Iodine deficiency in pregnancy and effects of maternal iodine supplementation on the offspring: a review. Am J Clin Nutr 2009;8 (suppl:668S-72S).
33. de Escobar DM, Obregon MJ, del Rey FF. maternal thyroid hormones early in pregnancy and fetal brain development. Best Pract Res Clin Endoriconl Metab 2004;18:225-48.
34. Caldwell KL, Miller GA, Wang RY, Jain RB, Jones, RL. Iodine status of the U.S. population, National Health and Nutrition Examination Survey 2003-2004. Thyroid 2008;18:1207-14.
35. Becker DV, Braverman LE, Delange F, et al. Iodine supplementation for pregnancy and lactation – United States and Canada: recommendations of the American Thyroid Association. Thyroid 2006;16:949-51.
36. Leung AM, Pearce EN, Braverman, LE. Iodine content of prenatal vitamins in the United States. N Engl J Med 2009;360-9.
37. Centers for Disease Control and Prevention, Division of Birth Defects and Developmental Disabilities. Folic acid and the prevention of spina bifida and anencephaly: 10 years after the U.S. Public Health Service recommendation. MMWR 2002;51:(RR-13)1-3.
38. Centers for Disease Control and Prevention. National Center for Environmental Health, Division of Birth Defects and Developmental Disabilities. Preventing neural tube birth defects: a prevention model and resource guide. Atlanta: CDC, 1998.
39. Centers for Disease Control and Prevention. Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. MMWR 1992;41:RR-14.
40. Evans MI, Llurba E, Landsberger EJ, O'Brien JE, Harrison HH. Impact of folic acid fortification the Untied States: markedly diminish high maternal serum alpha-fetoprotein values. Am Col Obstetr Gynecol. 2004;103(3):447.
41. Chacko MR, Anding R, Kozinetz CA, Grover JL. Neural tube defects: knowledge and preconceptional prevention practices in minority young women. Pediatrics. 2003;112(3):536-542.
42. Centers for Disease Control and Prevention, Diagnosis and Management of Foodborne Illness, A Primer for Physicians. MMWR 2001;50:RR-2.
43. Food Safety and Inspection Service, USDA. Listeriosis and Pregnancy: What is Your Risk? [cited August 11, 2004] Available from: http://www.fsis.usda.gov.

Websites for additional information:

427.1 References - Supplements/Herbs
http://www.marchofdimes.com
http://www.dietary-supplements.info.nih.gov/
http://www.vm.cfsan.fda.gov/
http://www.herbalgram.org

427.2 References - Highly Restrictive Eating/ Nutrient Malabsorption
http://www.eatright.org
http://www.nimh.nih.gov
http://www.eatright.org/
http://www.llu.edu/llu/vegetarian/
http://www.nal.usda.gov/fnic/pubs/bibs/gen/vegetarian.htm
http://www.gastric-bypass-treatment.com/long-term-weight-loss-surgery-compications.aspx

427.3 References - Non-Food Ingestion
http://www.nieh.nih.gov/
http://www.epa.gov/

427.4 References - Folic Acid
http://www.cdc.gov/
http://www.aap.org/
http://www.iom.edu/

427.5 References - Listeriosis
http://www.cdc.gov/foodsafety
http://www.cdc.gov/ncidod/dbmd/diseaseinfo/listeriosis_g.htm
http://www.cfsan.fda.gov
http://www.foodsafety.gov
http://www.fightbac.org
http://www.ific.org

Nutrition Risk Criteria **428** **DIETARY RISK ASSOCIATED WITH COMPLEMENTARY FEEDING PRACTICES**

Category, Priority and Referral

Category	Priority	Referral
INFANTS 4 TO 12 MONTHS	4	-
CHILDREN 12 THROUGH 23 MONTHS	5	-

Definition

An infant or child who has begun to or is expected to begin to 1) consume complementary foods and beverages, 2) eat independently, 3) be weaned from breast milk or infant formula, or 4) transition from a diet based on infant/toddler foods to one based on the *Dietary Guidelines for Americans*, is at risk of inappropriate complementary feeding.

A complete nutrition assessment, including for risk #411, Inappropriate Nutrition Practices for Infants, or #425, Inappropriate Nutrition Practices for Children, must be completed prior to assigning this risk.

Required Documentation

Complete Nutrition Assessment

Justification

Overview

Complementary feeding is the gradual addition of foods and beverages to the diet of the infant and young child. (1, 2) The process of adding complementary foods should reflect the physical, intellectual, and behavioral stages as well as the nutrient needs of the infant or child. Inappropriate complementary feeding practices are common and well documented in the literature. Caregivers often do not recognize signs of developmental readiness and, therefore, offer foods and beverages that may be inappropriate in type, amount, consistency, or texture. Furthermore, a lack of nationally accepted feeding guidelines for children under the age of two might lead caregivers to assume that all foods are suitable for this age range.

The 2000 WIC Participant and Program Characteristics study (PC 2000) shows greater percentages of anthropometric and biochemical risk factors in children ages 6 to 24 months than in children 24 to 60 months of age. (3)
These differences could reflect physical manifestations of inappropriate complementary feeding practices. Although PC 2000 shows a lower dietary risk in the 6 to 24 month age group, this risk is probably under-reported due to the high incidence of other higher priority nutrition risks.

AGE	ANTHOPOMETRIC RISK (%)	BIOCHEMICAL RISK (%)	DIETARY RISK (%)
6-11 MOS	40	16	55
1 YEAR	41	14	76
2 YEAR	37	12	80
3 YEAR	32	9	80
4 YEAR	35	7	79

The Institute of Medicine (IOM), in their report, *Summary of Proposed Criteria for Selecting the WIC Food Packages* identified specific nutrients with potential for inadequacy or excess for WIC participants. For breast-fed infants 6 through 11 months, the nutrients of concern for potential inadequacy are iron and zinc while those for children 12 through 23 months are iron, vitamin E, fiber and potassium. The nutrients of concern for excessive intake in children 12 through 23 months are zinc, preformed vitamin A, sodium and energy. (4)

To manage complementary feeding successfully, caregivers must make decisions about what, when, where, and how to offer foods according to the infant's or child's:

- Requirement for energy and nutrients;
- Fine, gross, and oral motor skills;
- Emerging independence and desire to learn to self-feed; and
- Need to learn healthy eating habits through exposure to a variety of nutritious foods. (1, 2, 5, 6, 7)

How WIC Can Help

The WIC Program plays a key role not only in the **prevention** of nutrition-related health problems, but also in the **promotion** of lifelong healthy eating behaviors. The process of introducing complementary foods provides a unique opportunity for WIC staff to assist caregivers in making appropriate feeding decisions for young children that may have lifelong implications.

Prevention of Nutrition-Related Health Problems

- Zinc deficiency: Zinc is critical for growth and immunity, as well as brain development and function. The concentration of zinc in breast milk declines to a level considered inadequate to meet the needs of infants 7 to 12 months of age. (8, 9) Complementary food sources of zinc, such as meats or zinc-fortified infant cereals, should be introduced to exclusively breastfed infants by 7 months.

- Iron deficiency: Hallberg states, "The weaning period in infants is especially critical because of the especially high iron requirements and the importance of adequate iron nutrition during this crucial period of development."(10) According to the Centers for Disease Control and Prevention (CDC), children less than 24 months of age, especially those between 9 and 18 months, have the highest rate of iron deficiency of any age group. (11) In the third National Health and Nutrition Examination Survey (NHANES III), children ages 1 to 2, along with adolescent girls, had the highest rates of overt anemia, while 9 % were iron deficient. (12) Meanwhile, the Pediatric Nutrition Surveillance 2003 Report noted anemia rates of 16.2 % in 6 to 11 month-old infants, 15.0 % in 12 to 17 month-olds, and 13.5 % in 18-23 month old children. (13)

  Picciano et al. reported that the intake of iron decreased from 98% of the recommended amount at 12 months to 76% at 18 months of age. (14) In WIC clinics, Kahn et al. found that the incidence of anemia was significantly more common in 6 to 23 month old children than in 23 to 59 month-olds. The 6 to 23 month-old was also more likely than the older child to develop anemia despite a normal hemoglobin test at WIC certification. (15)

Clarification

Feeding practices that may prevent iron deficiency include:

- Breastfeeding infants exclusively until 4 to 6 months of age;
- Feeding only iron-fortified infant formula as a substitute for or supplement to breast milk until age 1;
- Offering a supplemental food source of iron to infants, between 4 to 6 months or when developmentally ready;
- Avoiding cow's milk until age 12 months; and
- Limiting milk consumption to no more than 24 oz per day for children aged 1 to 5 years. (11)

- Obesity: Much of the literature on obesity indicates that learned behaviors and attitudes toward food consumption are major contributing factors. Proskitt states, "The main long term effect of weaning on nutritional status could be through attitudes toward food and meals learned by infants through the weaning process. This may be a truly critical area for the impact of feeding on later obesity." (16)

  Birch and Fisher state, "An enormous amount of learning about food and eating occurs during the transition from the exclusive milk diet of infancy to the omnivore's diet consumed by early childhood." The authors believe that parents have the greatest influence on assuring eating behaviors that help to prevent future overweight and obesity. (17)

  The American Academy of Pediatrics (AAP) states, "…prevention of overweight is critical, because long-term outcome data for successful treatment approaches are limited…" and, "Families should be educated and empowered through anticipatory guidance to recognize the impact they have on their children's development through lifelong habits of physical activity and nutritious eating." (1) Parents can be reminded that they are role models and teachers who help their children adopt healthful eating and lifestyle practices.

- Tooth decay: Children under the age of 2 are particularly susceptible to Early Childhood Caries (ECC), a serious public health problem. (18) In some communities, the incidence of ECC can range from 20% to 50%. (19) Children with ECC appear to be more susceptible to caries in permanent teeth at a later age. (1, 20)

  Dental caries can be caused by many factors, including prolonged use of a bottle and extensive use of sweet and sticky foods. (21)
  The Avon Longitudinal Study of Pregnancy and Childhood examined 1,026 children aged 18 months and found that baby bottles were used exclusively for drinking by 10 % of the children and for at least one feeding by 64% of the children. Lower income families were found to use the bottle more frequently for carbonated beverages than higher income families. (22)

  Complementary feeding practices that caregivers can use to prevent oral health problems include:
  - Avoiding concentrated sweet foods like lollipops, candy and sweetened cereals
  - Avoiding sweetened beverages. Introducing fruit juice after 6 months of age (1) and only feeding it in a cup; and limiting fruit juice to 4-6 oz/day
  - Weaning from a bottle to a cup by 12 to 14 months (23)

Promotion of Lifelong Healthy Eating Behaviors

- Timing of introduction of complementary foods:
  The AAP, Committee on Nutrition (CON) states that, "… complementary foods may be introduced between ages 4 and 6 months…" but cautions that actual timing of introduction of complementary foods for an individual infant may differ from this (population based) recommendation. Furthermore, the AAP-CON acknowledges a difference of opinion with the AAP, Section on Breastfeeding, which recommends exclusive breastfeeding for at least 6 months. (1)

  Early introduction of complementary foods before the infant is developmentally ready (i.e., before 4-6 months of age) is associated with increased respiratory illness, allergy in high-risk infants, and decreased breast milk production (7).

  Infants with a strong family history of food allergy should be breastfed for as long as possible and should not receive complementary foods until 6 months of age. The introduction of the major food allergens such as eggs, milk, wheat, soy, peanuts, tree nuts, fish and shellfish should be delayed until well after the first year of life as guided by the health care provider. (7, 24)

  Delayed introduction of complementary foods, on the other hand, is also associated with feeding difficulties. Northstone et al found that introduction of textured foods after 10 months of age resulted in more feeding difficulties later on, such as picky eating and/or refusal of many foods. To avoid these and other developmental problems, solid foods should be introduced no later than 7 months, and finger foods between 7 and 9 months of age. (25)

- Choosing Appropriate Complementary Foods and Beverages:
  Complementary foods should supply essential nutrients and be developmentally appropriate. (7) The WIC Infant Feeding Practices Study (WIC-IFPS) found that by 6 months of age, greater than 80% of mothers introduced inappropriate dairy foods (i.e., yogurt, cheese, ice cream and pudding), 60% introduced sweets/snack foods (defined as chips, pretzels, candy, cookies, jam and honey), and 90% introduced high protein foods (beans, eggs and peanut butter) to their infants. This study also found that, among the infants who received supplemental drinks by 5 months of age, three-quarters had never used a cup, concluding that most infants received supplemental drinks from the bottle. By one year of age, almost 90% of WIC infants received sweetened beverages and over 90% received sweet/snack foods. (26)

  The Feeding Infants and Toddlers Study (FITS) found that WIC infants and toddlers consumed excess energy but inadequate amounts of fruits and vegetables. In addition, WIC toddlers consumed more sweets, desserts and sweetened beverages than non-WIC toddlers. (27)

  Sixty-five percent of all food-related choking deaths occur in children under the age of 2. Children in this age group have not fully developed their oral-motor skills for chewing and swallowing. For this reason, they should be fed foods of an appropriate consistency, size, and shape. Foods commonly implicated in choking include hot dogs, hard, gooey or sticky candy, nuts and seeds, chewing gum, grapes, raisins, popcorn,
  peanut butter and hard pieces of raw fruits and vegetables and chunks of meat or cheese. (1, 28, 29)

- Introducing a cup: Teaching an infant to drink from a cup is part of the process of acquiring independent eating skills. A delay in the initiation of cup drinking prolongs the use of the nursing bottle that can lead to excess milk and juice intake and possible Early Childhood Caries (ECC). Weaning from a bottle to a cup should occur by 12 to14 months of age. (23)

- Helping the child establish lifelong healthy eating patterns:
  Lifelong eating practices may have their roots in the early years. Birch and Fisher state that food exposure and accessibility, the modeling behavior of parents and siblings, and the level of parental control over food consumption influence a child's food preferences. Inappropriate feeding practices may result in under- or over-feeding and may promote negative associations with eating that continue into later life.

  Normal eating behaviors such as spitting out or gagging on unfamiliar food or food with texture are often misinterpreted as dislikes or intolerances leading to a diminished variety of foods offered. Infants have an innate preference for sweet and salty tastes. Without guidance, an infant may develop a lifelong preference for highly sweetened or salty foods rather than for a varied diet. (17)

  A young child gradually moves from the limited infant/toddler diet to daily multiple servings from each of the basic food groups as described in the Dietary Guidelines. (30) The toddler stage (ages 1-2 years) may frustrate caregivers since many toddlers have constantly changing food preferences and erratic appetites. In addition, toddlers become skeptical about new foods and may need to experience a food 15-20 times before accepting it. (31)

  Caregivers can be guided and supported in managing common toddler feeding problems. Feeding practices that caregivers can use to facilitate a successful transition to a food group-based diet include:
  - Offering a variety of developmentally appropriate nutritious foods
  - Reducing exposure to foods and beverages containing high levels of salt and sugar
  - Preparing meals that are pleasing to the eye and include a variety of colors and textures, setting a good example by eating a variety of foods;
  - Offering only whole milk from age 1-2 (lower fat milk can be introduced after that age)
  - Providing structure by scheduling regular meal and snack times
  - Allowing the child to decide how much or whether to eat
  - Allowing the child to develop eating/self-feeding skills
  - Eating with the child in a pleasant mealtime environment without coercion

References

1. American Academy of Pediatrics. Committee on Nutrition. Kleinman RE, editor. Pediatric Nutrition Handbook. 5th ed. 2004.
2. Pelto GH, Levitt E, and Thairu L. Improving feeding practices: Current patterns, common constraints, and the design of interventions. Food and Nutrition Bulletin, 2003; 24(1): 45-82.
3. United States Department of Agriculture. Study of WIC participant and program characteristics. 2000.
4. Institute of Medicine. Food and Nutrition Board. Proposed criteria for selecting the WIC food packages. The National Academies Press, Washington DC, 2004.

5. Hervada AR, Hervada-Page M. Infant Nutrition: The first two years. In: Childhood Nutrition. Lifshitz F, editor. CRC Press; 1995.
6. Pipes PL, Trahms CM. Nutrient needs of infants and children. In: PipesPL, Trahms CM editors. Nutrition in infancy and childhood 5th ed. Mosby Publishing Co. 1993.
7. Hendricks KM, Weaning: Pathophysiology, practice and policy. In: Nutrition in Pediatrics, 3rd edition. B.C. Decker Inc, 2003.
8. Institute of Medicine. Food and Nutrition Board. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. National Academies Press, Washington DC, 2001.
9. Clinical Nutrition Services; Warren Grant Magnuson Clinical Center, Office of Dietary Supplements. Facts about dietary supplements: zinc. National Institutes of Health. Bethesda Maryland; 2002.
10. Hallberg L. Perspectives on nutritional iron deficiency. Annu Rev Nutr. 2001; 21:1-21.
11. Centers for Disease Control and Prevention. Recommendations to prevent and control iron deficiency in the United States. MMWR. April 1998:18-21.
12. Looker AC, Dallman PR, Carroll MD, Gunter EW, Johnson CL. Prevalence of iron deficiency in the United States. JAMA. 1997; 277:973-6.
13. Centers for Disease Control and Prevention. Pediatric surveillance system 2003 annual report, Atlanta: U.S. Department of Health and Human Services, Center for Disease Control and Prevention, 2004. Available at http://www.cdc.gov/pednss (accessed 11/04).
14. Picciano MF, Smiciklas-Wright H, Birch LL, Mitchel DC, Murray-Kolb L, McConchy KL. Nutritional guidance is needed during dietary transition in early childhood. Ped. 2000; 106: 109-114.
15. Kahn JL, Binns HJ, Chen T, Tanz RR, Listernick R. Persistence and emergence of anemia in children during participation in the Special Supplemental Nutrition Program for Women, Infants, and Children. Arch Pediatr Adolesc Med. 2002; 156:1028-32.
16. Proskitt EM. Early feeding and obesity. In: Boulton J, Laron Z and Rey J, editors. Long-term consequences of early feeding. Nestle Nutrition Workshops Series; 1996, Nestle Ltd., Vevey/Lippincott-Raven Publishers, Philadelphia; Vol. 36.
17. Birch LL, Fisher JO. Development of eating behaviors among children and adolescents. Ped. 1998; 101:539-549.
18. Bertness J, Holt K, editors. Promoting awareness, preventing pain: Facts on early childhood caries (ECC) 2nd. Ed. [Fact Sheet on the Internet]. Washington (DC); National Maternal & Child Oral Health Resource Center; 2004. Available from: http://www.mchoralhealth.org.
19. American Academy of Pediatric Dentistry. Baby bottle tooth decay/early childhood caries. Pediatr Dent. 2001 Mar-Apr; 23 (2): 18.
20. al-Shalan TA, Erickson PR, Hardie NA. Primary incisor decay before age 4 as a risk factor for future dental caries. J Pediatr Dent. 1997 Jan-Feb; 9 (1): 37-41.
21. Casamassimo P ed. 1996. Bright futures in practice: oral health. Arlington, VA: National Center for Education in Maternal and Child Health.
22. Northstone K, Rogers I, Emmett P. Drinks consumed by 18-month-old children: Are current recommendations being followed? Eur J Clin Nutr. 2002; 56:236-44.
23. American Academy of Pediatric Dentistry. Policy on early childhood caries (EEC): Classifications, consequences, and prevention strategies. Pediatr Dent; Reference manual 2003-2004: 2004; 25(7):25

24. Butte N, Cobb K, Dwyer J, Graney L, Heird W, Rickard K. The start healthy feeding guidelines for infants and toddlers. J Am Diet Assoc, 2004; 104 (3) 442-454.
25. Northstone, K, Emmett P, Nethersole F. The effect of age of introduction to lumpy solids on foods eaten and reported difficulties at 6 and 15 months. J Hum Nutr Dietet. 2001; 14: 43-54.
26. Baydar N, McCann M, Williams R, Vesper E, McKinney P. WIC infant feeding practices study. USDA Office of Analysis and Evaluation. November 1997.
27. Ponza M, Devaney B, Ziegler P, Reidy K, and Squatrito C. Nutrient intake and food choices of infants and toddlers participating in WIC. J Am Diet Assoc 2004; 104: s71-s79.
28. Harris CS, Baker SP, Smith GA, Harris RM. Childhood asphyxiation by food: A national analysis and overview. JAMA. 1984; 251:2231-5.
29. Lucas B. Normal nutrition from infancy through adolescence. In: Handbook of pediatric nutrition. 2nd ed. Gaithersburg, Maryland: Aspen Publishers, Inc. 1999.
30. United States Department of Agriculture and the United States Department of Health and Human Services. Dietary guidelines for Americans, 5th ed. 2000. Available from:http://www.usda.gov.cnpp.
31. Story M, Holt K, Sofka D, editors. Bright futures in practice: nutrition. 2nd ed. Arlington, VA: National Center for Education in Maternal and Child Health; 2002.

Nutrition Risk Criteria **501 POSSIBILITY OF REGRESSION**

Category, Priority and Referral

Category	Priority	Referral
BREASTFEEDING WOMEN	1	
NON-BREASTFEEDING WOMEN	6	
INFANTS	1	
CHILDREN	3	

Definition

A participant who has previously been certified eligible for the Program may be considered to be at nutritional risk in the next certification period if the competent professional authority determines there is a possibility of regression in nutritional status without the benefits that the WIC Program provides.

***Possibility of Regression may not be used for consecutive certifications.**

***Use when no other risk can be identified. This code is the only code the participant has when used.**

Required Documentation

Document what previous risk code qualifies the applicant for regression risk use.

Justification

On occasion, a participant's nutritional status may be improved, to the point that s/he rises slightly above the cutoff of the initial risk condition by the end of the certification period. This occurs most frequently with those conditions that contain specific cutoffs or thresholds, such as anemia or inappropriate growth. Removal of such individuals from the Program can result in a "revolving-door" situation where the individual's recently improved nutritional status deteriorates quickly, so that s/he then re-enters the Program at equal or greater nutrition risk status than before. Therefore, WIC Program regulations permit State agencies to certify previously certified individuals who do not demonstrate a current nutrition risk condition against the possibility of their reverting to the prior existing risk condition if they do not continue to receive WIC benefits. This policy is consistent with the preventive nature of the WIC Program, and enables State and local agencies to ensure that their previous efforts to improve a participant's nutrition status, as well as to provide referrals to other health care, social service, and/or public assistance programs are not wasted.

Competent Professional Authorities and other certifying staff should keep in mind that every nutrition risk condition does not necessarily lead itself to the possibility of regression. For example, gestational diabetes or gingivitis of pregnancy are not conditions to which a new mother could regress, since they are directly associated with pregnancy, and the breastfeeding or non-breastfeeding women cannot regress to being pregnant if she is no longer receiving WIC benefits.

Clarification

After April 1, 1999, any certification for regression must be based on the new set of risk criteria. For example, a person deemed anemic under a State’s more inclusive criteria prior to April 1, 1999, may only be certified for regression after April 1, 1999, if his/her blood values would have met the revised CDC criteria for anemia published in the April 1998 MMWR tables.

Further, regression may only be used as a certifying nutrition risk when it complies with the policies established by the State agency for its use, as set forth in the WIC Nutrition Services Standards issued by FNS in 1988. Such policies must include:

1. A requirement for a nutritional assessment to rule out the existence of another current risk factor before using eligibility on regression.
2. A requirement for written identification of the risk factor to which the participant may regress.
3. A list of risk factors and priority levels for which eligibility based regression may be applied; and
4. A limit on the number of times regression for a given risk factor may be consecutively applied.

References

WIC Program Regulations, Sect. 246.7(e)(1)(iii).

Nutrition Risk Criteria **502★ TRANSFER OF CERTIFICATION**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	as needed
BREASTFEEDING WOMEN	4	“
NON-BREASTFEEDING WOMEN	6	“
INFANTS	4	“
CHILDREN	5	“

Definition

Person with current valid Verification of Certification (VOC) document from another State or local agency. The VOC is valid until the certification period expires, and shall be accepted as proof of eligibility for program benefits. If the receiving local agency has waiting lists for participation, the transferring participant shall be placed on the list ahead of all other waiting applicants.

This criterion would be used primarily when the VOC card/document does not reflect another (more specific) nutrition risk condition at the time of transfer or if the participant was initially certified based on a nutrition risk condition not in use by the receiving State agency.

Required Documentation

VOC document from the certifying WIC clinic*

*See policy related to VOC transfers for more information

Justification

Local agencies must accept Verification of Certification (VOC) documents from participants. A person with a valid VOC document shall not be denied participation in the receiving State because the person does not meet that State’s particular eligibility criteria. Once a WIC participant has been certified by a local agency, the service delivery area into which s/he moves is obligated to honor that commitment.

References

1. FNS Instruction 803-11, Rev.1.
2. WIC Program Regulations; Section 246.7(k).

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1,2, or 4*	-
BREASTFEEDING WOMEN	1, 2, or 4*	-

*Must be the same priority as at-risk infant.

Definition

A breastfeeding woman whose breastfed infant has been determined to be at nutritional risk.

Required Documentation

The infant's nutrition risk must be documented on the mother's certification record.

Justification

A breastfed infant is dependent on the mother's milk as the primary source of nutrition. Special attention should therefore be given to the health and nutritional status of the mother (5). Lactation requires an additional approximately 500 Kcal per day as increased protein, calcium, and other vitamins and minerals (3,1). Inadequate maternal nutrition may result in decreased nutrient content of the milk (1).

References

1. Institute of Medicine. Nutrition During Lactation. National Academy Press, Washington, D.C.; 1991.
2. Lawrence RA. Breastfeeding a guide for the medical profession. St. Louis: Mosby, 1994.
3. National Research Council (U.S.), Subcommittee on the Tenth Edition of the RDAs, National Institutes of Health, Committee on Dietary Allowances. Recommended dietary allowances. Washington, D.C.: National Academy Press, 1989.
4. WIC Program Regulations, Sect. 246.7(e)(1)(iii).
5. Worthington-Roberts BS, Williams SR. Nutrition in Pregnancy and Lactation. St. Louis: Mosby, 1993.

Nutrition Risk Criteria **602 BREASTFEEDING COMPLICATIONS OR POTENTIAL COMPLICATIONS (WOMEN)**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	**1**	**RD/LC**
BREASTFEEDING WOMEN	**1**	**RD/LC**

Definition

A breastfeeding woman with any of the following complications or potential complications for breastfeeding:

- Severe breast engorgement
- Recurrent plugged ducts
- Mastitis (fever or flu-like symptoms with localized breast tenderness)
- Flat or inverted nipples
- Cracked, bleeding or severely sore nipples
- Age ≥ 40 years
- Failure of milk to come in by 4 days postpartum
- Tandem nursing (breastfeeding two siblings who are not twins)

Required Documentation

Document breastfeeding complication(s)

Justification

- **Severe breast engorgement**
  Severe engorgement is often caused by infrequent nursing and/or ineffective removal of milk. This severe breast congestion causes the nipple-areola area to become flattened and tense, making it difficult for the baby to latch-on correctly. The result can be sore, damaged nipples and poor milk transfer during feeding attempts. This ultimately results in diminished milk supply. When the infant is unable to latch-on or nurse effectively, alternative methods of milk expression are necessary, such as using an electric breast pump.
- **Recurrent plugged ducts**
  A clogged duct is a temporary back-up of milk that occurs when one or more of the lobes of the breast do not drain well. This usually results from incomplete emptying of milk. Counseling on feeding frequency or method or advising against wearing an overly tight bra or clothing can assist.
- **Mastitis**
  Mastitis is a breast infection that causes a flu-like illness accompanied by an inflamed, painful area of the breast - putting both the health of the mother and successful breastfeeding at risk. The woman should be referred to her health care provider for antibiotic therapy.
- **Flat or inverted nipples**
  Infants may have difficulty latching-on correctly to nurse when nipples are flat or inverted. Appropriate interventions can improve nipple protractility and skilled help guiding a baby in proper breastfeeding technique can facilitate proper attachment.
- **Cracked, bleeding or severely sore nipples**
  Severe nipple pain, discomfort lasting throughout feedings, or pain persisting beyond one week postpartum is atypical and suggests the baby is not positioned correctly at the breast. Improper infant latch-on not only causes sore nipples, but impairs milk flow and leads to diminished milk supply and

inadequate infant intake. There are several other causes of severe or persistent nipple pain, including Candida or staph infection. Referrals for lactation counseling and/or examination by the woman's health care provider are indicated.

- **Age ≥ 40 years**
  Older women (over 40) are more likely to experience fertility problems and perinatal risk factors that could impact the initiation of breastfeeding. Because involutional breast changes can begin in the late 30s, older mothers may have fewer functioning milk glands resulting in greater difficulty producing an abundant milk supply.
- **Failure of milk to come in by 4 days postpartum**
  Failure of milk to come in by 4 days postpartum may be a result of maternal illness or perinatal complications. This may place the infant at nutritional and/or medical risk, making temporary supplementation necessary until a normal breast milk supply is established.
- **Tanden nursing (breastfeeding two siblings who are not twins)**
  With tandem nursing the older baby may compete for nursing privileges, and care must be taken to assure that the younger baby has first access to the milk supply. The mother who chooses to tandem nurse will have increased nutritional requirements to assure her adequate milk production.

References

1. Alexander JM, Grant AM, Campbell MJ. Randomised controlled trial of breast shells and Hoffman's exercises for inverted and non-protractile nipples. BMJ 1992;304:1030-2.
2. Akre J. Infant Feeding. The physiological basis. Bull.World Health Organ 1989;67 Suppl:1-108.
3. Amier, L, Garland, SM, Dennerstein, L, et al.: Candida albicans: Is it associated with nipple pain in lactating women? Gynecol Obstetr Invest; 1996; 41:30-34.
4. De Coopman J. Breastfeeding after pituitary resection: support for a theory of autocrine control of milk supply? J.Hum.Lact. 1993;9:35-40.
5. Lawrence RA. Breastfeeding a guide for the medical profession. St. Louis: Mosby, 1994.
6. Livingstone VH, Willis CE, Berkowitz J. Staphylococcus aureus and sore nipples. Can.Fam.Physician 1996;42:654-9.
7. Mohrbacher N, Stock J, La LL, I. The breastfeeding answer book. Schaumburg, Ill: La Leche League International, 1997.
8. Neifert M. Early assessment of the breastfeeding infant. Contemporary Pediatr. 1996 Oct;13:142.
9. Neifert MR. The optimization of breast-feeding in the perinatal period. Clin.Perinatol. 1998;25:303-26.
10. Neifert MR, Seacat JM, Jobe WE. Lactation failure due to insufficient glandular development of the breast. Pediatrics 1985;76:823-8.
11. Riodan J, Auerbach KG. Breastfeeding and human lactation. Boston: Jones and Bartlett Publishers, 1993.
12. The Main Trial Collaborative Group: Preparing for breastfeeding: treatment of inverted and non-protractile nipples in pregnancy; Midwifery; 1994; 10:200.
13. Woolridge MW. Aetiology of sore nipples. Midwifery 1986; 2:172-6.

Nutrition Risk Criteria	603	BREASTFEEDING COMPLICATIONS/POTENTIAL COMPLICATIONS (INFANTS)

Category, Priority and Referral

Category	Priority	Referral
INFANTS	1	RD/LC

Definition

A breastfed infant with any of the following complications or potential complications for breastfeeding:

- jaundice
- weak or ineffective suck
- difficulty latching onto mother's breast
- inadequate stooling (for age, as determined by a physician or other health care professional), and/or less than 6 wet diapers per day

Required Documentation

Document breastfeeding complication(s)

Justification

- Jaundice occurs when bilirubin accumulates in the blood because red blood cells break down too quickly, the liver does not process bilirubin as efficiently as it should, or intestinal excretion of bilirubin is impaired. The slight degree of jaundice observed in many healthy newborns is considered physiologic. Jaundice is considered pathologic if it appears before 24 hours, lasts longer than a week or two, reaches an abnormally high level, or results from a medical problem such as rapid destruction of red blood cells, excessive bruising, liver disease, or other illness. When jaundice occurs in an otherwise healthy breastfed infant, it is important to distinguish "breastmilk jaundice" from "breastfeeding jaundice" and determine the appropriate treatment.

  In the condition known as "breastmilk jaundice," the onset of jaundice usually begins well after the infant has left the hospital, 5 to 10 days after birth, and can persist for weeks and even months. Early visits to the WIC clinic can help identify and refer these infants to their primary health care provider. Breastmilk jaundice is a normal physiologic phenomenon in the thriving breastfed baby and is due to a human milk factor that increases intestinal absorption of bilirubin. The stooling and voiding pattern is normal. If the bilirubin level approaches 18-20 mg%, the health care provider may choose to briefly interrupt breastfeeding for 24-36 hours which results in a dramatic decline in bilirubin level.

  - Resumption of breastfeeding usually results in cessation of the rapid fall in serum bilirubin concentration, and in many cases a small increase may be observed, followed by the usual gradual decline to normal.
  - "Breastfeeding jaundice", is an exaggeration of physiologic jaundice, which usually peaks between 3 and 5 days of life, though it can persist longer. This type of jaundice is a common marker for inadequate breastfeeding. An infant with breastfeeding jaundice is underfed and displays the following symptoms: infrequent or ineffective breastfeeding; failure to gain appropriate weight; infrequent stooling with delayed

appearance of yellow stools (i.e., prolonged passage of meconium); and scant dark urine with urate crystals. Improved nutrition usually results in a rapid decline in serum bilirubin concentration.

- A weak or ineffective suck may cause a baby to obtain inadequate milk with breastfeeding and result in a diminished milk supply and an underweight baby. Weak or ineffective suckling can be due to prematurity, low birth weight, a sleepy baby, or physical/medical problems such as heart disease, respiratory illness, or infection. Newborns who receive bottle feedings before beginning breastfeeding or who frequently use a pacifier may have trouble learning the proper tongue and jaw motions required for effective breastfeeding.

  Difficulty latching onto the mother's breast may be due to flat or inverted nipples, breast engorgement, or incorrect positioning and breastfeeding technique. Early exposure to bottle feedings can predispose infants to "nipple confusion" or difficulty learning to attach to the breast correctly and effectively extract milk. A referral for lactation counseling should be made.

- Inadequate stooling or less than 6 wet diapers are probable indicators that the breastfed infant is not receiving adequate milk. Not only is the baby at risk for failure to thrive, but the mother's milk is at risk for rapidly diminishing due to ineffective removal of milk. The breastfed infant with inadequate caloric intake must be identified early and the situation remedied promptly to avoid long-term consequences of dehydration or nutritional deprivation. Although failure to thrive can have many etiologies, the most common cause in the breastfed infant is insufficient milk intake as a result of infrequent or ineffective nursing. Inadequate breastfeeding can be due to infant difficulties with latching on or sustaining suckling, use of a nipple shield over the mother's nipple, impaired let down of milk, a non-demanding infant, excessive use of a pacifier, or numerous other breastfeeding problems.

  The literature regarding inadequate stooling varies widely in terms of quantification; this condition is best diagnosed by the pediatrician or other health care practitioner.

References

1. Auerbach KG, Gartner LM. Breastfeeding and human milk: their association with jaundice in the neonate. Clin.Perinatol. 1987;14:89-107.
2. Barros FC, Victora CG, Semer TC, Tonioli FS, Tomasi E, Weiderpass E. Use of pacifiers is associated with decreased breast-feeding duration. Pediatrics 1995;95:497-9.
3. Bocar DL. The lactation consultant: part of the health care team. NAACOGS.Clin.Issu.Perinat.Womens Health Nurs. 1992;3:731-7.
4. Cooper WO, Atherton HD, Kahana M, Kotagal UR. Increased incidence of severe breastfeeding malnutrition and hypernatremia in a metropolitan area. Pediatrics 1995;96:957-60.Neifert MR. The optimization of breast-feeding in the perinatal period. Clin.Perinatol. 1998;25:303-26.
5. De Carvalho M, Robertson S, Friedman A, Klaus M. Effect of frequent breast-feeding on early milk production and infant weight gain. Pediatrics 1983;72:307-11.
6. Kurinij N, Shiono PH. Early formula supplementation of breast-feeding. Pediatrics 1991;88:745-50.
7. Lawrence RA. Breastfeeding a guide for the medical profession. St. Louis: Mosby, 1994.

8. Maisels MJ, Newman TB. Kernicterus in otherwise healthy, breast-fed term newborns. Pediatrics 1995;96:730-3.
9. Meier PP, Engstrom JL, Fleming BA, Streeter PL, Lawrence PB. Estimating milk intake of hospitalized preterm infants who breastfeed. J.Hum.Lact. 1996;12:21-6.

Nutrition Risk Criteria **701** **BORN TO WIC MOM OR A POTENTIAL WIC MOM**

Category, Priority and Referral	**Category**	**Priority**	**Referral**
	INFANTS	2	-

Definition

An infant under six months age whose mother was a WIC participant during pregnancy or whose mother's medical records document that the woman was at nutritional risk during pregnancy because of detrimental or abnormal nutritional conditions detectable by biochemical or anthropometric measurements or other documented nutritionally related medical conditions.

Required Documentation

Justification

Federal regulations designate these conditions for WIC eligibility (3).

WIC participation during pregnancy is associated with improved pregnancy outcomes. An infant whose nutritional status has been adequately maintained through WIC services during gestation and early infancy may decline in nutritional status if without these services and return to a state of elevated risk for nutrition related health problems. Infants whose mother was at medical/nutritional risk during pregnancy, but did not receive those services, may also be thought of as a group at elevated risk for morbidity and mortality in the infant period (1, 2).

WIC participation in infancy is associated with lower infant mortality, decreased anemia for infants and improvements in growth (head circumference, height and weight). Infants on WIC are more likely to consume iron-fortified formula and cereal and less likely to consume cow's milk before one year, thus lowering the risk of developing iron deficiency anemia (1, 2).

References

1. Disbrow DD. The costs and benefits of nutrition services: a literature review. J.Am.Diet.Assoc. 1989;89:S3-66.
2. Ryan AS, Martinez GA, Malec DJ. The effect of the WIC program on nutrient intakes of infants, 1984. Med.Anthropol. 1985;9153-72.
3. WIC Program Regulations; Section 246.7(e)(1)(ii).

	Category	Priority	Referral
Category, Priority and Referral	INFANTS	1, 2 or 4* *Must be the same priority as at-risk mother.	-

Definition

Breastfeeding infant of a mother who is at nutritional risk.

Required Documentation

The mother's nutrition risk must be documented on the infant's certification record.

Justification

A breastfed infant is dependent on the mother's milk as the primary source of nutrition. Lactation requires the mother to consume an additional 500 Kcal per day (approximately) as well as increased protein, calcium, and other vitamins and minerals (2, 1). Inadequate maternal nutrition may result in decreased nutrient content of the milk (1). Special attention should therefore be given to the health and nutritional status of breastfed infants whose mothers are at nutritional risk (4).

References

1. Institute of Medicine. Nutrition During Lactation. National Academy Press, Washington, D.C.; 1991.
2. National Research Council (U.S.), Subcommittee on the Tenth Edition of the RDAs, National Institutes of Health, Committee on Dietary Allowances. Recommended dietary allowances. Washington, D.C.: National Academy Press, 1989.
3. WIC Program Regulations; Section 246.7(e)(1)(i).
4. Worthington-Roberts BS, Williams SR. Nutrition During Pregnancy and Lactation. St. Louis: Mosby, 1989.

Nutrition Risk Criteria **801★ HOMELESSNESS**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	-
BREASTFEEDING WOMEN	4	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	4	-
CHILDREN	5	-

Definition

A woman, infant or child who lacks a fixed and regular nighttime residence; or whose primary nighttime residence is:

- A supervised publicly or privately operated shelter (including a welfare hotel, a congregate shelter, or a shelter for victims of domestic violence) designed to provide temporary living accommodations
- An institution that provides a temporary residence for individuals intended to be institutionalized
- A temporary accommodation of not more than 365 days in the residence of another individual
- A public or private place not designed for, or ordinarily used as, a regular sleeping accommodation for human beings

Required Documentation

Justification

Homeless individuals comprise a very vulnerable population with many special needs. WIC program regulations specify homelessness as a predisposing nutrition risk condition. Today's homeless population contains a sizeable number of women and children – over one-third of the total homeless population in the U.S. Studies show 43% of today's homeless are families, and an increasing number of the "new homeless" include economically displaced individuals who have lost their jobs, exhausted their resources, have recently entered into the ranks of the homeless and consider their condition to be temporary.

Reference

WIC Program Regulations; Sect. 246.7(e)(2)(iv).

Nutrition Risk Criteria **802★ MIGRANCY**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	-
BREASTFEEDING WOMEN	4	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	4	-
CHILDREN	5	-

Definition

Categorically eligible women, infants and children who are members of families which contain at least one individual whose principal employment is in agriculture on a seasonal basis, who has been so employed within the last 24 months, and who establishes, for the purposes of such employment, a temporary abode.

Required Documentation

Justification

Data on the health and/or nutritional status of migrants indicate significantly higher rates or incidence of infant mortality, malnutrition, and parasitic disease (among migrant children) than among the general U.S. population. Therefore, migrancy has long been stipulated as a condition that predisposes persons to inadequate nutritional patterns or nutritionally related medical conditions.

Reference

WIC Program Regulations; Sect. 246.7(e)(2)(iv).

Nutrition Risk Criteria **902** **WOMAN OR INFANT/CHILD OF PRIMARY CAREGIVER WITH LIMITED ABILITY TO MAKE FEEDING DECISIONS/PREPARE FOOD**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	4	-
BREASTFEEDING WOMEN	4	-
NON-BREASTFEEDING WOMEN	6	-
INFANTS	4	-
CHILDREN	5	-

Definition

Woman (pregnant, breastfeeding, or non-breastfeeding), or infant/child whose primary caregiver is assessed to have a limited ability to make appropriate feeding decisions and/or prepare food. Examples may include individuals who are:

- ≤ 17 years of age;
- mentally disabled/delayed and/or have a mental illness such as clinical depression (diagnosed by a physician or licensed psychologist);
- physically disabled to a degree which restricts or limits food preparation abilities; or
- currently using or having a history of abusing alcohol or other drugs.

Required Documentation

Justification

The mother or caregiver ≤ 17 years of age generally has limited exposure and application of skills necessary to care for and feed a total dependent. Cognitive limitation in a parent or primary caregiver has been recognized as a risk factor for failure to thrive, as well as for abuse and neglect. The mentally handicapped caregiver may not exhibit the necessary parenting skills to promote beneficial feeding interactions with the infant. Maternal mental illnesses such as severe depression and maternal chemical dependency are also strongly associated with abuse and neglect. In 22 states, 90% of caregivers reported for child abuse are active substance abusers. Certain physical handicaps such as blindness, para- or quadriplegia, or physical anomalies restrict/limit the caregiver's ability to prepare and offer a variety of foods. Education, referrals and service coordination with WIC will aid the mother/caregiver in developing skills, knowledge and/or assistance to properly care for a total dependent.

References

1. Accardo PJ, Whitman BY. Children of mentally retarded parents. Am.J.Dis.Child 1990;144:69-70.
2. Grand RJ, Sutphen JL, Dietz WH. Pediatric nutrition theory and practice. Boston: Butterworths, 1987.
3. Institute of Medicine. WIC nutrition risk criteria a scientific assessment. National Academy Press, Washington, D.C.; 1996.
4. Pollitt E, Wirtz S. Mother-infant feeding interaction and weight gain in the first month of life. J.Am.Diet.Assoc. 1981;78:596-601.
5. WIC Program regulations; Sect. 246.7(e)(2).

Nutrition Risk Criteria **903★ FOSTER CARE**

Category, Priority and Referral

Category	Priority	Referral
INFANTS	4	-
CHILDREN	5	-

Definition

Entering the foster care system during the previous six months or moving from one foster care home to another foster care home during the previous six months.

Required Documentation

Justification

"Foster children are among the most vulnerable individuals in the welfare system. As a group, they are sicker than homeless children and children living in the poorest sections of inner cities." This statement from a 1995 Government Accounting Office report on the health status of foster children confirms research findings that foster children have a high frequency of mental and physical problems, often the result of abuse and neglect suffered prior to entry into the foster care system. When compared to other Medicaid-eligible children, foster care children have higher rates of chronic conditions such as asthma, diabetes and seizure disorders. They are also more likely than children in the general population to have birth defects, inadequate nutrition and growth retardation including short stature.

Studies focusing on the health of foster children often point out the inadequacy of the foster care system in evaluating the health status and providing follow-up care for the children for whom the system is responsible. Because foster care children are wards of a system which lacks a comprehensive health component, the social and medical histories of foster children in transition, either entering the system or moving from one foster care home to another, are frequently unknown to the adults applying for WIC benefits for the children. For example, the adult accompanying a foster child to a WIC clinic for a first-time certification may have no knowledge of the child's eating patterns, special dietary needs, chronic illnesses or other factors which would qualify the child for WIC. Without any anthropometric history, failure to grow, often a problem for foster children, may not be diagnosed even by a single low cutoff percentile.

Since a high proportion of foster care children have suffered from neglect, abuse or abandonment and the health problems associated with these, entry into foster care or moving from one foster care home to another during the previous six months is a nutritional risk for certification in the WIC Program. Certifiers using this risk should be diligent in evaluating and documenting the health and nutritional status of the foster child to identify other risks as well as problems that may require follow-up or referral to other health care programs. This nutritional risk cannot be used for consecutive certifications while the child remains in the same foster home. It should be used as the sole risk criterion only if careful assessment of the applicant's nutritional status indicates that no other risks based on anthropometric, medical or nutritional risk criteria can be identified.

The nutrition education, referrals and service coordination provided by WIC will support the foster parent in developing the skills and knowledge to ensure that the foster child receives appropriate nutrition and health care. Since a foster parent frequently has inadequate information about a new foster child's health needs, the WIC nutritionist can alert the foster parent to the nutritional risks that many foster care children have and suggest ways to improve the child's nutritional status.

References

1. American Medical News: America's Sickest Children; January 10, 1994; 15-19.
2. Chernoff R, Combs-Orme T, Risley-Curtiss C, Heisler A. Assessing the health status of children entering foster care. Pediatrics 1994;93:594-601.
3. DuRouseau PC, Moquette-Magee E, Disbrow D. Children in foster care: are they at nutritional risk? J.Am.Diet.Assoc. 1991 Jan;91(1):83-85.
4. Government Accounting Office. Foster care health needs of many young children are unknown and unmet: report to the ranking minority member, Subcommittee on Human Resources, Committee on Ways and Means, House of Representatives. Washington D.C.: The Office; 1995 May. Report No.:GA 1.13:HEHS-95-114.
5. Halfon N, Mendonca A, Berkowitz G. Health status of children in foster care. The experience of the Center for the Vulnerable Child. Arch.Pediatr.Adolesc.Med. 1995;149:386-92.
6. Schor EL. The foster care system and health status of foster children. Pediatrics 1982;69:521-8.
7. Takayama JI, Wolfe E, Coulter KP. Relationship between reason for placement and medical findings among children in foster care. Pediatrics 1998;101:201-7.
8. Wyatt DT, Simms MD, Horwitz SM. Widespread growth retardation and variable growth recovery in foster children in the first year after initial placement. Arch.Pediatr.Adolesc.Med. 1997;151:813-6.

Nutrition Risk Criteria **904 EXPOSURE TO ENVIRONMENTAL TOBACCO SMOKE (ETS)**

Category, Priority and Referral

Category	Priority	Referral
PREGNANT WOMEN	1	-
BREASTFEEDING WOMEN	1	-
INFANTS	1	-
CHILDREN	3	-
NON-BREASTFEEDING WOMEN	6	-

Definition

Environmental tobacco smoke (ETS) exposure is defined (for WIC eligibility purposes) as exposure to smoke from tobacco products inside the home. *(1,2,3)
* See Clarification for background information.

Required Documentation

Document tobacco smoke exposure

Justification

ETS is a mixture of the smoke given off by a burning cigarette, pipe, or cigar (side stream smoke), and the smoke exhaled by smokers (mainstream smoke). ETS is a mixture of about 85% side stream and 15% mainstream smoke (4) made up of over 4,000 chemicals, including Polycyclic Aromatic Hydrocarbons (PAHs) and carbon monoxide (5). Side stream smoke has a different chemical make-up than main-stream smoke. Side stream smoke contains higher levels of virtually all carcinogens, compared to mainstream smoke (6). Mainstream smoke has been more extensively researched than side stream smoke, but they are both produced by the same fundamental processes.

ETS is qualitatively similar to mainstream smoke inhaled by the smoker. The 1986 Surgeon General's report: *The Health Consequences of Involuntary Smoking. A Report of the Surgeon General* concluded that ETS has a toxic and carcinogenic potential similar to that of the mainstream smoke (7). The more recent 2006 Surgeon General's report, *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General,* reaffirms and strengthens the findings of the 1986 report, and expands the list of diseases and adverse health effects caused by ETS (8).

ETS is a known human carcinogen (2). Women who are exposed to ETS are at risk for lung cancer and cardiovascular diseases (9). Prenatal or postnatal ETS exposure is related to numerous adverse health outcomes among infants and children, including sudden infant death syndrome (SIDS) (10, 11), upper respiratory infections (12), periodontal disease (13), increased severity of asthma/wheezing (12), metabolic syndrome (14), decreased cognitive function (15), lower birth weight and smaller head circumference (16). Infants born to women exposed to ETS during pregnancy have a small decrease in birth weight and a slightly increased risk of intrauterine growth retardation compared to infants of unexposed women (17).

Studies suggest that the health effects of ETS exposure at a young age could last into adulthood. These include cancer (18), specifically lung cancer (19, 20), and cardiovascular diseases (14, 21, 22,). There is strong evidence that ETS

exposure to the fetus and/or infant results in permanent lung damage (23, 24, 25, 26).

ETS exposure increases inflammation and oxidative stress (27, 28, 29). Inflammation is associated with asthma (30), cardiovascular diseases (31, 32), cancer (33), chronic obstructive pulmonary disease (34), and metabolic syndrome (14, 35). PAHs are the major class of compounds that contribute to the ETS-related adverse health outcomes. These compounds possess potent carcinogenic and immunotoxic properties that aggravate inflammation.

Oxidative stress is a general term used to describe the steady state of oxidative damage caused by highly reactive molecules known as free radicals. The free radicals can be generated both during the normal metabolic process and from ETS and other environmental pollutants. When free radicals are not neutralized by antioxidants, they can cause oxidative damage to the cells. This damage has been implicated in the cause of certain diseases. ETS provokes oxidant damage similar to that of active smoking (36).

Antioxidants may modulate oxidative stress-induced lung damage among both smokers and non-smokers (22, 27-29, 37-40). Fruits and vegetables are the major food sources of antioxidants that may protect the lung from oxidative stress (1). Research indicates that consuming fruits and vegetables is more beneficial than taking antioxidant supplements (1). This suggests that other components of fruits and vegetables may be more relevant in protecting the lung from oxidative stress. Dietary fiber is also thought to contribute to the beneficial health effects of fruits and vegetables (1).

The Institute of Medicine (IOM) reports that an increased turnover in vitamin C has been observed in nonsmokers who are regularly exposed to tobacco smoke (41). The increased turnover results in lowered vitamin C pools in the body. Although there are insufficient data to estimate a special requirement for non-smokers regularly exposed to ETS, the IOM urges those individuals to ensure that they meet the Recommended Dietary Allowance for vitamin C (36, 41).

The WIC food package supplements the participant intake of vitamin C. In addition, many WIC State Agencies participate in the WIC Farmers' Market Nutrition Program, which provides coupons for participants to purchase fresh fruits and vegetables. WIC Program benefits also include counseling to increase fruit and vegetable consumption, and to promote a healthy lifestyle, such as protecting participants and their children from ETS exposure. WIC staff may also make appropriate referrals to participants, and/or their caregivers, to other health and social services, such as smoking cessation programs.

In a comprehensive scientific report, the Surgeon General concluded that there is no risk-free level of exposure to secondhand smoke (8). However, for the purpose of risk identification, the definition used for this risk criterion is based on the Centers for Disease Control and Prevention (CDC) Pediatric Nutrition Surveillance System (PedNSS) and the Pregnancy Nutrition Surveillance System (PNSS) questions to determine Environmental Tobacco Smoke (ETS) exposure:

1. Does anyone living in your household smoke inside the home? (infants, children)
2. Does anyone else living in your household smoke inside the home? (women)

Clarification

Because the definition used by other Federal agencies for ETS exposure is specific to “inside the home” and has been validated (3), the definition used for WIC eligibility must also be as specific. In addition, FNS encourages the use of the PedNSS and PNSS ETS exposure questions for WIC nutrition assessment.

There are other potential sources of ETS exposure, such as work and day care environments. However, no other validated questions/definitions could be found that were inclusive of other environments and applicable to WIC.

References

1. Lesley Butler, et al. RISC/WIC Report on Environmental Tobacco Smoke Exposure. February 2006. Unpublished.
2. Respiratory Health Effects of Passive Smoking (Also Known as Exposure to Secondhand Smoke or Environmental Tobacco Smoke ETS). U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Washington, DC, EPA/600/6-90/006F, 1992. http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=2835 Accessed March 2006.
3. Pirkle, JL, KM Flegal, DJ Brody, RA Etzel, and KR Maurer. Exposure of the U.S. Population to Environmental Tobacco Smoke. The Third National Health and Nutrition Examination Survey, 1988 to 1991. *JAMA;* 1996; 275,16; 1233-1240.
4. Witschi, H, JP Joad, and KE Pinkerton. The toxicology of environmental tobacco smoke. *Annu Rev Pharmacol Toxicol* 1997; 37: 29-52.
5. Seifert, JA, CA Ross, and JM Norris. Validation of a five-question survey to assess a child’s exposure to environmental tobacco smoke. *Ann Epidemiol* 2002; 12:273-277.
6. Adams, JD, KJ O’Mara-Adams, and D Hoffmann. Toxic and carcinogenic agents in undiluted main-stream smoke and sidestream smoke of different types of cigarettes. *Carcinogenesis* 1987-8:729-731.
7. U.S. Department of Health and Human Services. *The Health Consequences of Involuntary Smoking: A Report of the Surgeon General.* Rockville (MD): U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Health Promotion and Education, Office on Smoking and Health, 1986. DHHS Publication No. (CDC) 87-8398.
8. U.S. Department of Health and Human Services*. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General—Executive Summary.* U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006.
9. National Cancer Institute. *Health Effects of Exposure to Environment Tobacco Smoke.* Smoking and Tobacco Control Monograph No. 10 (PDF - 71k). Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute; 1999. NIH Pub. No. 99-4645. Accessed: March 2006.

10/1/2016