



**Idaho STD and HIV Infection by  
Public Health District of Residence  
April - June 2017**

District	HIV Infection¶	Chlamydia	Gonorrhea	Early Syphilis**	Congenital Syphilis	Late or Unknown Duration Syphilis†	TOTAL
1	3	207	20	1	0	1	232
2	0	67	10	2	0	0	79
3	2	245	29	1	0	0	277
4	1	518	61	6	0	1	587
5	2	189	10	0	0	0	201
6	1	104	31	0	0	0	136
7	0	113	20	1	0	0	134
<b>Total</b>	<b>9</b>	<b>1,443</b>	<b>181</b>	<b>11</b>	<b>0</b>	<b>2</b>	<b>1,646</b>

**Idaho STD and HIV Infection by  
Public Health District of Residence  
April - June 2016**

District	HIV Infection¶	Chlamydia	Gonorrhea	Early Syphilis**	Congenital Syphilis	Late or Unknown Duration Syphilis†	TOTAL
1	0	125	12	4	0	4	145
2	0	76	0	0	0	0	76
3	1	231	4	2	0	0	238
4	4	404	91	5	0	8	512
5	1	129	17	2	0	1	150
6	1	138	12	3	0	2	156
7	0	103	10	0	0	0	113
<b>Total</b>	<b>7</b>	<b>1,206</b>	<b>146</b>	<b>16</b>	<b>0</b>	<b>15</b>	<b>1,390</b>

\*Data are provisional. Percentages may not sum to 100% due to rounding.


\*\*Syphilis of ≤ 1 year duration: primary, secondary, and early latent syphilis

†Syphilis of > 1 year or unknown duration: late symptomatic, late latent, and unknown duration

¶Newly diagnosed HIV infection among residents of Idaho, including concurrent HIV Stage 3 (AIDS) diagnosis.

HIV infection as defined by CDC is inclusive of AIDS, where AIDS is now referred to as "HIV Stage 3."  
HIV Stage 1, 2, 3, and Unknown are surveillance case definitions which take into account CD4 levels and clinical criteria.

For more information, visit <https://www.cdc.gov/nndss/conditions/>



Idaho District Health Departments		
District 1	Coeur d'Alene	415-5100
District 2	Lewiston	799-3100
District 3	Caldwell	455-5300
District 4	Boise	375-5211
District 5	Twin Falls	734-5900
District 6	Pocatello	233-9080
District 7	Idaho Falls	522-0310

Reported Idaho HIV and AIDS Cases Presumed Living <sup>1</sup> by Diagnostic Status as of June 30, 2017							
District	HIV Cases <sup>2</sup>	HIV Cases Reported Deceased	Presumed Living with HIV	AIDS Cases <sup>3</sup>	AIDS Cases Reported Deceased	Presumed Living with AIDS	Reported HIV and AIDS Cases Presumed Living
1	102	11	91	194	84	110	201
2	60	9	51	82	31	51	102
3	122	16	106	186	66	120	226
4	448	48	400	598	240	358	758
5	91	9	82	130	50	80	162
6	89	13	76	146	60	86	162
7	68	6	62	114	49	65	127
<b>TOTAL</b>	<b>980</b>	<b>112</b>	<b>868</b>	<b>1,450</b>	<b>580</b>	<b>870</b>	<b>1,738</b>

<sup>1</sup> Presumed living: Cases ever reported in Idaho and not determined to be deceased.

<sup>2</sup> Number of reported cases of HIV Infection (not yet HIV Stage 3 [AIDS]) EVER REPORTED in Idaho, excluding those who were concurrently diagnosed with HIV Stage 3 (AIDS) or have progressed to HIV Stage 3 (AIDS).

<sup>3</sup> Number of reported cases of Stage 3 HIV Infection (AIDS) ever reported in Idaho.

**HIV and AIDS (HIV Stage 3) Among Residents of Idaho at Diagnosis by Case Status, January 1, 1984 - June 30, 2017**

		HIV		AIDS		COMBINED	
<b>Sex</b>							
Male		493	83%	728	85%	1,221	84%
Female		100	17%	128	15%	228	16%
Total		593	100%	856	100%	1,449	100%
<b>Pediatric/Adult &amp; Adolescent</b>							
Adults/Adolescents (13 or above)		590	99%	849	99%	1,439	99%
Pediatric (<13 years)		3	1%	7	1%	10	1%
Total		593	100%	856	100%	1,449	100%
<b>Race/Ethnicity</b>							
Hispanic - All Races		68	11%	104	12%	172	12%
Not Hispanic	American Indian/Alaska Native	10	2%	12	1%	22	2%
	Asian	8	1%	2	0%	10	1%
	Black	35	6%	34	4%	69	5%
	Native Hawaiian/Pacific Islander	0	0%	0	0%	0	0%
	White	451	76%	690	81%	1,141	79%
	Legacy Asian/Pacific Islander	0	0%	2	0%	2	0%
	Multi-race	5	1%	7	1%	12	1%
	Unknown	16	3%	7	1%	21	1%
	Total	593	100%	856	100%	1,449	101%
<b>Exposure Categories - Adults/Adolescents</b>							
Men who have sex with men (MSM)		288	49%	431	51%	719	50%
Injection drug use		55	9%	90	11%	145	10%
MSM and inject drugs		62	11%	78	9%	140	10%
Hemophilia/Coagulation disorder		1	0%	18	2%	19	1%
High-risk heterosexual contact		64	11%	103	12%	167	12%
Presumed heterosexual contact		15	3%	17	2%	32	2%
Receipt of blood, components, or tissue		2	0%	11	1%	13	1%
Other/risk not reported or identified		103	17%	101	12%	204	14%
Total		590	100%	849	100%	1,439	100%
<b>Exposure Categories - Pediatric</b>							
Hemophilia/Coagulation disorder		0	0%	0	0%	0	0%
Mother with/at risk for HIV infection		3	100%	4	57%	7	70%
Receipt of blood, components, or tissue		0	0%	2	29%	2	20%
Other/risk not reported or identified		0	0%	1	14%	1	10%
Total		3	100%	7	100%	10	100%
<b>Cases by Age Group</b>							
Under 5		3	1%	3	0%	6	0%
5-12		0	0%	1	0%	1	0%
13-19		16	3%	8	1%	24	2%
20-29		242	41%	124	14%	366	25%
30-39		168	28%	334	39%	502	35%
40-49		114	19%	248	29%	362	25%
Over 49		50	8%	138	16%	188	13%
Unknown		0	0%	0	0%	0	0%
Total		593	100%	856	100%	1,449	100%

This table excludes cases of HIV or AIDS (HIV Stage 3) first diagnosed in other states. Data are provisional. Percentages may not sum to 100% due to rounding. Reports of HIV and AIDS cases should be made to the Idaho Department of Health & Welfare as required by Idaho Code 39-602. Data collection and publication are made possible through a federal grant from the Centers for Disease Control and Prevention – Department of Health and Human Services. Idaho statistics are posted on the Internet four times per year. Interested parties may e-mail Allen Haumann at [allen.haumann@dhw.idaho.gov](mailto:allen.haumann@dhw.idaho.gov) to be placed on an e-mail alert group to be signaled when each new issue of statistics is published.