



## Long-Acting Narcotic Analgesics Therapeutic Class Review (TCR)

---

**March 15, 2020**

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Magellan Rx Management.

All requests for permission should be mailed to:

Magellan Rx Management  
Attention: Legal Department  
6950 Columbia Gateway Drive  
Columbia, Maryland 21046

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to [PSTCReDitor@magellanhealth.com](mailto:PSTCReDitor@magellanhealth.com).

---

March 2020

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.  
© 2004–2020 Magellan Rx Management. All rights reserved.

**MagellanRx**  
MANAGEMENT<sup>SM</sup>

## FDA-APPROVED INDICATIONS

Drug	Manufacturer	Schedule	Indication(s)
buprenorphine (Belbuca™) <sup>1</sup>	Biodelivery Sciences	CIII	<ul style="list-style-type: none"> <li>Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate</li> </ul>
buprenorphine (Butrans®) <sup>2</sup>	generic, Purdue	CIII	
fentanyl transdermal (Duragesic®) <sup>3</sup>	generic, Janssen	CII	
hydrocodone extended-release (Hysingla® ER) <sup>4</sup>	Purdue	CII	
hydrocodone extended-release (Zohydro® ER) <sup>5</sup>	generic, Pernix	CII	
hydromorphone extended-release (Exalgo®) <sup>6</sup>	generic, Mallinckrodt	CII	
methadone (Dolophine®, Diskets®, Methadose™) <sup>7</sup>	generic, West-Ward/Hikma; Mallinckrodt	CII	<ul style="list-style-type: none"> <li>Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate;</li> <li>Detoxification and maintenance treatment of opioid addiction</li> </ul>
morphine sulfate controlled-release (MS Contin®) <sup>8</sup>	generic, Rhodes	CII	<ul style="list-style-type: none"> <li>Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate</li> </ul>
morphine sulfate extended-release* (Arymo ER™) <sup>9</sup>	Egalet/Zyla	CII	
morphine sulfate extended-release† (Kadian®) <sup>10</sup>	generic, Actavis/Allergan	CII	
morphine sulfate extended-release (Morphabond™ ER) <sup>11</sup>	Daiichi Sankyo	CII	
morphine sulfate extended-release and naltrexone (Embeda®) <sup>‡12</sup>	Pfizer	CII	
oxycodone controlled-release (Oxycontin®) <sup>§13</sup>	generic <sup>‡</sup> , Purdue	CII	
oxycodone extended-release (Xtampza ER™) <sup>14</sup>	Collegium	CII	
oxymorphone extended-release <sup>¶15</sup>	generic	CII	
tapentadol extended-release (Nucynta® ER) <sup>16</sup>	Depomed/Collegium	CII	<ul style="list-style-type: none"> <li>Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate;</li> <li>Neuropathic pain associated with diabetic peripheral neuropathy (DPN) in adults severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate</li> </ul>

**FDA-Approved Indications (continued)**

Drug	Manufacturer	Schedule	Indication(s)
tramadol extended-release (Conzip®) <sup>17</sup>	generic, Vertical	CIV	<ul style="list-style-type: none"> <li>Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate</li> </ul>
tramadol extended-release <sup>18,19</sup>	generic	CIV	

\* Effective September 28, 2018, Egalet/Zyla announced that they are discontinuing the promotion and manufacture of Arymo ER.<sup>20</sup> Product within the expiry date may be available.

† Pfizer has discontinued manufacture of Avinza® non-abuse-deterrent extended-release morphine sulfate capsules. Generic products, approved with Avinza as the reference product, are still available.

‡ In October 2019, Pfizer announced they will be stopping the sale and distribution of Embeda® ER (morphine/naltrexone) capsules as of November 15, 2019; product may remain available until supplies are depleted.<sup>21</sup>

§ All generic products of Oxycontin are authorized generics.

¶ In July 2017, Endo announced that they will voluntarily remove Opana® ER (oxymorphone ER) from the market in response to a request from the Food and Drug Administration (FDA) who determined that the product’s risks outweigh its benefits.<sup>22,23</sup> Although Opana ER had abuse-deterrent *properties* following its reformulation in 2012, it was not considered an abuse-deterrent *formulation* by the FDA. In addition, oxymorphone extended-release products were approved based on the original formulation of Opana ER (originally approved in 2006), which did not contain abuse-deterrent properties.

|| Generics were approved to the reference product of either Ryzolt® (previously manufactured by Purdue Pharma) or Ultram® ER (previously manufactured by Janssen).

lonsys®, a fentanyl iontophoretic transdermal system for pain management in hospitalized patients only, will not be addressed in this review.

**OVERVIEW**

Pain of multiple etiologies remains a substantial problem for many patients presenting in the clinical setting.<sup>24</sup> Pain management must be individualized for each of these patients. There are many opioid analgesic products available, differing in specific opioid, dosage form, abuse-deterrent properties, and duration of action. In this review, the terms “narcotic” and “opioid” are used interchangeably.

Data from 2012 demonstrated that approximately 11.2% of adults report daily pain, which is greatly misunderstood. Historically, data have suggested that pain may be undertreated, but newer estimates imply that opioid treatment for pain may be overutilized.<sup>25</sup> An estimated 20% of patients who presented to outpatient providers with noncancer pain or pain-related diagnoses, whether acute or chronic, received an opioid prescription.<sup>26</sup> Caregivers’ misconceptions regarding opiate doses, duration of analgesic effect, and fear of addiction may be responsible for the historical undertreatment of pain, and some of the historical increase in opioid use is thought to be related to prescribers worrying they are undertreating their patients’ pain.<sup>27,28</sup> Unfortunately, approximately 67,367 people have died from overdoses related to opioid pain medications in the United States (US) in 2018.<sup>29</sup> Likewise, drug related deaths have tripled from 1999 to 2015, and during 2015 alone 33,091 persons in the US died from opioid related overdoses.<sup>30</sup> Opioid related overdose deaths were higher among males (20.4%) in comparison to females (9.4%).<sup>31</sup> In 2018, 15% of the US population received ≥ 1 opioid prescription. Annually from 2012, there has been a continued decreased in opioid prescribing.<sup>32</sup> Likewise, the yearly rate for high-dose opioid prescriptions has decreased by 66.1% from 2006 to 2018. Regardless, persistent pain that is uncontrolled may have clinical, psychological, and social consequences; thus, it is critical to weigh the risks and benefits of opioid use and reevaluate patients routinely for appropriate dose, duration, and treatment choice, including both pharmacologic and nonpharmacologic modalities.

Different management techniques are utilized for acute and chronic pain. When properly used, long-acting opioids decrease administration frequency, decrease the incidence of adverse effects, and increase periods of consistent pain control. While definitions vary, chronic pain is generally defined as pain lasting > 3 months or past the time required for normal tissue healing. It has various etiologies, including injury, inflammation, and underlying medical conditions.<sup>33</sup>

## Treatment Guidelines

The World Health Organization's (WHO) guidelines for cancer pain management for adults recommend a 3-stepped approach with consideration for the type of pain and response to therapy.<sup>34</sup> If pain occurs, WHO recommends prompt oral administration of drugs in the following order: non-opioids (acetaminophen, non-steroidal anti-inflammatory drugs [NSAIDs]); then, as necessary, mild opioids (codeine); then strong opioids, until the patient is free of pain. These agents can be used alone or in combination to achieve or maintain effective and safe pain management; any opioid can be considered for pain relief in the maintenance setting.<sup>35</sup> Analgesics should be given around the clock, rather than on-demand.<sup>36</sup> Psychosocial care and adjuvant therapies (e.g., corticosteroids, antidepressants, anticonvulsants, bisphosphonates) should also be considered as part of a comprehensive care plan.

In 2016, the American Society of Clinical Oncology (ASCO) published guidelines for chronic pain management in adult cancer survivors.<sup>37</sup> For pharmacologic treatment, systemic non-opioid analgesics and adjuvant analgesics are suggested for chronic pain and to improve function. They state that opioids may be used in those who do not respond to more conservative management.

Similarly, the National Comprehensive Cancer Network (NCCN) published guidelines on the treatment of cancer pain in adults in 2019 specific to pain scale ratings.<sup>38</sup> On a scale of 0 to 10, the pain scale is rating of 1 to 3 is mild pain, 4 to 7 is moderate pain, and  $\geq 8$  is severe pain. The recommendation in opioid-naïve patients with mild pain (1 to 3) is non-opioid and adjuvant therapies unless contraindications due to adverse effects or drug interactions are present. For moderate pain (4 to 7), NCCN recommends adding a short-acting opioid as needed. NCCN defines opioid tolerant patients by the FDA definition: patients receiving at least 25 mcg/h fentanyl patch, at least 60 mg of morphine daily, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily, or an equianalgesic dose of another opioid for a week or longer. The NCCN recommendation for opioid-tolerant patients is the same as opioid-naïve patients, except that they specify titrating short-acting opioid dose by 30% to 50% daily. Recommendations for both opioid-naïve and opioid-tolerant patients include opioid rotation if limiting adverse effects are noted and opioid reassessment of efficacy and adverse effects in 1 to 4 weeks. Long-acting opioids are recommended if 3 to 4 daily doses of short-acting opioid are consistently needed. If pain is persistent, NCCN recommends that the opioid should be scheduled with rescue dose as needed. Consideration for treatment in the hospital or hospice setting for patient-specific goals is recommended for any severe pain ( $\geq 8$ ). NCCN recommends against the use of meperidine (due to central nervous system [CNS] toxicity) and mixed agonist-antagonists (limited usefulness) for cancer pain.<sup>39</sup> Also, they recommend to consider supplementing with doses of short-acting opioid when using methadone as a long-acting opioid. NCCN also provides extensive guidance on dosing, adverse effect management, and pain assessment.

A position statement released by the American Academy of Neurology (AAN) states that the risks of opioids outweigh their benefits for treating chronic non-cancer pain.<sup>40</sup> AAN states that, while opioids

may provide short-term pain relief, there is no proof that they maintain pain relief or improve patients' ability to function over long periods of time without a serious risk of overdose, dependence, or addiction.

In 2016, the Centers for Disease Control and Prevention (CDC) released guidelines for prescribing opioids for chronic pain outside of active cancer, palliative, and end-of-life care.<sup>41</sup> These CDC guidelines on opioid prescribing are not intended to deny opioid therapy for pain management in patients with chronic pain, sickle cell disease, undergoing cancer treatment, and cancer survivors with chronic pain; rather, the guidelines aim to ensure that clinicians and patients consider all safe and effective treatment options. These guidelines include 12 recommendations: 3 regarding when to initiate or continue opioids for chronic pain; 4 regarding opioid selection, dosage, duration, follow-up, and discontinuation; and 5 regarding assessing risk and addressing harms of opioid abuse.<sup>42</sup> The guidelines prefer nonpharmacologic and nonopioid pharmacologic therapy for chronic pain, and recommend a full individual assessment, including risk evaluation and realistic treatment goal setting, prior to prescribing opioids for chronic pain. If opioids are deemed appropriate for a patient's chronic pain, they recommend initial treatment with immediate-release opioids instead of extended-release opioids, which should be prescribed at the lowest effective dose. They further specify that doses of  $\geq 50$  morphine milligram equivalents (MME)/day should prompt reassessment of the individual's benefits and risks and use of  $\geq 90$  MME/day should be avoided without justification. They state that long-term opioid use often begins with acute pain treatment; thus, opioids for acute pain should be immediate-release, at the lowest effective dose, and the quantity should not exceed the expected duration of pain severe enough to require opioids (typically 3 days and with  $> 7$  days rarely needed). They recommend reassessment within 1 to 4 weeks to determine benefits, harms, and appropriate dosing and continued follow up at least every 3 months. At these visits, efforts should be made to optimize other therapies and taper or discontinue opioids as able and as risks outweigh the individual's benefits. In order to decrease risks, the guidelines recommend avoiding concurrent use of benzodiazepines when possible and risk management strategies, such as offering naloxone in high-risk individuals (e.g., history of overdose, history of substance abuse, doses  $\geq 50$  MME/day, concurrent benzodiazepine use). Likewise, they recommend urine drug testing at baseline and annually and review of state prescription drug monitoring programs (PDMPs) at baseline and every 3 months. Prescribers should also offer treatment for opioid use disorder (e.g., buprenorphine or methadone in combination with behavioral therapies).

In 2017, the American College of Physicians (ACP) updated their guidelines on noninvasive treatments for acute, subacute, and chronic low back pain.<sup>43</sup> The guidelines recommend nonpharmacologic treatment (e.g., superficial heat, massage, acupuncture, spinal manipulation) in most patients with acute or subacute low back pain. In cases when a pharmacologic treatment is preferred, an NSAID or skeletal muscle relaxant is recommended. For the treatment of chronic low back pain, nonpharmacologic treatment is also preferred. Those with an inadequate response to nonpharmacologic therapy may be treated with an NSAID as first-line therapy and tramadol or duloxetine as second-line therapy. The guidelines state that opioids should only be considered in those who have failed these prior therapies and if the benefits outweigh the risks on an individualized basis.

Also in 2017, the American Society of Interventional Pain Physicians (ASIPP) updated opioid prescribing guidelines for the management of patients with chronic, non-cancer pain.<sup>44</sup> ASIPP recommends that medical necessity of opioids should be established based on an average moderate to severe (pain or disability level  $\geq 4$  on a 0 to 10-point scale). Regarding specific products, ASIPP recommends initiation of opioids at low doses with short-acting agents and appropriate monitoring. They consider daily doses of

≤ 40 morphine milligram equivalents (MME) as low dose, 41 to 90 MME as moderate dose, and ≥ 91 MME as high dose opioids. They recommend methadone only after failure of other opioid therapy and state long-acting opioids should be avoided during opioid initiation and recommend long-acting or high dose opioids only in special circumstances in which there is severe, intractable pain. They state there is similar effectiveness for long-acting and short-acting opioids, but there are greater risks of adverse events with long-acting opioids. However, they do not recommend 1 specific short-acting opioid over another. ASIPP also recommends that all patients should be screened for opioid abuse and that providers should use urine drug testing and prescription drug monitoring programs to monitor for abuse.

The HIV Medicine Association (HIVMA) of the Infectious Disease Society of America (IDSA) has issued the first comprehensive guidelines for managing chronic pain in people living with the human immunodeficiency virus (HIV) that covers musculoskeletal, arthritic, and neuropathic pain types (not cancer pain).<sup>45</sup> It is recommended that all persons living with HIV be screened for chronic pain and if positive for pain, multidisciplinary treatment focused on nondrug therapies should be offered, followed by non-opioid drugs (gabapentin [preferred], serotonin norepinephrine reuptake inhibitors, tricyclic antidepressants, or pregabalin). Other options include capsaicin, medical cannabis (in select patients), and alpha lipoic acid. The guidelines recommend against use of lamotrigine and against opioids as first-line.

The American Association of Oral & Maxillofacial Surgeons (AAOMS) issued a White Paper regarding opioid prescribing for acute and postoperative pain management.<sup>46</sup> NSAIDs are recommended over opioids as first-line therapy to manage acute and post-operative pain. If an opioid is needed, the lowest dose for the shortest duration should be used and extended-release formulations avoided.

The Institute for Clinical and Economic Review (ICER) has published a final report on abuse-deterrent formulation (ADF) opioids.<sup>47</sup> The group concluded that the evidence is adequate to suggest a reduced risk of abuse among individual patients prescribed Oxycontin compared to non-ADF opioids; however, evidence is not sufficient to show a reduced risk of abuse for individual patients being prescribed any other abuse-deterrent ER opioid assessed in the report (Hysingla ER, Morphabond, Xtampza ER, Arymo ER).

In 2018, the Scientific and Medical Advisory Board of the Restless Legs Syndrome (RLS) Foundation published new guidance stating that opioid therapy is a mainstay of refractory RLS management and low total daily doses of opioids may be life-transforming for patients with refractory RLS.<sup>48</sup> Therapy should be started with a short-acting formulation to determine response, but long-acting opioids are preferred. Most commonly used agents are oxycodone and methadone, but tramadol, codeine, morphine, and hydrocodone can also be considered. Controlled-release medication should be used for evening dosage and short-acting drugs, if needed, during the day.

In March 2020, the American Academy of Pain Medicine (AAPM) published evidence-based clinical guidelines on the diagnosis and treatment of low back pain.<sup>49</sup> There is limited information provided regarding opioids in this setting as the opioid crisis occurred after the guideline development was already underway. However, it is suggested that use of opioid pain medications for treating low back pain be limited and restricted to a short duration (grade of recommendation: B). In contrast, non-selective NSAIDs are suggested for the treatment of low back pain (grade of recommendation: B).

To help curb the potentially fatal effects of opioid overdoses, the Department of Health and Human Services (HHS) released guidelines recommending that naloxone should be prescribed to individuals at

risk for opioid overdose, including individuals who are on relatively higher doses of opioid ( $\geq 50$  MME/day), those with select respiratory comorbidities (e.g., sleep apnea, chronic obstructive pulmonary disease [COPD]), take other medications which enhance opioid complications (e.g., benzodiazepines), or have other non-opioid substance abuse disorders or mental health disorders.<sup>50</sup> In October 2019, the HHS published a new guideline for clinicians on dosage reduction or discontinuation of long-term opioid analgesics.<sup>51</sup> This guidance discusses the risks of opioid taper and advises that opioids should not be quickly tapered or discontinued abruptly due to the potential for opioid withdrawal which can result in acute withdrawal symptoms, pain exacerbation, psychological distress, and suicidal ideation in patients who are physiologically dependent. Except for life-threatening circumstances (e.g., impending overdose), it is not recommended to abruptly reduce or discontinue opioid dosing. Guidance details situations when it may be appropriate to taper to a reduced dosage (e.g., pain improvement, patient request, no clinically meaningful improvement in pain or function with opioids, increasing doses without improvements in pain, signs of opioid misuse, side effects impacting function or quality of life, risks for an impending overdose/serious event, concurrent medications or comorbidities increasing the risk for adverse events, extended treatment period without clear benefits versus harms). Other key recommendations include: referring patients with serious mental illness, high suicide risk, or suicidal ideation to a behavioral health provider prior to taper; assessing patients for opioid use disorder if they show signs of opioid misuse and offering medication-assisted treatment if appropriate; advising patients of risks for overdose if they abruptly return to their higher dose; tapering by 5% to 20% every 4 weeks is common, but longer tapering schedules may be required; and considering transition to buprenorphine for patients on high doses and unable to taper.

## Opioid Regulation

Soon after its approval in the US in 1995, diversion and abuse of tramadol were reported. This led to the addition of warnings regarding the abuse potential of tramadol to the product labeling by the Food and Drug Administration (FDA). Tolerance, dependence, and addiction to tramadol have been demonstrated and abrupt discontinuation of the drug can result in withdrawal symptoms. Effective in August 2014, tramadol-containing products were placed into Schedule IV of the Controlled Substance Act.<sup>52</sup>

In 2013, the FDA announced class-wide safety labeling changes and new post-market study requirements for all extended-release and long-acting (ER/LA) opioid analgesics intended to treat pain. The updated indication states that ER/LA opioids are indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment for which alternative treatment options are inadequate.<sup>53</sup>

In April 2015, the FDA released guidance to the industry regarding the development of opioid drug products with abuse-deterrent properties to limit the misuse, abuse, and potentially death associated with these products.<sup>54</sup> This guidance states that abuse-deterrent properties discourage abuse but may not fully prevent abuse. Examples of abuse-deterrent properties include physical/chemical barriers to prevent breakdown of the product, agonist/antagonist combinations, added aversive substances, unique drug delivery systems (e.g., depot injections, implants), or new molecular entities or prodrugs with novel effects (e.g., altered receptor binding or require enzymatic activation). In order for the FDA to approve a formulation as an abuse-deterrent formulation, an abuse-deterrent property is not sufficient; the manufacturer should demonstrate that the product deters abuse in studies. The types of studies the FDA requires are laboratory-based *in vitro* manipulation and extraction studies, pharmacokinetic studies, and

clinical abuse potential studies, which commonly measure drug-liking by subjects without pain and with current addiction behaviors. The following agents have demonstrated abuse-deterrence in studies, thus meeting FDA requirements for abuse-deterrent formulations: Arymo ER tablets, Embeda capsules, Hysingla ER tablets, Morphabond tablets, Oxycontin biconcave tablets, oxycodone ER (authorized generics of Oxycontin) tablets, and Xtampza ER capsules.<sup>55,56,57</sup> As a result, the labeling of these products includes information regarding abuse-deterrence, reflecting the FDA's approval of these products as abuse-deterrent formulations. The FDA also published information on abuse deterrence studies to guide new product development and evaluations of generic formulations.<sup>58</sup>

In February 2016, the FDA announced plans to reassess their approach to opioid medications with a focus on policies to reverse the epidemic of deaths associated with opioid use.<sup>59</sup> Select components of the action plan include the use of an expert advisory committee prior to the approval of an opioid without abuse-deterrent properties, the formation of a Pediatric Advisory Committee to review pediatric labeling for new products, an update of risk evaluation and mitigation strategy (REMS) requirements and improvement in access to abuse-deterrent formulations, naloxone, and other treatment options for patients with opioid-use disorders.

In April 2017, the FDA announced the restricted the use of codeine and tramadol medicines in children due to serious risks, including slowed or difficult breathing and death, which are a greater risk in children younger than 12 years.<sup>60</sup> These medications should also be limited in some older children. The FDA also recommended against the use of codeine and tramadol medicines in breastfeeding mothers due to possible harm to their infants.

In September 2017, the FDA determined that a REMS is necessary for immediate-release (IR) opioid analgesics to ensure that the drug benefits outweigh the risks. As a result, the FDA sent letters to IR opioid analgesic manufacturers informing them that their products that are intended for outpatient setting use will be subject to the same REMS requirement as the ER/LA opioid analgesics.

In October 2018, the Substance Use-Disorder Prevention that Promotes Opioid Recovery and Treatment for Patients and Communities (SUPPORT) Act was passed to reduce unnecessary opioid exposure, and lower the rate of new opioid addiction.<sup>61</sup> The provision requires the Drug Enforcement Administration (DEA) to provide drug manufacturers and distributors access to drug supply chain information from point of manufacture to point of sale.<sup>62</sup> This system, Automated Reports and Consolidated Orders System (ARCOS), helps drug manufacturers and distributors identify, report, and stop suspicious orders of opioids and reduce diversion rates. Additionally, the Act gave authority to the FDA to require manufacturers to develop disposal technologies to allow patients easier disposal of unused medications out of their homes.<sup>63</sup>

In April 2019, the FDA required label changes to guide prescribers on gradual, individualized tapering of opioid pain medications due to harm reported from sudden discontinuation.<sup>64</sup> The FDA was prompted to make these changes after receiving several reports of serious harm in patients physically dependent on opioid medications undergoing sudden discontinuation or rapid dose decrease. These effects include serious withdrawal symptoms, uncontrolled pain, psychological distress, and suicide. Withdrawal and uncontrolled pain from rapid dose decreases can lead patients to seek illicit substances to treat pain, which could be confused with abuse drug-seeking behaviors. Tapering schedules should be patient-specific based on multiple patient factors such as duration of therapy, type of pain treated, and physical

and psychological patient attributes. The FDA suggests tapering by an increment of no more than 10% to 25% every 2 to 4 weeks.

In April 2019, the FDA announced as a part of their “Remove the Risk” campaign for removing an properly disposing of unused prescription opioids from the home, they were targeting females aged 35 to 64 years since the FDA considers them as common gatekeepers in the home for healthcare decisions and medication management.<sup>65</sup> The campaign involves a toolkit of materials for news media, healthcare providers, consumer groups, and other organizations.

The FDA continues to evaluate and issue guidance on the use and development of opioids.<sup>66</sup> Part of the FDA’s efforts include commissioning the National Academy of Sciences, Engineering, Medicine (NASEM) to develop a consensus report on "Framing Opioid Prescribing Guidelines for Acute Pain." This report was released in December 2019 and provides a framework for assessing opioid prescribing in order to develop a clinical practice guideline and determine areas of future research. The analysis also includes an evaluation of current opioid prescribing guidelines with identification of areas for additional research.<sup>67</sup> Similarly, the DEA has announced reductions in the quantity of Schedule II opioid prescriptions that may be manufactured in anticipation of a decline in need.<sup>68,69,70</sup>

## PHARMACOLOGY<sup>71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90</sup>

Opioid agonists reduce pain by acting primarily through interaction with opioid mu-receptors located in the brain, spinal cord, and smooth muscle. The primary site of therapeutic action is the central nervous system (CNS). Opioid agonists produce respiratory depression by direct action on the brain stem respiratory center. Buprenorphine (Belbuca, Butrans) differs in that it is a partial agonist/antagonist of opioid receptors.

Naltrexone, a component of Embeda, is a centrally-acting mu-receptor antagonist that reverses the analgesic effects of mu-receptor agonists by competing for binding sites with opioids.

Tapentadol (Nucynta ER) is a centrally-acting synthetic analgesic and exerts its analgesic effects without a pharmacologically active metabolite. The exact mechanism of action is unknown. Tapentadol also inhibits norepinephrine reuptake.

In addition to binding to the mu-opioid receptors, tramadol (Conzip, generic) exerts its effect through weak inhibition of norepinephrine and serotonin reuptake.

## PHARMACOKINETICS<sup>91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110</sup>

Drug	Half-Life (hr)	Tmax (hr)	Excretion
buprenorphine (Belbuca)	27.6	3	69% metabolized and eliminated in feces and approximately 30% excreted in urine
buprenorphine (Butrans)	26	about 48-72	70% metabolized and eliminated in feces and approximately 27% excreted in urine
fentanyl transdermal (Duragesic)	3–12	28.8–35.8	75% metabolized and renally eliminated
hydrocodone ER (Hysingla ER)	7–9	14–16	primarily metabolized and renally eliminated
hydrocodone ER (Zohydro ER)	8	5	highly metabolized; 26% eliminated in urine
hydromorphone ER (Exalgo)	8–15	12–16	highly metabolized; 75% eliminated in urine
methadone (Dolophine, Diskets, Methadose)	8–59	1–7.5	highly metabolized; eliminated in urine and feces
morphine sulfate CR (MS Contin)	2–15	~ 1.5	90% metabolized and renally eliminated
morphine sulfate ER (Arymo ER)	2–15	4.5–6.5	90% metabolized and renally eliminated
morphine sulfate ER (Kadian)	2–15	8.6–10.3	90% metabolized and renally eliminated
morphine sulfate ER (Morphabond)	nr	2–4	90% metabolized and renally eliminated
morphine sulfate ER/naltrexone (Embeda)	29	7.5	90% metabolized and renally eliminated
oxycodone CR (Oxycontin)	4.5	1.6–3.2	primarily metabolized and renally eliminated
oxycodone ER (Xtampza ER)	~5.6	4–4.5	primarily metabolized and renally eliminated
oxymorphone ER	9.4–11.3	1–2	highly metabolized; eliminated in urine and feces
tapentadol ER (Nucynta ER)	5	3–6	97% metabolized and renally eliminated
tramadol ER (Conzip)	tramadol 10 metabolites 11	tramadol 5.9 metabolites 11	30% excreted as tramadol, 60% excreted as active metabolites in the urine
tramadol ER	tramadol 7.9 metabolites 8.8	tramadol 12 metabolites 15	30% excreted as tramadol, 60% excreted as active metabolites in the urine

hr = hours; Tmax = time to maximum serum concentration; nr = not reported

## CONTRAINDICATIONS/WARNINGS<sup>111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130</sup>

### Contraindications

The agents in this review are contraindicated in patients with a known hypersensitivity to any component of the product. Hydromorphone (Exalgo) is also contraindicated in patients with known hypersensitivity to sulfites. Oxymorphone ER is contraindicated in patients with a known hypersensitivity to morphine analogs, such as codeine.

Long-acting opioids are contraindicated in patients who are not opioid-tolerant; patients who have acute or severe bronchial asthma; situations of significant respiratory depression (in the absence of resuscitative equipment or monitors); and patients with known or suspected paralytic ileus.

In general, long-acting opioids are not indicated as an as-needed analgesic; for example, for use in the management of acute pain, mild pain, or intermittent pain, or in patients who require opioid analgesia for a short period of time in the management of post-operative pain, including use after outpatient or day surgeries (e.g., tonsillectomies).

Oxymorphone ER is contraindicated in patients with moderate and severe hepatic impairment.

Monoamine oxidase inhibitors (MAOIs) may intensify the actions of opioids; these long-acting opioids should not be taken in patients taking MAOIs or within 14 days of stopping MAOI treatment.

Tramadol ER (Conzip, generic) is contraindicated in patients who have acute intoxication with alcohol, hypnotics, centrally-acting analgesics, opioids, or psychotropic drugs, and situations where opioids may be contraindicated. In April 2017, the FDA released a Drug Safety Communication regarding the use of tramadol in pediatric patients and is requiring labeling revisions for these products.<sup>131</sup> Due to case reports of respiratory depression, some leading to death, the FDA required a new contraindication for the use of tramadol to treat pain following tonsillectomy and/or adenoidectomy in patients < 18 years old. An additional contraindication in children < 12 years old was also added. Notably, tramadol formulations within this class review are approved only in adults. Additional warnings were also required and are described below.

## Warnings

### **Boxed Warnings**

Buprenorphine transdermal (Belbuca, Butrans), methadone (Dolophine, Diskets, Methadose), fentanyl transdermal (Duragesic), morphine sulfate ER/SR (Arymo ER, Kadian, Morphabond, MS Contin), morphine sulfate ER/naltrexone (Embeda), hydrocodone ER (Hysingla ER, Zohydro ER), hydromorphone ER (Exalgo), morphine sulfate controlled-release (MS Contin), oxymorphone ER, oxycodone CR (Oxycontin), oxycodone ER (Xtampza ER), tapentadol ER (Nucynta ER), and tramadol ER (Conzip, generic) boxed warnings include: monitoring for signs of misuse, abuse, and addiction during therapy; fatal respiratory depression may occur, with the highest risk at initiation and with dose increases; and accidental exposure can result in the fatal overdose of the above medications in children. Concomitant use of opioids with benzodiazepines or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. These should only be used in combination when alternative treatment options are inadequate and doses and duration should be limited. Patients should be monitored closely for any signs and symptoms of respiratory depression and sedation.

In August 2016, the FDA issued a Drug Safety Communication regarding the risk of death when combining opioid pain medications or cough medicine with benzodiazepines and other CNS depressants. The FDA required an additional boxed warning for all opioids describing this risk (as described above).<sup>132</sup>

The boxed warning for methadone (Dolophine, Diskets, Methadose) indicates that cardiac and respiratory deaths have been reported during initiation and conversion of pain patients to methadone treatment from other opioid agonists. Cases of QT interval prolongation and serious arrhythmia have also been observed.

Chronic maternal use of opioids during pregnancy can result in life-threatening neonatal opioid withdrawal syndrome; infants may require treatment if exposed to opioids during gestation.

Respiratory depression is the chief hazard of opioid agonists. Serious or life-threatening hypoventilation may occur at any time during the use of long-acting opioids, especially during the initial 24 to 72 hours following initiation of therapy and following increases in dose. Respiratory depression is more likely to occur in elderly or debilitated patients, usually following large initial doses in non-tolerant patients or when opioids are given in conjunction with other agents that depress respiration, such as alcohol and other central nervous system depressants. Respiratory depression from opioids is manifested by a reduced urge to breathe and a decreased rate of respiration, often associated with the “sighing” pattern of breathing (deep breaths separated by abnormally long pauses). Carbon dioxide retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids. This makes overdoses involving drugs with sedative properties and opioids especially dangerous. Opioids can also lead to central sleep apnea (CSA) in a dose-dependent manner as well as sleep-related hypoxemia; a reduction in dosage may be warranted for patients with CSA.

When the patient no longer requires therapy with agents in this class, doses should be tapered gradually to prevent signs and symptoms of withdrawal in the physically-dependent patient. These agents should not be abruptly discontinued in physically dependent patients; withdrawal syndrome, as well as return of pain, may occur if these agents are discontinued abruptly. Gradually taper the opioid dosage at the time of discontinuation to minimize the risk for withdrawal symptoms.

Patients should not consume alcohol or any products containing alcohol while taking morphine sulfate ER/naltrexone (Embeda) because co-ingestion can result in fatal plasma morphine levels.

Oxycodone CR and ER (OxyContin, Xtampza ER) also carry a boxed warning for drug interactions associated with cytochrome P450 3A4 inhibitor or inducer initiation or discontinuation potentially altering drug exposure which could lead to overdose.

## Other Warnings

Long-acting opioids may worsen increased intracranial pressure and obscure its signs. Patients with increased intracranial pressure, brain tumors, head injury, or impaired consciousness may be more susceptible. Patients at increased risk of hypotension and those in circulatory shock could experience worsened conditions with treatment. Orthostatic hypotension and syncope have also been reported with opioids. Patients at higher risk of hypotension include those with hypovolemia or those taking concurrent products that compromise vasomotor tone (e.g., phenothiazines, general anesthetics).

All agents in this review, except tramadol ER, may cause spasm of the sphincter of Oddi. Monitor patients with biliary tract disease, including acute pancreatitis, for worsened symptoms.

All opioids contain a warning regarding serotonin syndrome when used concomitantly with any serotonergic drug (e.g., monoamine oxidase [MAO] inhibitors, selective serotonin reuptake inhibitors [SSRIs], serotonin and norepinephrine reuptake inhibitors [SNRI], tricyclic antidepressants [TCAs], triptans, linezolid, lithium, or certain muscle relaxants [e.g., cyclobenzaprine, metaxalone]). Serotonin syndrome typically occurs within several hours to a few days following use.<sup>133</sup>

Monitor patients with a history of seizure disorders for worsened seizure control during therapy with buprenorphine, hydromorphone ER, morphine ER/CR, morphine ER/naltrexone, oxycodone CR/ER,

oxymorphone ER, tapentadol ER, and tramadol ER since these agents can aggravate seizure disorder and may induce seizures in some clinical settings. This is not reported for fentanyl transdermal or hydrocodone ER medications.

Opioids inhibit the secretion of adrenocorticotrophic hormone (ACTH) and cortisol. All opioids carry a warning for adrenal insufficiency; if an opioid causes adrenal insufficiency, treat with corticosteroids and withdraw the opiate as appropriate.<sup>134</sup>

Initiation of concurrent therapy with CYP3A4 inhibitors, or discontinuation of CYP3A4 inducers with buprenorphine, fentanyl transdermal, hydrocodone ER, and oxycodone CR/ER, can result in a fatal opioid overdose.

All tablets or capsules of oral long-acting opioids, with the exception of oxycodone ER (Xtampza ER), should be swallowed whole. Taking broken, chewed, or crushed tablets or capsules can lead to a rapid release and absorption of a potentially fatal dose of most drugs in this class.

Buprenorphine and fentanyl patches are for transdermal use on intact skin only. Avoid exposure to direct heat while wearing the patch because an increased absorption of the drug can occur. Monitor patients who develop increased body temperature for opioid side effects and adjust dose if signs of respiratory or central nervous system depression are seen. Buprenorphine patches should not be used if the pouch seal is broken or the patch is cut, damaged, or changed in any way. Damaged or cut fentanyl patches should not be used.

Use of opioids with other CNS depressants may result in hypotension, sedation, respiratory depression, coma, or death.

Due to the risk of sedation with opioids, this may impair a patient's ability to drive or operate heavy machinery.

If urine testing for clinical management is needed, the assay should be assessed to ensure the sensitivity and specificity are appropriate.

### ***buprenorphine buccal (Belbuca) and buprenorphine transdermal (Butrans)***

Avoid using buprenorphine in patients with long QT syndrome or a family history of the disease. Use caution in patients with hypokalemia, hypomagnesemia, or clinically-unstable cardiac disease. These patients should not exceed a dose of one 20 mcg/hr buprenorphine transdermal patch due to the risk of QTc prolongation.

Cases of cytolytic hepatitis and hepatitis with jaundice have been observed in individuals receiving sublingual buprenorphine for the treatment of opioid dependence. For patients with an increased risk of hepatotoxicity (e.g., history of excessive alcohol intake, liver disease), monitor liver enzyme levels before and periodically during treatment with buprenorphine. The risk of overdose may be greater in patients with moderate to severe hepatic impairment.

Cancer patients with oral mucositis may absorb buprenorphine buccal (Belbuca) more quickly than intended. A dose reduction is recommended in patients with known or suspected mucositis.

Rare cases of severe application site skin reactions have been reported with use of buprenorphine transdermal patches. Onset can occur within days to months after treatment starts. Discontinue therapy if this occurs.

Buprenorphine transdermal has not been studied for use in the management of addictive disorders.

### ***fentanyl transdermal (Duragesic)***

In 2005 and again in 2007, the FDA issued safety communications that highlight important information on appropriate prescribing, dose selection, and the safe use of the fentanyl transdermal system (patch).<sup>135</sup> However, the FDA continues to receive reports of death and life-threatening adverse events related to fentanyl overdose that have occurred when the fentanyl patch was used to treat pain in opioid-naïve patients and when opioid-tolerant patients have applied more patches than prescribed, changed the patch too frequently, and exposed the patch to a heat source. Directions for prescribing and using the fentanyl patch must be followed exactly to prevent death or other serious side effects from fentanyl overdose.

Fentanyl may cause bradycardia; monitor patients with bradyarrhythmias.

### ***hydrocodone bitartrate ER (Zohydro ER/Hysingla ER)***

QTc prolongation has been observed with daily doses of hydrocodone ER 160 mg and should be considered when prescribing this agent in patients with congestive heart failure, bradyarrhythmias, electrolyte imbalances, or who are taking medications known to cause QTc prolongation. Avoid hydrocodone ER in patients with congenital long QT syndrome. A dose reduction or a change to an alternative analgesic should be made in patients who develop QTc prolongation.

Alcohol or alcohol-containing products should not be consumed while taking hydrocodone bitartrate ER as co-ingestion can result in fatal plasma hydrocodone levels.

### ***morphine ER/naltrexone (Embeda)***

Morphine sulfate ER/naltrexone (Embeda) capsules contain pellets of morphine sulfate with a sequestered core of naltrexone. Tampering with the pellets by crushing or chewing causes rapid release and absorption of both morphine and naltrexone, resulting in a potentially fatal morphine dose, particularly in opioid-naïve individuals. In opioid-tolerant patients, the absorption of naltrexone may increase the risk of precipitating withdrawal. Morphine sulfate ER/naltrexone 100/4 mg capsules are for opioid-tolerant patients. Patients should not consume alcoholic beverages or use prescription or non-prescription medications containing alcohol while on this therapy.

Naltrexone may or may not interfere with enzymatic methods for the detection of opioids in the urine depending on the specificity of the test. Consult the test manufacturer for specific details.

### ***oxycodone CR (Oxycontin)***

A reformulation of oxycodone CR (Oxycontin) was approved in 2010 by the FDA, as well as the first reformulated generic version. These new formulations were designed to decrease the likelihood that this medication will be misused or abused, and result in overdose. The reformulation added tamper-resistant features aimed at preserving the controlled release of the active ingredient, oxycodone.<sup>136</sup> Any attempt to dissolve the tablets in liquid results in a gummy substance. The oxycodone CR 60 mg and 80 mg tablets (or a single dose > 40 mg) are to be used in opioid-tolerant patients only, since fatal respiratory depression may occur when administered to patients who are not tolerant to the respiratory depressant effects of opioids.

### ***oxycodone ER (Xtampza ER)***

The risk of overdose may be greater in patients with moderate to severe hepatic impairment compared to patients with normal hepatic function; a lower starting dose is recommended in this population.

### ***oxymorphone ER***

In October 2012, the FDA issued a safety warning that thrombotic thrombocytopenic purpura appears to occur with the use of oxymorphone ER when it is abused and injected intravenously. TTP resulting in kidney failure requiring dialysis have been reported.<sup>137</sup> Oxymorphone ER should only be taken orally.

Endo has voluntarily removed Opana ER from the market, per FDA request.<sup>138,139</sup> Postmarketing data illustrated that the reformulated Opana ER drew concerns that the benefits of the drug no longer outweighed the risks as there was a significant shift in the route of abuse of Opana ER from nasal to injection.

### ***tramadol extended-release (Conzip, generic)***

Tramadol ER should not be prescribed for patients who are suicidal or addiction-prone. Tramadol extended-release formulations are for oral use only, should be swallowed whole, and should not be chewed, dissolved, split, or crushed.

The effects of concomitant use or discontinuation of drugs affecting cytochrome P450 should be taken into consideration in patients using tramadol.

Due to the risk of respiratory depression with tramadol in pediatric patients described above, the FDA is requiring an additional warning recommending against the use of tramadol in adolescents ages 12 to 18 years who are obese or have conditions which may increase the risk of serious breathing problems (e.g., obstructive sleep apnea, severe lung disease).<sup>140</sup> The FDA is also strengthening the warning regarding its use while breastfeeding; tramadol use by women while breastfeeding is not recommended as it may cause significant adverse reactions (e.g., excess sleepiness, difficulty feeding, breathing difficulty) in breastfed infants.

## **Risk Evaluation and Mitigation Strategy (REMS)<sup>141</sup>**

In 2012, the revised REMS for extended-release (ER) and long-acting (LA) opioid medications was introduced. The FDA communicated that this class contains highly potent medications whose approved indication is to treat moderate-to-severe persistent pain for serious chronic conditions. In addition, misuse and abuse of these medications was addressed referencing a serious public health crisis of addiction, overdose, and death. The REMS introduced new safety measures (including dispensing of medication guides and prescriber training). In September 2017, the FDA determined that a REMS is necessary for immediate-release (IR) opioid analgesics to ensure that the drug benefits outweigh the risks. As a result, the FDA sent letters to IR opioid analgesic manufacturers informing them that their products that are intended for outpatient setting use will be subject to the same REMS requirement as the ER/LA opioid analgesics. In September 2018, the REMS program name was changed to the Opioid Analgesic REMS and now applies to immediate-release opioid analgesics for outpatient use as well.

Minor changes to this REMS program have occurred over the years. The current requirements of the REMS program include a medication guideline and healthcare provider training, including prescriber

letters, blueprints for education points, patient counseling documents, and organization and licensing board education letters.

## DRUG

### INTERACTIONS<sup>142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161</sup>

All long-acting narcotics should be used with caution and in reduced doses in patients who are concurrently receiving other narcotic analgesics, muscle relaxants, general anesthetics, phenothiazines, tranquilizers, sedative-hypnotics, tricyclic antidepressants, or other CNS depressants (including alcohol). Respiratory depression, hypotension, and profound sedation or coma may result.

Mixed agonist/antagonist analgesics (e.g., pentazocine, nalbuphine, butorphanol) and partial agonists (buprenorphine) should not be administered to patients who have received or are receiving a course of therapy with a pure opioid agonist analgesic, including morphine ER/naltrexone (Embeda). In these patients, the mixed agonist/antagonist or partial agonist may alter the analgesic effect or may precipitate withdrawal symptoms.

Monoamine oxidase inhibitors (MAOIs) may intensify the actions of fentanyl transdermal, hydrocodone ER, hydromorphone ER, morphine CR/ER, morphine/naltrexone, tapentadol ER, and tramadol ER; these long-acting opioids should not be used in patients taking MAOIs or within 14 days of stopping MAOI treatment.

All opioids contain a warning regarding serotonin syndrome when used concomitantly with any serotonergic drug as described above.<sup>162</sup>

Concomitant use of diuretics and anticholinergics with opioids may reduce the efficacy of the diuretic and increase the risk of urinary retention respectively; monitor and adjust the doses as needed.

Drugs that induce or inhibit cytochrome P450 3A4 enzymes may affect the metabolism of buprenorphine, fentanyl transdermal, hydrocodone ER, oxycodone CR/ER, and tramadol ER. Initiation of concurrent therapy with CYP3A4 inhibitors, or discontinuation of CYP3A4 inducers with these agents, can increase the risk of serious adverse events. However, co-administration of the strong CYP3A4 inhibitor ketoconazole 200 mg with buprenorphine transdermal 10 mcg/hr did not result in changes in the buprenorphine pharmacokinetic profile. Concurrent use of a CYP3A4 inducer may result in a reduced analgesic effect.

Methadone is metabolized by several CYP isoforms, including CYP3A4, CYP2B6, CYP2C19, CYP2C9, and CYP2D6. As a result, use with inhibitors of these enzymes may require a decrease in methadone dose and inducers may result in decreased methadone efficacy. Methadone should be used cautiously in others taking concomitant medications that can affect cardiac conduction or electrolytes; monitoring of cardiac conduction is recommended.

Tramadol is metabolized by CYP2D6 to form an active metabolite. Concomitant administration of CYP2D6 inhibitors, such as quinidine, fluoxetine, paroxetine, and amitriptyline, may reduce metabolic clearance of tramadol increasing the risk for serious adverse events including seizures. Drugs that induce or inhibit cytochrome P450 3A4 enzymes may affect the metabolism of oxycodone CR/ER; monitor and adjust dosage as required if concomitant use is warranted.

The use of buprenorphine, fentanyl transdermal, hydrocodone ER, hydromorphone ER, morphine CR/ER, morphine ER/naltrexone, oxycodone CR, oxymorphone ER, or tapentadol ER with anticholinergic

products may increase the risk of urinary retention and/or severe constipation, which may lead to paralytic ileus. P-glycoprotein (P-gp) inhibitors, such as quinidine, may increase the exposure of morphine by 2-fold. Cimetidine can potentiate respiratory depression when given with oxymorphone or morphine. Morphine and oxycodone can reduce the effect of diuretics.

## ADVERSE EFFECTS<sup>163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182</sup>

Drug	Asthenia	Constipation	Dizziness	Dyspnea	Headache	Nausea	Rash	Somnolence	Vomiting
buprenorphine (Belbuca)	< 1	4–13 (3)	2–6 (1)	< 1	2–6 (3)	10–50 (7)	< 5	1–7 (< 1)	4–8 (< 1)
buprenorphine (Butrans)	1–5	3–13 (1–5)	2–15 (1–7)	≥ 1– < 5	5–16 (5–11)	13–23 (8–11)	3–9 (6)	2–13 (2–4)	4–9 (1)
fentanyl transdermal (Duragesic)	> 10	> 10	3–10	3–10	3–10	> 10	> 1	> 10	> 10
hydrocodone ER (Hysingla ER)	≥ 1 – < 5	≥ 5	≥ 5	< 1	≥ 5	≥ 5	≥ 1– < 5	≥ 5	≥ 5
hydrocodone ER (Zohydro ER)	nr	8 (0)	2 (1)	1–10 (nr)	0 (1)	7 (3)	1–10 (nr)	1 (0)	7 (1)
hydromorphone ER (Exalgo)	1–4 (4)	7–15 (4)	2–4 (1)	reported	5–8 (7)	9–12 (7)	reported	1–9 (0)	6 (4)
methadone (Dolophine, Diskets, Methadose)	reported	reported	reported	reported	reported	reported	reported	reported	reported
morphine sulfate CR (MS Contin)	reported	reported	reported	nr	reported	reported	reported	reported	reported
morphine sulfate ER (Arymo ER)	reported	reported	reported	nr	reported	reported	reported	reported	reported
morphine sulfate ER (Kadian)	2	9	6	3	< 2	7	3	9	2
morphine sulfate ER (Morphabond)	reported	reported	reported	nr	reported	reported	reported	reported	reported
morphine sulfate ER / naltrexone (Embeda)	reported	31.2	4.1	nr	6.9	22.2	< 1	7.3	8
oxycodone CR (Oxycontin)	6 (nr)	23 (7)	13 (9)	1-5	7 (7)	23 (11)	1-5	23 (4)	12 (7)
oxycodone ER (Xtampza ER)	nr	5.2–13 (0.5)	1.6–5.7 (0)	< 1	6.2–13.9 (11.7)	10.9–16.6 (4.6)	1–5	< 1–8.8– (< 1)	4.1–6.4 (1.5)

Adverse effects are reported as a percentage. Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. Incidences for placebo group are reported in parentheses. nr = not reported.

**Adverse Effects (continued)**

Drug	Asthenia	Constipation	Dizziness	Dyspnea	Headache	Nausea	Rash	Somnolence	Vomiting
oxymorphone ER	nr	27.6 (13.2)	17.8 (7.6)	1–10	12.2 (5.6)	33.1 (13.2)	nr	17.2 (2.2)	15.6 (4.1)
tapentadol ER (Nucynta ER)	2 (1)	17 (7)	17 (6)	1 (0)	15 (13)	21 (7)	1 (<1)	12 (4)	8 (3)
tramadol ER	3.5–6.5	12.2–29.7	15.9–28.2	1– < 5	11.5–15.8 (10.6)	15.1–26.2	nr	7.3–20.3 (1.7)	5–9.4
tramadol ER (Conzip)	3.5–8.6	9.3–21.3	9.6–13.6	nr	19.0–23.1	16.1–25.1	1-5	11.7–16.1	6.5–10.4

Adverse effects are reported as a percentage. Adverse effects data are obtained from package inserts and are not meant to be comparative or all inclusive. Incidences for placebo group are reported in parentheses. nr = not reported.

Opioids have been associated with a decrease in sex hormone levels. Laboratory assessment is recommended in patients who report low libido, impotence, erectile dysfunction, lack of menstruation, or infertility.<sup>183</sup>

## **SPECIAL POPULATIONS**<sup>184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200,201,202,203</sup>

### **Pediatrics**

Fentanyl transdermal (Duragesic) is approved for use in patients as young as 2 years of age who are opioid-tolerant. Oxycodone CR (Oxycontin) is approved for use in opioid-tolerant pediatric patients  $\geq$  11 years of age who are already receiving and tolerating at least 20 mg oxycodone orally (or its equivalent).

Safety and efficacy of buprenorphine transdermal (Butrans), buprenorphine buccal (Belbuca), hydrocodone bitartrate ER (Hysingla ER, Zohydro ER), methadone (Dolophine, Diskets, Methadose), morphine sulfate CR/ER (Arymo ER, Kadian, MS Contin, Morphabond), morphine sulfate ER/naltrexone (Embeda), oxycodone ER (Xtampza ER), oxymorphone ER, tramadol ER (Conzip, generic), and tapentadol ER (Nucynta ER) have not been established in patients younger than 18 years of age. Safety and efficacy of hydromorphone ER (Exalgo) has not been established in patients younger than 17 years of age.

While not approved in this population, the FDA announced in April 2017 that they are requiring additional contraindications and warnings to the labeling of tramadol regarding its use in pediatric patients. This is a result of their investigation of case reports involving respiratory depression.<sup>204</sup>

### **Geriatrics**

Elderly patients may be more sensitive to the opioid agonist effects than younger patients. Monitor geriatric patients closely for signs of sedation and respiratory depression, particularly when initiating opioid therapy and when given in conjunction with other drugs that depress respiration.

Administration of oxymorphone ER in elderly patients resulted in plasma levels that were 40% higher than those in younger subjects.

In clinical trials with buprenorphine buccal (Belbuca), adverse effects were higher in elderly patients than in younger adults.

In clinical trials with oxycodone CR (OxyContin) and oxycodone ER (Xtampza ER), oxycodone exposure was higher in elderly patients than in younger adults, but no unexpected or untoward adverse effects occurred. Use cautiously in this population.

### **Pregnancy**

Prolonged use of opioid analgesics during pregnancy may cause neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated. Opioids cross the placenta and may produce respiratory depression in neonates. Opioids are not for use in women during and immediately prior to labor, when shorter-acting analgesics or other analgesic techniques are more appropriate. Opioid analgesics can prolong labor through actions that temporarily reduce the strength, duration, and frequency of uterine contractions.

Based on the Pregnancy and Lactation Labeling Rule (PLLR), newer agents and agents with labeling changes have had their Pregnancy Category designation replaced with descriptive text. There are no available or adequate data with buprenorphine (Belbuca, Butrans), fentanyl transdermal (Duragesic), hydrocodone ER (Hysingla ER, Zohydro ER), hydromorphone ER (Exalgo), morphine sulfate CR (MS

Contin), morphine sulfate ER (Kadian, Morphabond), morphine sulfate ER/naltrexone (Embeda), oxycodone ER, tapentadol ER (Nucynta ER), and tramadol ER (Conzip) in pregnant women to inform of a drug-associated risk for major birth defects and miscarriage. Previously, most agents within this class were assigned Pregnancy Category C.

Previously, oxycodone CR (Oxycontin) was assigned a Pregnancy Category B designation, but this was also replaced with descriptive information. While animal studies did not reveal evidence of fetal harm, there are no adequate or well-controlled studies in pregnant women; animal studies are not always predictive of human response. Therefore, oxycodone CR should be used during pregnancy only if clearly required. There are no adequate and well-controlled studies of oxycodone ER (Xtampza ER) in pregnant women.

Methadone's Pregnancy Category C designation was also replaced based on the PLLR. There are no adequate and well-controlled studies in pregnant women; it should only be used when the benefit outweighs the potential risk. Women on high-dose methadone maintenance already nursing should be counseled to wean breast-feeding gradually to prevent neonatal abstinence syndrome.

## **Hepatic Impairment**

Oxycodone ER is contraindicated in patients with moderate to severe hepatic impairment.

Avoid use of fentanyl transdermal, tapentadol ER, and tramadol ER in patients with severe hepatic impairment. Reduce dose of tapentadol ER in patients with moderate hepatic impairment.

Initiate hydrocodone ER therapy at 50% of the usual starting dose in patients with severe hepatic impairment and monitor closely for adverse events such as respiratory depression.

Hydromorphone ER should be initiated at 25% of the usual starting dose in patients with moderate hepatic impairment and closely monitor these patients for respiratory and central nervous system depression during initiation and dose titration. The pharmacokinetics of hydromorphone in severe hepatic impairment patients have not been studied.

Oxycodone CR/ER should be initiated at 33% to 50% of the usual starting dose and carefully titrated in patients with hepatic impairment. An alternative analgesic is recommended in those requiring a dose of < 9 mg.

Buprenorphine transdermal (Butrans) has not been evaluated in patients with severe hepatic impairment. Since it is intended for 7-day dosing, alternate analgesic therapy should be considered in patients with severe hepatic impairment.

The pharmacokinetics of morphine in patients with severe hepatic impairment have not been adequately studied; however, there are reports of altered pharmacokinetics in this population. Initiation at a lower dose is generally recommended, followed by cautious titration to monitor for adverse effects.

Buccal buprenorphine (Belbuca) has not been evaluated in patients with severe hepatic impairment; dosage adjustments are not required for patients with mild to moderate hepatic impairment.

## **Renal Impairment**

Avoid use of fentanyl transdermal, tapentadol ER, and tramadol ER in patients with severe renal impairment.

Use oxycodone CR/ER with caution in patients with renal impairment. In renally-impaired patients (estimated creatinine clearance [CrCl] < 60 mL/min), plasma exposure may be 50% higher than in those with normal renal function. An alternative analgesic is recommended in those requiring a dose of < 9 mg.

Titrate oxymorphone ER slowly in patients with moderate to severe renal impairment.

Although accumulation of morphine metabolites has been found in people with renal failure, adequate studies of the pharmacokinetics of morphine in patients with severe renal impairment have not been performed. Initiation at a lower dose is generally recommended, followed by cautious titration to monitor for adverse effects.

Start hydromorphone ER therapy at 50% of the usual starting dose in patients with moderate renal impairment and 25% of the usual starting dose in patients with severe renal impairment. Closely monitor patients for respiratory and central nervous system depression during initiation and during dose titration. Hydromorphone ER is only intended for once daily administration; consider an alternate analgesic that may provide more dosing flexibility in patients with severe renal impairment.

Initiate therapy with 50% of the initial dose of hydrocodone ER in patients with moderate or severe renal impairment or end-stage renal disease and monitor closely for adverse events such as respiratory depression.

No impact of creatinine clearance on steady state buprenorphine concentrations has been found.

No differences in pharmacokinetics of buccal buprenorphine (Belbuca) were found between patients on dialysis and those with normal renal function.

## DOSAGES<sup>205,206,207,208,209,210,211,212,213,214,215,216,217,218,219,220,221,222,223,224</sup>

Drug	Starting Dose	Titration	Available Strengths	Abuse-Deterrent Formulation*
buprenorphine buccal (Belbuca)	75 mcg buccally every 12 to 24 hours, as tolerated	For opioid-naïve patients: 75 mcg buccally every 12 to 24 hours, as tolerated, for $\geq 4$ days prior to increasing to 150 mcg every 12 hours; can proceed in 150 mg increments every 12 hours, no more than every 4 days For opioid-experienced patients, there is a dosing conversion chart in the prescribing information; do not initiate until patient has been tapered to an oral morphine sulfate equivalent of 30 mg or less; Maximum dose: 900 mcg every 12 hours due to risk of QTc prolongation	75 mcg, 150 mcg, 300 mcg, 450 mcg, 600 mcg, 750 mcg, and 900 mcg buccal film	No
buprenorphine transdermal (Butrans)	1 patch changed every 7 days	For opioid-naïve patients: Initiate treatment with buprenorphine transdermal 5 mcg/hour and titrate as needed after 72 hours For opioid-experienced patients, refer to the dosing conversion chart in the prescribing information; do not initiate until patient's around-the-clock opioids have been tapered to an oral morphine equivalent of $\leq 30$ mg for up to days Do not exceed a dose of one 20 mcg/hour patch due to the risk of QTc prolongation	5 mcg/hr, 7.5 mcg/hr, 10 mcg/hr, 15 mcg/hr, and 20 mcg/hr patches	No
fentanyl transdermal (Duragesic)	25 mcg/hr patch changed every 3 days for opioid-tolerant patients	For patients on other opioids converting to fentanyl, calculate their total 24-hour analgesic requirement, then convert this amount to an equivalent analgesic oral morphine dose (see Package Insert for dosing table); however, using the conversion table to convert <i>from</i> fentanyl transdermal to other opioids will result in an unsafe overestimation of the dose Dosage increase may occur 3 days after the first dose and then every 6 days by adding up the rescue medication dosage Initial doses should be reduced in elderly or debilitated patients More than 1 patch may be used for patients requiring more than 100 mcg/hr	12 mcg/hr, 25 mcg/hr, 37.5 mcg/hr (generic only), 50 mcg/hr, 62.5 mcg/hr (generic only), 75 mcg/hr, 87.5 mcg/hr (generic only), and 100 mcg/hr patches	No
hydrocodone ER (Hysingla ER)	20 mg orally every 24 hours	Adjust the dose in increments of 10 mg to 20 mg every 3 to 5 days, as needed, to achieve adequate analgesia	20 mg, 30 mg, 40 mg, 60 mg, 80 mg, 100 mg, and 120 mg tablets	Yes (Formulation deters misuse and abuse via chewing, snorting, or injection)

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

## Dosages (continued)

Drug	Starting Dose	Titration	Available Strengths	Abuse-Deterrent Formulation*
hydrocodone ER (Zohydro ER)	10 mg every 12 hours	Adjust dosage in increments of 10 mg every 12 hours every 3 to 7 days until adequate pain relief with minimal adverse reactions is achieved; refer to the package insert for conversion factors  For opioid-tolerant patients, use the conversion chart to calculate the approximate hydrocodone daily dose and then divide in half for administration every 12 hours; always round down, if necessary, to the appropriate hydrocodone ER strengths available	10 mg, 15 mg, 20 mg, 30 mg, 40 mg, and 50 mg capsules <sup>†</sup>  (Viscous gel develops if crushed and dissolved in liquids or solvents)	No
hydromorphone ER (Exalgo)	8 to 64 mg once daily for opioid-tolerant patients	Adjust dosage not more often than every 3 to 4 days; individually titrate dosing to allow pain control while minimizing adverse reactions	8 mg (generic only), 12 mg, 16 mg, and 32 mg tablets  (Uses Osmotic extended-Release Oral delivery System or OROS, making it difficult to crush or extract for injection)	No
methadone (Dolophine, Diskets, Methadose)	2.5 mg to 10 mg every 8 to 12 hours	Adjust dosage according to the severity of pain and patient response; for exceptionally severe pain, or in those tolerant of opioid analgesia, it may be necessary to exceed the usual recommended dosage	5 mg and 10 mg tablets; 40 mg tablets for oral suspension; 1 mg/mL and 2 mg/mL oral solutions (generic only); 10 mg/mL oral concentrate	No
morphine sulfate CR (MS Contin)	15 mg every 8 to 12 hours	In adjusting dosing regimens, attention should be given to daily dose, degree of opioid tolerance, if any, and general condition and mental status of the patient  Do not crush, chew, break, or dissolve tablets  The 100 mg and 200 mg tablets should only be used in opioid-tolerant patients	15 mg, 30 mg, 60 mg, 100 mg, and 200 mg tablets	No

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

† Reformulated Zohydro ER with abuse-deterrent *properties* is available; originally-marketed formulations are no longer available.

**Dosages (continued)**

Drug	Starting Dose	Titration	Available Strengths	Abuse-Deterrent Formulation*
morphine sulfate ER (Arymo ER)	15 mg every 8 to 12 hours	In adjusting dosing regimens, attention should be given to daily dose, degree of opioid tolerance, if any, and general condition and mental status of the patient  A single dose of > 60 mg or a total daily dose > 120 mg can be used in patients who are tolerant to an opioid of comparable potency  Do not crush, chew, break, or dissolve tablets	15 mg, 30 mg, and 60 mg tablets	Yes  (Polymer matrix tablet utilizing a physical and chemical barrier to deter abuse; increased resistance to cutting, crushing, or grinding the tablet; when in contact with liquid, it becomes a viscous hydrogel to deter from injectable abuse; physical and chemical properties are expected to reduce abuse via the intranasal route)
morphine sulfate ER (Kadian)	1 capsule every 12 to 24 hours based on previous opioid requirements	Titrate to pain control; do not exceed upward titration of more than 20 mg every other day  Swallow capsules whole; do not crush, chew, or dissolve capsules or contents of capsules; may sprinkle pellets on applesauce  The 100 mg, and 200 mg capsules should only be used in opioid-tolerant patients	10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 80 mg, 100 mg, and 200 mg (brand only) capsules <sup>‡</sup>  30 mg, 45 mg, 60 mg, 75 mg, 90 mg, and 120 mg capsules <sup>§</sup>	No
morphine sulfate ER (Morphabond)	15 mg every 12 hours	Titrate to pain control; initiate treatment therapy at 15 mg every 8 to 12 hours for opioid-naïve patients or patients who are not opioid tolerant  For patients who are converting from another oral morphine product, therapy should begin with half of the total established daily morphine dose being given twice daily, or with one-third of the dose given 3 times daily  For patients also taking a CNS depressant, a dose of 15 mg every 12 hours should be used  Do not crush or chew the tablet	15 mg, 30 mg, 60 mg, and 100 mg tablets	Yes  (Uses SentryBond™ technology involving physical and chemical barriers to abuse)

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

‡ The morphine sulfate ER (Kadian) 130 mg and 150 mg formulations have been discontinued.

§ Select strengths are generics for Avinza (morphine sulfate extended-release), which is no longer marketed.

**Dosages (continued)**

Drug	Starting Dose	Titration	Available Strengths	Abuse-Deterrent Formulation*
morphine sulfate ER/naltrexone (Embeda)	1 capsule every 24 hours	<p>Titrate to pain control; initiate therapy at the lowest dose for patients who use Embeda as their first opioid analgesic</p> <p>For patients who are converting from another oral morphine product, therapy should begin with half of the total established daily morphine dose being given twice daily, or with the full dose being given once a day</p> <p>For patients also taking a CNS depressant, a dose of one-third to one-half of recommended starting dose should be used</p> <p>Swallow capsules whole or may sprinkle on applesauce; do not crush, chew, or dissolve capsules or contents of capsules</p> <p>The 100/4 mg capsules are reserved for opioid-tolerant patients</p>	20/0.8 mg, 30/1.2 mg, 50/2 mg, 60/2.4 mg, 80/3.2 mg, and 100/4 mg capsules	<p>Yes</p> <p>(Features a sequestered core of naltrexone to deter crushing, chewing, or dissolving)</p>
oxycodone CR (Oxycontin)	10 mg every 12 hours	<p>Except for the increase from 10 mg to 20 mg every 12 hours, the total daily oxycodone CR dose can be increased by 25 to 50% at each increase; patients should be titrated so that they need no more than 2 supplemental analgesia doses per day; a conversion chart is found in the package insert for patients on other opioid therapy</p> <p>For elderly, debilitated, and patients with hepatic impairment, the dosage should be reduced by 33% to 50%</p> <p>For patients with creatinine clearance &lt; 60 mL/min, dosage may need to be lowered by up to 50%</p> <p>Pediatric dosing is similar to adult dosing</p> <p>Single doses &gt; 40 mg or total dose &gt; 80 mg should only be used in opioid-tolerant patients</p>	10 mg, 15 mg, 20 mg, 30 mg, 40 mg, 60 mg, and 80 mg tablets	<p>Yes</p> <p>(Reformulated to diminish the ease of cutting, breaking, chewing, crushing, or dissolving and to form viscous gel when mixed with aqueous liquid)</p>

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

**Dosages (continued)**

<b>Drug</b>	<b>Starting Dose</b>	<b>Titration</b>	<b>Available Strengths</b>	<b>Abuse-Deterrent Formulation*</b>
oxycodone ER (Xtampza ER)	9 mg orally every 12 hours taken with food (may sprinkle capsule contents onto food or in a cup for administration by mouth, nasogastric tube, or gastrostomy tube)	Adjustments in dosing are recommended to occur no earlier than every 1 to 2 days Single doses of $\geq 36$ mg and total daily doses of $\geq 72$ mg should be reserved for patients whom have shown tolerance for comparable opioid potency analgesia Maximum daily dose is 288 mg (320 mg oxycodone HCl per day)	9 mg, 13.5 mg, 18 mg, 27 mg, and 36 mg capsules	Yes (Employs Deterx <sup>®</sup> technology: a microsphere-in-capsule formulation; in each microsphere, oxycodone is present as a solid solution of a fatty acid salt [oxycodone myristate] in a hydrophobic matrix that also contains waxes) <sup>225</sup>
oxymorphone ER	5 mg every 12 hours	Increase by 5 to 10 mg twice a day every 3 to 7 days based on patient pain intensity and adverse effects Do not attempt to break, crush, chew, or dissolve tablets In patients with creatinine clearance $< 50$ mL/min, oxymorphone should be started with the lowest dose and titrated slowly while carefully monitoring adverse effects	5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, and 40 mg tablets	No
tapentadol ER (Nucynta ER)	50 mg every 12 hours	Titrate to pain control within therapeutic range of 100 to 250 mg twice a day; in patients previously taking other opioid therapy, initiate with 50 mg then titrate to effective and tolerable dose within range of 100 to 250 mg twice a day; titrate dose by no more than 50 mg/dose twice a day (100 mg/day) no more than every 3 days Do not exceed maximum daily dose of 500 mg (250 mg twice a day)	50 mg, 100 mg, 150 mg, 200 mg, and 250 mg tablets	No (Resistant to breaking or crushing)

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

### Dosages (continued)

Drug	Starting Dose	Titration	Available Strengths	Abuse-Deterrent Formulation*
tramadol ER (Conzip)	100 mg daily	Initiate at 100 mg daily then titrate at 100 mg increments every 5 days as needed to relieve pain, up to max 300 mg/day; dose no more frequently than every 24 hours For patients maintained on tramadol immediate-release tablets, calculate the 24-hour tramadol dose, and initiate a total daily dose of the extended-release capsules rounded down to the next lowest 100 mg increment; individualize dose as needed, up to a maximum of 300 mg/day Consider lower doses in elderly; The concomitant use of the extended-release capsules with other tramadol products is not recommended	100 mg, 150 mg (authorized generic only), 200 mg, and 300 mg capsules	No
tramadol ER	1 tablet daily	Initiate at 100 mg daily then titrate at 100 mg increments every 5 days as needed to relieve pain, up to a max of 300 mg/day; do not use in patients with severe renal or hepatic impairment	100 mg, 200 mg, and 300 mg tablets (generics with reference product of Ultram ER)  100 mg, 200 mg, and 300 mg tablets (generics with reference product of Ryzolt)	No

\* Has met requirements by the FDA to be approved as an abuse-deterrent formulation.

The following medications are available as abuse-deterrent formulations: morphine sulfate tablets (Arymo ER, Morphabond), morphine sulfate ER/naltrexone (Embeda) capsules; hydrocodone ER (Hysingla ER) tablets; oxycodone CR (Oxycontin) tablets; oxycodone ER (authorized generics for OxyContin) tablets; and oxycodone ER capsules (Xtampza ER). Although hydromorphone ER (Exalgo), tapentadol ER (Nucynta ER), and hydrocodone ER (Zohydro ER) have abuse-deterrent properties, they have not been approved by the FDA as abuse-deterrent formulations.

The naltrexone component of morphine sulfate ER/naltrexone (Embeda) is formulated such that, if the capsule is swallowed whole or opened and sprinkled over applesauce, the morphine component will be released while the naltrexone will remain sequestered in a film coating that is resistant to digestion. However, if the capsule contents are chewed or crushed, the naltrexone is released, reversing the effects of the morphine, thus reducing the likelihood that the product will be abused by disabling the extended-release mechanism. No studies have established whether this hypothesized reduction of morphine effects following chewing or crushing were clinically significant; the product labeling states that there is no evidence that the naltrexone in Embeda reduces the abuse liability.

The tapentadol ER tablet formulation is designed to provide a high degree of mechanical resistance to crushing or chewing.

Due to significant pharmacokinetic variability among patients, close observation and slow titration of methadone (Dolophine, Diskets, Methadose) is required. The peak respiratory depressant effect lasts longer than the peak therapeutic effect; extra caution when dosing is recommended.

Patients receiving other oxycodone products may convert on the basis of administering one-half of the current daily dose every 12 hours with oxycodone ER (Xtampza ER). For other opioids, see prescribing information. A 9 mg dose of Xtampza ER is equivalent to a 10 mg dose of other oxycodone HCl products (ratio of 9:10 for other doses as well).

Oxymorphone ER should be given on an empty stomach at least 1 hour prior to or 2 hours after eating. Maximum serum concentration was increased by 50% when given with food. An *in vivo* study with oxymorphone ER showed that the maximum concentration increased 31% to 70%, on average, following concomitant administration with ethanol. Co-administration must be avoided.

Apply buprenorphine buccal film to the inside of the cheek. It will dissolve in approximately 30 minutes. Individual titration should be in increments of no more than 150 mcg every 12 hours and should not occur any sooner than every 4 days. Reduce dose (and titration increments) in half for patients with severe hepatic impairment or mucositis.

Apply buprenorphine transdermal patch (Butrans) to the upper outer arm, upper chest, upper back, or the side of the chest. Rotate application sites, waiting a minimum of 21 days before reapplying to the same skin site. Apply to a hairless or nearly hairless, dry skin site. Do not apply to irritated skin. Do not use soaps, lotions, oils, or alcohol on the skin before the patch is applied. If the buprenorphine transdermal patch falls off during the 7-day dosing interval, dispose of the transdermal system properly and place a new patch on at a different skin site. During the dose titration, 2 patches can be used for a total max dose of 20 mcg.

Fentanyl transdermal patch (Duragesic) patch should be applied to the chest, back, flank, or upper arm on dry, intact, hairless skin. Do not use soaps, lotions, oils, or alcohol on the skin before the patch is applied. If the patch falls off before 3 days of use, discard and apply a new patch at a different skin site. In an effort to minimize the risk of accidental exposure to fentanyl patches, the FDA is requiring the manufacturer of Duragesic to print the name and strength of the drug on the patch in long-lasting ink, in a color that is clearly visible to patients and caregivers.<sup>226</sup> The current ink color varies by strength and is not always easy to see. This change is intended to enable patients and caregivers to more easily find patches on patients' bodies and see patches that have fallen off, which children or pets could accidentally touch or ingest. The manufacturers of generic fentanyl patches are being requested to make similar changes.

Patients using morphine sulfate ER (Morphabond ER) may notice a tablet shell ("ghost" tablet) in stool.

Opioids should be stored securely, out of sight and reach of children, in a location not accessible to other individuals. Once the medication is expired or no longer being used it should be disposed of properly through a drug take-back program, if available or in another appropriate manner as determined by the FDA: [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal).<sup>227</sup>

## Opioid Morphine Equivalent Conversions<sup>228</sup>

This table is intended to provide an estimate of overall opioid exposure; it should not be used for dosing determinations (e.g., converting a patient from 1 opioid to another). Conversion factors may vary based on individual pharmacokinetics and duration of use (e.g., opioid-naïve versus chronic dosing). The same conversion is used for immediate- and extended-release oral products with the same opioid component unless otherwise specified. This table includes medications that are not reviewed in this class review for reference purposes. Likewise, some medications are not included in this table due to limited data. See prescribing information for detailed recommendations, when available, for converting among formulations and active ingredients. While previously assigned a MME conversion factor, the CDC has since removed buprenorphine's MME assignment and nalbuphine has been removed from the list.

Opioid	MME Conversion Factor
butorphanol	7
codeine	0.15
dihydrocodeine	0.25
fentanyl buccal, SL tablet, or lozenge*	0.13
fentanyl film or oral spray*	0.18
fentanyl nasal spray*	0.16
fentanyl patch†	7.2
hydrocodone	1
hydromorphone	4
levorphanol tartrate	11
mepiridine	0.1
methadone	3
morphine‡	1
opium	1
oxycodone hydrochloride <sup>§</sup>	1.5
oxymorphone	3
pentazocine	0.37
tapentadol	0.4
tramadol	0.1

\* Multiply conversion factor by the number of micrograms in the dose.

† Based on total micrograms exposure over 24 hours and assumes 1 mg parenteral fentanyl = 100 mg oral morphine (e.g., 25 mcg/hr patch = 180 MME over 3 days = 60 MME/day).

‡ Extended-release formulations of morphine sulfate are not bioequivalent. A ratio of approximately 3 mg oral to 1 mg parenteral morphine sulfate may be used to estimate exposure, but not for dosing conversions as a more cautious approach is recommended.

§ Strengths of oxycodone ER capsules (Xtampza ER) should be converted to the oxycodone HCl equivalent prior to calculation using a ratio of 9 mg Xtampza ER to 10 mg oxycodone HCl.

## CLINICAL TRIALS

### Search Strategies

Articles were identified through searches performed on PubMed and review of information sent by manufacturers. Search strategy included the FDA-approved use of all drugs in this class. Randomized, controlled, comparative trials for FDA-approved indications are considered the most relevant in this category. Studies included for analysis in the review were published in English, performed with human participants, and randomly allocated participants to comparison groups. In addition, studies must contain clearly stated, predetermined outcome measure(s) of known or probable clinical importance, use data analysis techniques consistent with the study question, and include follow-up (endpoint assessment) of at least 80% of participants entering the investigation. Despite some inherent bias found in all studies including those sponsored and/or funded by pharmaceutical manufacturers, the studies in this therapeutic class review were determined to have results or conclusions that do not suggest systematic error in their experimental study design. While the potential influence of manufacturer sponsorship/funding must be considered, the studies in this review have also been evaluated for validity and importance.

There are no published comparative clinical studies with the following agents: buprenorphine (Belbuca), buprenorphine (Butrans), hydrocodone ER (Hysingla ER, Zohydro ER), hydromorphone ER (Exalgo), morphine sulfate ER (Arymo ER, Morphabond), morphine sulfate ER/naltrexone (Embeda), oxycodone ER (Xtampza ER), tapentadol ER (Nucynta ER), and tramadol ER (generic, Conzip).<sup>229,230,231,232,233,234, 235, 236,237,238,239,,240,241,242,243,244</sup>

The following agents have demonstrated abuse-deterrence in placebo-controlled and/or active-comparator studies, which is commonly measured by a measure of drug-liking using a visual analog scale (VAS), although other measurements have also been used: Arymo ER tablets, Embeda capsules, Hysingla ER tablets, Morphabond tablets, Oxycontin biconcave tablets, oxycodone ER (authorized generics of OxyContin) tablets, and Xtampza ER capsules.

### methadone (Dolophine) versus morphine sulfate SR

A total of 103 patients with pain requiring initiation of strong opioids were randomly assigned to treatment with methadone 7.5 mg every 12 hours and 5 mg every 4 hours, as needed, or morphine 15 mg sustained release every 12 hours and 5 mg every 4 hours, as needed.<sup>245</sup> After 4 weeks, patients receiving methadone had more opioid-related discontinuations than those receiving morphine (22 versus 6%;  $p=0.019$ ). The opioid escalation index at days 14 and 28 were similar between the 2 groups. More than three-fourths of patients in each group reported a 20% or more reduction in pain intensity by day 8; at 4 weeks, the proportion of patients with a 20% or more reduction in pain was similar: 0.49 in the methadone group and 0.56 in the morphine group.

### oxycodone controlled-release (Oxycontin) versus oxycodone immediate-release

A multicenter, randomized, double-blind, parallel-group study was performed in 111 patients with cancer pain.<sup>246</sup> Patients were being treated with fixed-combination opioid/nonopioid analgesics at study entry. Patients received oxycodone CR 30 mg every 12 hours or oxycodone IR 15 mg 4 times daily for 5 days. No titration or supplemental analgesic medications were permitted. The mean baseline pain intensity (0 = none, 1 = slight, 2 = moderate, 3 = severe) was 1.5 for the oxycodone CR-treated group and

1.3 for the oxycodone IR group ( $p>0.05$ ). The 5-day mean pain intensity was 1.4 and 1.1 for the CR and IR groups, respectively ( $p>0.05$ ). Discontinuation rates were equivalent (33%). There was no significant difference between treatment groups in the incidence of adverse events.

Cancer patients who required therapy for moderate to severe pain were randomized to oxycodone CR every 12 hours ( $n = 81$ ) or oxycodone IR 4 times daily ( $n = 83$ ) for 5 days in a multicenter, double-blind study.<sup>247</sup> Rescue medication was allowed. Pain intensity remained slight during the study, with mean oxycodone CR doses of 114 mg/day and mean oxycodone IR doses of 127 mg/day. Acceptability of therapy was fair to good with both treatments. Discontinuation rates for lack of acceptable pain control were 4% with oxycodone CR and 19% with oxycodone IR. Fewer adverse events were reported with oxycodone CR than with oxycodone IR ( $p=0.006$ ).

### **oxymorphone ER versus oxycodone CR (Oxycontin)**

A multicenter, randomized, double-blind, placebo- and active-controlled trial was conducted to compare the analgesic efficacy and safety of oxymorphone ER with placebo and oxycodone CR in patients with moderate to severe chronic low back pain requiring opioid therapy.<sup>248</sup> Patients ( $n=213$ ) were randomized to receive oxymorphone ER (10 to 110 mg) or oxycodone CR (20 to 220 mg) every 12 hours during a 7- to 14-day dose-titration phase. Patients achieving effective analgesia at a stable opioid dose entered an 18-day double-blind treatment phase and either continued opioid therapy or received placebo. With stable dosing throughout the treatment phase, oxymorphone ER (79.4 mg/day) and oxycodone CR (155 mg/day) were superior to placebo for change from baseline in pain intensity as measured on a visual analog scale ( $p=0.0001$ ). Adverse events for the active drugs were similar; the most frequently reported were constipation and sedation. Oxymorphone ER was equianalgesic to oxycodone CR at one-half the milligram daily dosage with comparable safety.

### **tapentadol ER (Nucynta ER) versus oxycodone CR (Oxycontin)**

A randomized, double-blind study was conducted to compare the safety and efficacy of tapentadol ER in patients with moderate-to-severe chronic osteoarthritis knee pain.<sup>249</sup> A total of 990 patients with pain requiring initiation of strong opioids were randomly assigned to tapentadol ER, oxycodone CR, or placebo for a double-blind, 3-week titration period and a 12-week maintenance period. After 15 weeks, the primary endpoints were not significantly different for patients receiving tapentadol ER or for oxycodone CR versus placebo at week 12 ( $p=0.152$ ). Significantly more patients rated their pain to be “much improved” with tapentadol ER at the end of treatment (56% versus 42.5%, respectively). Tapentadol ER was associated with better gastrointestinal tolerability with constipation, nausea, and/or vomiting ( $p<0.001$ ). However, the study did not demonstrate assay sensitivity or significant pain reduction.

## **META-ANALYSES**

A meta-analysis compared tapentadol ER and oxycodone/naloxone ER (not available) to oxycodone CR (3 trials of tapentadol ER versus oxycodone CR with or without a placebo group;  $n=3,105$ ).<sup>250</sup> Compared to oxycodone CR, tapentadol ER resulted in a better risk ratio reduction for discontinuation due to adverse effects (RR, 0.526 [95% CI, 0.456 to 0.607]), discontinuation due to nausea and vomiting (RR, 0.526 [95% CI, 0.471 to 0.588]), and constipation (RR, 0.609 [95% CI, 0.545 to 0.68]).

A Cochrane review of opioids (immediate and extended release formulations) for chronic low back pain (15 trials; n=5,540), found that tramadol was more effective for pain control than placebo (standardized mean difference [SMD], -0.55 [95% CI, -0.66 to -0.44]; 5 trials; n=1,378).<sup>251</sup> Transdermal buprenorphine had a small, but significant impact on pain control (SMD, -2.47 [95% CI, -2.69 to -2.25]; 2 trials; n=653), and stronger opioids (morphine, hydromorphone, oxycodone, oxymorphone, and tapentadol) were also superior to placebo for pain control (SMD, -0.43 [95% CI, -0.52 to -0.33]; 6 trials; n=1,887).

A systematic review compared transdermal buprenorphine to transdermal fentanyl using 17 clinical trials, including trials for indirect comparison.<sup>252</sup> Transdermal fentanyl was associated with a greater risk of nausea (odds ratio [OR], 4.66 [95% CI, 1.07 to 20.39]) and treatment discontinuations due to adverse effects compared to transdermal buprenorphine (OR, 5.94 [95% CI, 1.78 to 19.87]). No differences were found in efficacy.

## SUMMARY

Pain of multiple etiologies remains a substantial problem for many patients presenting in the clinical setting. No clinical data exist that distinguish analgesic efficacy of any of these products from the others. Pain management must be individualized, and patients who do not respond to one opioid may respond to another.

Full opioid agonists do not typically have a ceiling for their analgesic effectiveness but the dose is limited by their adverse effects. Morphine is the standard drug of comparison. Methadone (Dolophine, Diskets, Methadose) may provide an effective alternative in palliative care of most patients with cancer pain referred for poor pain control and/or adverse effects. It is also useful in the treatment of opioid dependence. Morphine sulfate has been available as a twice-daily sustained-release oral dosage form (MS Contin) for many years, and 2 newly-approved formulations, Arymo ER and Morphabond, are approved as abuse-deterrent formulations. A morphine extended-release dosage form (Kadian) is available for once-daily use. Hydrocodone bitartrate extended-release (Hysingla ER, Zohydro ER) is available without an acetaminophen component. Both Hysingla ER and a new formulation of Zohydro ER contain properties to limit abuse, but only Hysingla ER has met FDA abuse-deterrent formulation requirements. Hysingla ER is dosed once every 24 hours, while Zohydro ER is taken every 12 hours. Hydromorphone ER (Exalgo) is available for a once-daily dosing. Morphine ER/naltrexone (Embeda) is a formulation of morphine that theoretically reduces abuse potential with the addition of naltrexone and dosed once daily.

Tapentadol ER (Nucynta ER) and tramadol ER (Conzip) are centrally-acting oral analgesics that bind to mu-opioid receptors and inhibit norepinephrine re-uptake. The tapentadol ER tablet formulation is designed to provide a high degree of mechanical resistance to crushing or chewing. Tapentadol ER is dosed twice daily, while tramadol ER is dosed once daily.

Like the controlled-release forms of morphine, oxycodone CR (Oxycontin) allows for less frequent (12-hour) dosing of an opioid. Oxycodone CR has a significant potential for abuse and has been associated with increases in crime and deaths due to illicit use. However, all opioids can be abused and are subject to illicit use. Oxycodone CR has been reformulated to include a delivery system that causes a gummy substance when tablets are crushed; the effects of this redesign on illicit use have yet to be seen. A new formulation of oxycodone ER, Xtampza ER capsules, offers another twice-daily treatment option as an

abuse-deterrent formulation. These capsules may also be opened and administered in alternative ways for patients unable to swallow them whole.

Oxymorphone hydrochloride, an opioid agonist, is a metabolite of oxycodone, which is taken twice daily.

The following medications are available as abuse-deterrent formulations: Arymo ER tablets, Embeda capsules, Hysingla ER tablets, Morphabond tablets, Oxycontin tablets, oxycodone ER (authorized generics of Oxycontin) tablets, and Xtampza ER capsules. In 2018, however, the manufacturer of Arymo ER announced that they are discontinuing its promotion and manufacture, and in 2019, the manufacturer of Embeda announced the product would be discontinued.

Two agents in this review provide analgesia via transdermal routes. Buprenorphine (Butrans), a partial opioid agonist, and fentanyl (Duragesic), a full agonist, are available as transdermal patches. Buprenorphine transdermal patch should be changed every 7 days, while a new fentanyl transdermal patch is recommended every 72 hours.

One agent provides a buccal route for analgesia, buprenorphine buccal (Belbuca), which is dosed twice daily.

## REFERENCES

- 1 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 2 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 3 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.
- 4 Hysingla ER [package insert]. Stamford, CT, Purdue; October 2019.
- 5 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 6 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.
- 7 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 8 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.
- 9 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 10 Kadian [package insert]. Madison, NJ; Allergan; October 2019.
- 11 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.
- 12 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 13 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 14 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 15 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
- 16 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 17 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 18 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 19 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019 .
- 20 United States Securities and Exchange Commission. 001-36295. [https://www.sec.gov/Archives/edgar/data/1586105/000110465918059435/a18-36089\\_18k.htm](https://www.sec.gov/Archives/edgar/data/1586105/000110465918059435/a18-36089_18k.htm). Accessed March 28, 2020.
- 21 Morphine sulfate and naltrexone hydrochloride (Embeda) extended-release capsules. Available at: [https://www.accessdata.fda.gov/scripts/drugshortages/dsp\\_ActiveIngredientDetails.cfm?AI=Morphine%20Sulfate%20and%20Naltrexone%20Hydrochloride%20\(EMBEDA%20AE\)%20Extended-Release%20Capsules%20&st=d](https://www.accessdata.fda.gov/scripts/drugshortages/dsp_ActiveIngredientDetails.cfm?AI=Morphine%20Sulfate%20and%20Naltrexone%20Hydrochloride%20(EMBEDA%20AE)%20Extended-Release%20Capsules%20&st=d). Accessed March 28, 2020.
- 22 FDA news release. FDA requests removal of Opana ER for risks related to abuse. Available at: <https://www.fda.gov/newsevents/newsroom/pressannouncements/ucm562401.htm>. Accessed March 28, 2020.
- 23 Endo Press Release. July 6, 2017. Endo Provides Update On Opana ER. Available at: [http://investor.endo.com/newsroom/news-releases?field\\_nir\\_news\\_date\\_value%5Bmin%5D=2017#views-exposed-form-widget-news-widget-news-ul](http://investor.endo.com/newsroom/news-releases?field_nir_news_date_value%5Bmin%5D=2017#views-exposed-form-widget-news-widget-news-ul). Accessed March 28, 2020.
- 24 Barkin RL. Acetaminophen, aspirin, or Ibuprofen in combination analgesic products. *Am J Ther.* 2001; 8(6):433–442.
- 25 Marks RM, Sacher EJ. Undertreatment of medical inpatient pain with narcotic analgesics. *Ann Intern Med.* 1973; 78:173–181.
- 26 Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain – United States, 2016. *MMWR Recomm. Rep.* 2016; 65(1): 1-49. DOI: 10.15585/mmwr.rr6501e1. Available at: <http://www.cdc.gov/mmwr/volumes/65/rr/rr6501e1.htm>. Accessed March 28, 2020.
- 27 Lister BJ. Dilemmas in the treatment of chronic pain. *Am J Med.* 1996; 101(suppl 1A):2S-5S.
- 28 Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain – United States, 2016. *MMWR Recomm Rep.* 2016; 65(1): 1-49. DOI: 10.15585/mmwr.rr6501e1. Available at: <http://www.cdc.gov/mmwr/volumes/65/rr/rr6501e1.htm>. Accessed March 28, 2020.
- 29 CDC Drug Overdose Deaths. Available at: <https://www.cdc.gov/drugoverdose/data/statedeaths.html>. Accessed March 28, 2020.

- 30 Centers for Disease Control and Prevention. Annual Surveillance Report of Drug-Related Risks and Outcomes — United States, 2017. Surveillance Special Report 1. Centers for Disease Control and Prevention, U.S. Department of Health and Human Services. Published August 31, 2017. Available at: <https://www.cdc.gov/drugoverdose/pdf/pubs/2017-cdc-drug-surveillance-report.pdf>. Accessed March 28, 2020.
- 31 Centers for Disease Control and Prevention. 2019 Annual Surveillance Report of Drug-Related Risks and Outcomes — United States, 2019. Surveillance Special Report. Centers for Disease Control and Prevention, U.S. Department of Health and Human Services. Published November 1, 2019. Available at: <https://www.cdc.gov/drugoverdose/pdf/pubs/2019-cdc-drug-surveillance-report.pdf>. Accessed March 28, 2020.
- 32 Centers for Disease Control and Prevention. Annual Surveillance Report of Drug-Related Risks and Outcomes — United States, 2019. Surveillance Special Report. Centers for Disease Control and Prevention, U.S. Department of Health and Human Services. Published November 1, 2019. Available at: <https://www.cdc.gov/drugoverdose/pdf/pubs/2019-cdc-drug-surveillance-report.pdf>. Accessed March 28, 2020.
- 33 Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain — United States, 2016. MMWR Recomm Rep. 2016; 65(1): 1-49. DOI: 10.15585/mmwr.rr6501e1. Available at: <http://www.cdc.gov/mmwr/volumes/65/rr/rr6501e1.htm>. Accessed March 28, 2020.
- 34 WHO's cancer pain ladder for adults. Available at: <http://www.who.int/cancer/palliative/painladder/en/>. Accessed March 28, 2020.
- 35 WHO guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents. Available at: <https://www.who.int/ncds/management/palliative-care/cancer-pain-guidelines/en/>. Accessed March 29, 2020.
- 36 Available at: <https://www.who.int/cancer/palliative/painladder/en/>. Accessed March 29, 2020.
- 37 Paice JA, Protenov R, Lacchetti C, et al. Management of chronic pain in survivors of adult cancers: American Society of Clinical Oncology clinical practice guidelines. J Clin Oncol. 2016; 34(27): 3325-3345. DOI: 10.1200/JCO.2016.68.5206. Available at: <https://www.asco.org/research-guidelines/quality-guidelines/guidelines/patient-and-survivor-care/#/13021>. Accessed March 29, 2020.
- 38 Swarm RA, Paice JA, Angheliescu DL, et al. National Comprehensive Cancer Network (NCCN) Adult Cancer Pain V3.2019. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/pain.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf). Accessed March 29, 2020.
- 39 Swarm RA, Paice JA, Angheliescu DL, et al. National Comprehensive Cancer Network (NCCN) Adult Cancer Pain V3.2019. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/pain.pdf](https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf). Accessed March 29, 2020.
- 40 Goadsby P, Goldenberg JN, Hinson H, et al. Position Statement: Opioids. American Academy of Neurology 2018. Available at: <https://www.aan.com/policy-and-guidelines/policy/position-statements/>. Accessed March 29, 2020.
- 41 Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain — United States, 2016. MMWR Recomm Rep. 2016; 65(1): 1-49. DOI: 10.15585/mmwr.rr6501e1. Available at: <http://www.cdc.gov/mmwr/volumes/65/rr/rr6501e1.htm>. Accessed March 28, 2020.
- 42 Dowell D, Haegerich TM, Chou R. CDC guideline for prescribing opioids for chronic pain — United States, 2016. MMWR Recomm Rep. 2016; 65(1): 1-49. DOI: 10.15585/mmwr.rr6501e1. Available at: <http://www.cdc.gov/mmwr/volumes/65/rr/rr6501e1.htm>. Accessed March 28, 2020.
- 43 Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. Ann Intern Med. 2017; 166(7): 514-530. DOI: 10.7326/M16-2367. Available at: <https://www.acponline.org/clinical-information/guidelines>. Accessed March 29, 2020.
- 44 Manchikanti L, Kaye AM, Knezevic NN, et al. Responsible, safe, and effective prescription of opioids for chronic non-cancer pain: American Society of Interventional Pain Physicians (ASIPP) guidelines. Pain Physician. 2017; 20(2S): S3-S92. Available at: <https://www.asipp.org/opioidguidelines.htm>. Accessed March 29, 2020.
- 45 Bruce RD, Merlin J, Lum PJ, et al. 2017 HIVMA of IDSA Clinical practice guideline for the management of chronic pain in patients living With HIV. Clinical Infectious Diseases. 2017; 65(10);e1–e37. Available at: <http://www.idsociety.org/PracticeGuidelines/>. Accessed March 29, 2020.
- 46 American Association of Oral and Maxillofacial Surgeons. White Paper. Opioid prescribing: Acute and postoperative pain management. Available at: <https://www.aaoms.org/practice-resources/aaoms-advocacy-and-position-statements/white-papers>. Accessed March 29, 2020.
- 47 Institute for Clinical and Economic Review. Abuse-deterrent formulations of opioids: Effectiveness and value. Final Evidence Report. August 8, 2017. Available at: <https://icer-review.org/topics/#past-topics>. Accessed March 29, 2020.
- 48 Silber NH, Becker PM, Buchfuhrer MJ, et al. The Appropriate use of opioids in the treatment of refractory restless legs syndrome. Mayo Clinic, 2018 93(1):59-67. DOI: 10.1016/j.mayocp.2017.11.007. Available at: [https://www.mayoclinicproceedings.org/article/S0025-6196\(17\)30825-X/fulltext](https://www.mayoclinicproceedings.org/article/S0025-6196(17)30825-X/fulltext). Accessed March 29, 2020.
- 49 American Academy of Pain Medicine. North American Spine Society. Evidence-based clinical guidelines for multidisciplinary spine care: Diagnosis and treatment of low back pain. 2020. Available at: <https://painmed.org/clinician-resources/clinical-guidelines>. Accessed March 30, 2020.
- 50 How healthcare providers and patients can better utilize naloxone, the opioid reversal drug. Available at: <https://www.hhs.gov/opioids/prevention/safe-opioid-prescribing/index.html>. Accessed August March 29, 2020.
- 51 Guidance for clinicians on the appropriate dosage reduction or discontinuation of long-term opioids. Available at: <https://www.hhs.gov/opioids/treatment/clinicians-guide-opioid-dosage-reduction/>. Accessed February 21, 2020.
- 52 Drug Enforcement Administration. Schedules of controlled substances: Placement of tramadol Into Schedule IV. Available at: [http://www.deadiversion.usdoj.gov/fed\\_regs/rules/2014/fr0702.htm](http://www.deadiversion.usdoj.gov/fed_regs/rules/2014/fr0702.htm). Accessed March 29, 2020.
- 53 FDA announces safety labeling changes and postmarket study requirements for extended-release and long-acting opioid analgesics. September 10, 2013. Available at: <https://www.fda.gov/Drugs/DrugSafety/InformationbyDrugClass/ucm363722.htm>. Accessed March 29, 2020.
- 54 FDA Guidance Document. Abuse-deterrent opioids-evaluation and labeling. April 2015. Available at: <https://www.fda.gov/downloads/drugs/guidancecomplianceregulatoryinformation/guidances/ucm334743.pdf>. Accessed March 29, 2020. .
- 55 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 56 Hysingla ER [package insert]. Stamford, CT, Purdue; October 2019.
- 57 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 58 FDA. General principles for evaluating the abuse deterrence of generic solid oral opioid drug products guidance for industry. November 2017. Available at: <https://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM492172.pdf>. Accessed March 29, 2020.
- 59 FDA news release. Calif, FDA top officials call for sweeping review of agency opioids policies. Available at: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm484765.htm>. Accessed March 29, 2020.

- 
- 60 FDA Drug Safety Communication: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. April 20, 2017. Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm549679.htm>. Accessed March 29, 2020.
- 61 H.R.6 - SUPPORT for Patients and Communities Act. October 24, 2018. Available at: <https://www.congress.gov/bill/115th-congress/house-bill/6>. Accessed March 29, 2020.
- 62 DEA announces enhanced tool for registered drug manufacturers and distributors to combat opioid crisis. Available at: <https://www.dea.gov/press-releases/2019/02/26/dea-announces-enhanced-tool-registered-drug-manufacturers-and>. Accessed March 29, 2020.
- 63 FDA news release: FDA launches public education campaign to encourage safe removal of unused opioid pain medicines from homes. April 25, 2019. Available at: <https://www.fda.gov/news-events/press-announcements/fda-launches-public-education-campaign-encourage-safe-removal-unused-opioid-pain-medicines-homes>. Accessed March 29, 2020.
- 64 FDA Drug Safety Communication: FDA identifies harm reported from sudden discontinuation of opioid pain medicines and requires label changes to guide prescribers on gradual, individualized tapering. April 9, 2019. Available at: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-identifies-harm-reported-sudden-discontinuation-opioid-pain-medicines-and-requires-label-changes>. Accessed March 29, 2020.
- 65 FDA news release: FDA launches public education campaign to encourage safe removal of unused opioid pain medicines from homes. Available at: <https://www.fda.gov/news-events/press-announcements/fda-launches-public-education-campaign-encourage-safe-removal-unused-opioid-pain-medicines-homes>. Accessed March 29, 2020.
- 66 Timeline of selected FDA activities and significant events addressing opioid misuse and abuse. Available at: <https://www.fda.gov/Drugs/DrugSafety/InformationbyDrugClass/ucm338566.htm>. Accessed March 29, 2020.
- 67 FDA Statement. Statement on National Academies of Sciences, Engineering, and Medicine report on framing opioid prescribing guidelines for acute pain. Available at: <https://www.fda.gov/news-events/press-announcements/statement-national-academies-sciences-engineering-and-medicine-report-framing-opioid-prescribing>. Accessed March 29, 2020.
- 68 DEA. DEA reduces amount of opioid controlled substances to be manufactured in 2017. October 4, 2016. Available at: <https://www.dea.gov/divisions/hq/2016/hq100416.shtml>. Accessed March 28, 2020.
- 69 DEA. DEA proposes to reduce the amount of five opioids manufactured in 2020, marijuana quota for research increases by almost a third. September 11, 2019. Available at: <https://www.dea.gov/press-releases/2019/09/11/dea-proposes-reduce-amount-five-opioids-manufactured-2020-marijuana-quota>. Accessed March 28, 2020.
- 70 Established aggregate production quotas for Schedule I and II controlled substances and assessment of annual needs for the list I chemicals ephedrine, pseudoephedrine, and phenylpropanolamine for 2020. Available at: [https://www.deadiversion.usdoj.gov/fed\\_regs/quotas/2019/fr1202.htm](https://www.deadiversion.usdoj.gov/fed_regs/quotas/2019/fr1202.htm). Accessed March 28, 2020.
- 71 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 72 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 73 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.
- 74 Hysingla ER [package insert]. Stamford, CT; Purdue; October 2019.
- 75 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 76 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.
- 77 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 78 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.
- 79 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 80 Kadian [package insert]. Madison, NJ; Allergan; October 2019.
- 81 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.
- 82 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 83 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 84 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 85 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
- 86 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 87 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 88 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 89 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.
- 90 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.
- 91 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 92 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 93 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.
- 94 Hysingla ER [package insert]. Stamford, CT; Purdue; October 2019.
- 95 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 96 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.
- 97 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 98 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.
- 99 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 100 Kadian [package insert]. Madison, NJ; Allergan; October 2019.
- 101 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.
- 102 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 103 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 104 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 105 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
-

- 
- 106 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 107 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 108 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 109 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.
- 110 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.
- 111 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 112 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 113 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.
- 114 Hysingla ER [package insert]. Stamford, CT; Purdue; October 2019.
- 115 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 116 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.
- 117 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 118 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.
- 119 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 120 Kadian [package insert]. Madison, NJ; Allergan; October 2019.
- 121 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.
- 122 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 123 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 124 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 125 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
- 126 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 127 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 128 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 129 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.
- 130 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.
- 131 FDA Drug Safety Communication: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. April 20, 2017. Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm549679.htm>. Accessed March 29, 2020.
- 132 FDA Drug Safety Communication. FDA warns about serious risks and death when combining opioid pain or cough medicines with benzodiazepines; requires its strongest warning. August 31, 2016. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm518473.htm>. Accessed March 29, 2020.
- 133 FDA Drug Safety Communication. FDA warns about several safety issues with opioid pain medicines; requires label changes. March 22, 2016. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm489676.htm>. Accessed March 29, 2020.
- 134 FDA Drug Safety Communication. FDA warns about several safety issues with opioid pain medicines; requires label changes. March 22, 2016. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm489676.htm>. Accessed March 29, 2020.
- 135 Fentanyl Transdermal System (marketed as Duragesic and generics). December 21, 2007. Available at: <https://wayback.archive-it.org/7993/20170112170250/http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm152289.htm>. Accessed March 29, 2020.
- 136 FDA News Release. FDA approves new formulation of OxyContin. April 5, 2010. Available at: <https://wayback.archive-it.org/7993/20170112130258/http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm207480.htm>. Accessed August 13, 2019.
- 137 Abusing pain drug Opana can cause blood disorder: FDA . Available at: <https://www.reuters.com/article/us-opana-warning/abusing-pain-drug-opana-can-cause-blood-disorder-fda-idUSBRE89A13W20121011>. Accessed March 29, 2020.
- 138 FDA news release. FDA requests removal of Opana ER for risks related to abuse. June 8, 2017. Available at: <https://www.fda.gov/newsevents/newsroom/pressannouncements/ucm562401.htm>. Accessed March 29, 2020.
- 139 Endo Press Release. Endo Provides Update On Opana ER. July 6, 2017. Available at: <http://investor.endo.com/news-releases/news-release-details/endo-provides-update-opanar-er>. Accessed March 29, 2020.
- 140 FDA Drug Safety Communication. FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. April 20, 2017. Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm549679.htm>. Accessed March 29, 2020.
- 141 REMS@FDA. Available at: <http://www.accessdata.fda.gov/scripts/cder/rems/index.cfm>. Accessed March 29, 2020.
- 142 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 143 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 144 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.
- 145 Hysingla ER [package insert]. Stamford, CT; Purdue; October 2019.
- 146 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 147 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.
- 148 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 149 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.
- 150 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 151 Kadian [package insert]. Madison, NJ; Allergan; October 2019.
- 152 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.
- 153 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 154 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.
- 155 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 156 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
-

---

157 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.

158 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.

159 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.

160 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.

161 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.

162 FDA Drug Safety Communication: FDA warns about several safety issues with opioid pain medicines; requires label changes. March 22, 2016. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm489676.htm>. Accessed March 29, 2020.

163 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.

164 Butrans [package insert]. Stamford, CT; Purdue; October 2019.

165 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.

166 Hysingla ER [package insert]. Stamford, CT, Purdue; October 2019.

167 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.

168 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.

169 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.

170 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.

171 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.

172 Kadian [package insert]. Madison, NJ; Allergan; October 2019.

173 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.

174 Embeda [package insert]. New York, NY; Pfizer; October 2019.

175 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.

176 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.

177 Opana ER [package insert]. Malvern, PA; Endo; October 2019.

178 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.

179 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.

180 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.

181 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.

182 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.

183 FDA Drug Safety Communication. FDA warns about several safety issues with opioid pain medicines; requires label changes. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm489676.htm>. Accessed March 29, 2020.

184 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.

185 Butrans [package insert]. Stamford, CT; Purdue; October 2019.

186 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.

187 Hysingla ER [package insert]. Stamford, CT, Purdue; October 2019.

188 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.

189 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.

190 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.

191 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.

192 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.

193 Kadian [package insert]. Madison, NJ; Allergan; October 2019.

194 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.

195 Embeda [package insert]. New York, NY; Pfizer; October 2019.

196 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.

197 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.

198 Opana ER [package insert]. Malvern, PA; Endo; October 2019.

199 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.

200 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.

201 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.

202 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.

203 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.

204 FDA Drug Safety Communication: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm549679.htm>. Accessed March 29, 2020.

205 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.

206 Butrans [package insert]. Stamford, CT; Purdue; October 2019.

207 Duragesic [package insert]. Titusville, NJ; Janssen; October 2019.

208 Hysingla ER [package insert]. Stamford, CT, Purdue; October 2019.

209 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.

210 Exalgo [package insert]. Webster Groves, MO; Mallinckrodt; October 2019.

211 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.

212 MS Contin [package insert]. Coventry, RI; Rhodes; October 2019.

213 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.

214 Kadian [package insert]. Madison, NJ; Allergan; October 2019.

215 Morphabond ER [package insert]. Basking Ridge, NJ; Daiichi Sankyo; October 2019.

216 Embeda [package insert]. New York, NY; Pfizer; October 2019.

217 Oxycontin [package insert]. Stamford, CT; Purdue; October 2019.

---

- 
- 218 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 219 Opana ER [package insert]. Malvern, PA; Endo; October 2019.
- 220 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 221 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 222 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 223 Tramadol HCl extended-release [package insert]. Morgantown, WV; Mylan; October 2019.
- 224 Clinical Pharmacology. Available at: [www.clinicalpharmacology.com](http://www.clinicalpharmacology.com). Accessed March 29, 2020.
- 225 Available at: <https://www.collegiumpharma.com/about/company-overview/>. Accessed March 29, 2020.
- 226 FDA Drug Safety Communication: FDA requiring color changes to Duragesic (fentanyl) pain patches to aid safety—emphasizing that accidental exposure to used patches can cause death. Available at: <https://www.fda.gov/Drugs/DrugSafety/ucm368902.htm>. Accessed March 29, 2020.
- 227 FDA identifies harm reported from sudden discontinuation of opioid pain medicines and requires label changes to guide prescribers on gradual, individualized tapering. FDA Drug Safety Communication. Available at: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-identifies-harm-reported-sudden-discontinuation-opioid-pain-medicines-and-requires-label-changes>. Accessed February 21, 2020.
- 228 Centers for Medicare and Medicaid Services. Opioid morphine equivalent conversion factors, 2018 version. Available at: <https://www.cms.gov/Medicare/Prescription-Drug-Coverage/PrescriptionDrugCovContra/Downloads/Oral-MME-CFs-vFeb-2018.pdf>. Accessed March 29, 2020.
- 229 Belbuca [package insert]. Raleigh, NC; BioDelivery Sciences; December 2019.
- 230 Zohydro ER [package insert]. Morristown, NJ; Currax; October 2019.
- 231 Hysingla ER [package insert]. Stamford, CT; Purdue; October 2019.
- 232 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 233 Embeda [package insert]. New York, NY; Pfizer; October 2019.
- 234 Nucynta ER [package insert]. Stoughton, MA; Collegium; October 2019.
- 235 Buynak R, Shapiro DY, Okamoto A, et al. Efficacy and safety of tapentadol extended release for the management of chronic low back pain: results of a prospective, randomized, double-blind, placebo- and active-controlled Phase III study. *Expert Opin Pharmacother*. 2010; 11(11): 1787–1804.
- 236 Schwartz S, Etropolski M, Shapiro DY, et al. Safety and efficacy of tapentadol ER in patients with painful diabetic peripheral neuropathy: results of a randomized-withdrawal, placebo-controlled trial. *Curr Med Res Opin*. 2011; 27(1):151–162.
- 237 Afilalo M, Etropolski MS, Kuperwasser B, et al. Efficacy and safety of tapentadol extended release compared with oxycodone controlled release for the management of moderate to severe chronic pain related to osteoarthritis of the knee: a randomized, double-blind, placebo- and active-controlled phase III study. *Clin Drug Investig*. 2010; 30(8):489–505.
- 238 Babul N, Noveck R, Chipman H, et al. Efficacy and safety of extended-release, once-daily tramadol in chronic pain: a randomized 12-week clinical trial in osteoarthritis of the knee. *J Pain Symptom Manage*. 2004; 28(1):59–71.
- 239 Tramadol HCl extended-release [package insert]. Baltimore, MD; Lupin; October 2019.
- 240 Conzip [package insert]. Bridgewater, NJ; Vertical; October 2019.
- 241 Butrans [package insert]. Stamford, CT; Purdue; October 2019.
- 242 Xtampza ER [package insert]. Cincinnati, OH; Patheon; October 2019.
- 243 Arymo ER [package insert]. Wayne, PA; Egalet; October 2019.
- 244 Webster LR, Smith MD, Lawler J, Lindhardt K, Dayno JM. Human abuse potential of an abuse-deterrent (AD), extended-release (ER) morphine product candidate (morphine-ADER injection-molded tablets) vs extended-release morphine administered intranasally in nondependent recreational opioid users. *Pain Medicine: The Official Journal of the American Academy of Pain Medicine*. 2017;18(9):1695-1705. DOI:10.1093/pm/pnw219.
- 245 Dolophine [package insert]. Eatontown, NJ; West-Ward; October 2019.
- 246 Parris WC, Johnson BW Jr., Croghan MK, et al. The use of controlled-release oxycodone for the treatment of chronic cancer pain: a randomized, double-blind study. *J Pain Symptom Manage*. 1998; 16(4):205–211.
- 247 Kaplan R, Parris WC, Citron ML, et al. Comparison of controlled-release and immediate-release oxycodone tablets in patients with cancer pain. *J Clin Oncol*. 1998; 16(10):3230–3237.
- 248 Hale ME, Dvergsten C, Gimbel J. Efficacy and safety of oxymorphone extended release in chronic low back pain: results of a randomized, double-blind, placebo- and active-controlled phase III study. *J Pain*. 2005; 6(1):21–28.
- 249 Serrie A, Lange B, Steup A. Tapentadol prolonged-release for moderate-to-severe chronic osteoarthritis knee pain: a double-blind, randomized, placebo- and oxycodone controlled release-controlled study. *Curr Med Res Opin*. 2017 Aug;33(8):1423-1432. DOI: 10.1080/03007995.2017.1335189.
- 250 Coluzzi F, Ruggeri M. Clinical and economic evaluation of tapentadol extended release and oxycodone/naloxone extended release in comparison with controlled release oxycodone in musculoskeletal pain. *Curr Med Res Opin*. 2014; 30(6): 1139–1151. DOI: 1185/03007995.2014.894501.
- 251 Chaparro LE, Furlan AD, Deshpande A, et al. Opioids compared to placebo or other treatments for chronic low-back pain. *Cochrane Database Syst Rev*. 2103; 8: CD004959. DOI: 10.1002/14651858.CD004959.pub4.
- 252 Wolff RF, Aune D, Truysers C, et al. Systematic review of efficacy and safety of buprenorphine versus fentanyl or morphine in patients with chronic moderate to severe pain. *Curr Med Res Opin*. 2012; 28(5)L 833-845. DOI: 10.1185/03007995.2012.678938.
-